



# INDONESIA SECOND NATIONAL COMMUNICATION

UNDER THE UNITED NATIONS FRAMEWORK CONVENTION ON CLIMATE CHANGE



*Climate Change Protection for Present and Future Generation*



Ministry of Environment, Republic of Indonesia





# **Indonesia Second National Communication Under The United Nations Framework Convention on Climate Change (UNFCCC)**

**Jakarta, November 2010**

**Adviser**

State Minister of Environment

**Editor in Chief**

Masnellyarti Hilman, Deputy Minister for Nature Conservation Enhancement and Environmental Degradation Control

**Coordinating Lead Authors**

Rizaldi Boer, Sulistyowati, Irsal Las, Farida Zed, Nur Masripatin, Dana A. Kartakusuma, Dadang Hilman, and Haneda Sri Mulyanto

**Lead Authors**

Retno Gumilang Dewi, Rizaldi Boer, Ari Wibowo, Suryahadi, Muhammad Ardiansyah, Agus Bueno, Rini Hidayati, Feril Hariati, Prihasto Setyanto, Elza Surmaini, Arien Heryansyah, Kiki Kartikasari, Idat G. Permana, and Akhmad Faqih

**Contributing Authors:**

Sunaryo, Marwansyah Lobo Balia, Wendy Aritenang, Eddy Effendi Tedjakusuma, Subandono Diposaptono, Ida Kusuma, Elly Andriani Sinaga, Gatot Irianto, Hendradjat N., Yana Anggadireja, Sriworo B Harijono, Ano Herwana, Wan Alkadri, Endang Supraptini, Agus Wahyudi, Ghafur Akbar Dharmaputra, Iman Bonila Sombu, Lilih Handayaningrum, Saleh Abdurrahman, Sidik Boedoyo, Soendjoto, Agus Hermawan Atmadilaga, Thomas Djamaluddin, Tania June, Yanuar J. Purwanto, Adi Rakhman, Ucock Siagian, Mala H Nasution, Puji Lestari, Syahrina D Anggraeni, Yuyun Ismawati, Ujang Solihin, Aisyah Syafe'i, Upiek SA Kamil, Sisi Febriyanti, Diva Oktavariani, Fitriyani, Doddy Juli Irawan, Gito Ginting, Jessica and Andrea.

**Acknowledgment**

The State Minister of Environment would thank Ministry of Agriculture, National Development Planning Agency, Ministry of Marine Affairs and Fisheries, Ministry of Transportation, Ministry of Agriculture, Ministry of Health, The Ministry of Industry, Ministry of Foreign Affairs, Ministry of Home Affairs, Ministry of Energy & Mineral Resources, Ministry of Research and Technology, Ministry of Public Work, Centre for Climate Risk and Opportunity Management in Southeast Asia Pacific-Bogor Agriculture University (CCROM SEAP-IPB), Centre for Research on Energy-Bandung Institute of Technology (PPE-ITB), Agency for the Assessment and Application Technology (BPPT), National Agency for Meteorology, Climatology and Geophysics (BMKG), Central Bureau of Statistic (BPS) Indonesia, National Coordinating Agency for Survey and Mapping (Bakosurtanal), National Institute of Aeronautics and Space (LAPAN) and UNDP CO Indonesia for overall support and backstopping in the preparation of the Second National Communication under the United Nations Framework Convention on Climate Change (UNFCCC)

---

# Preface



It is with great pleasure that I have availed myself of the opportunity to present a foreword for the Second National Communication (SNC) of the Republic of Indonesia to the United Nations Framework Convention on Climate Change (UNFCCC).

The devastating impact of global warming is already noticeable in Indonesia and will likely worsen due to further unsustainable economic development and human-induced climate change. Increasingly frequent and severe floods, extreme weather events and prolonged droughts in some regions will lead to further environmental destruction and degradation, human injury and illness. The continuation of warmer temperatures will also increase the number of malaria and dengue fever cases and lead to an increase in other infectious diseases as a result of poor nutrition due to food production disruption.

Indonesia is estimated to have emitted 1.378 Giga tonnes of CO<sub>2</sub> equivalent in the year 2000 with a significant proportion of emissions due to deforestation and land-use change at 1.1 million hectares per year, which accounts for 51 % of the country's annual greenhouse gas emissions. Despite the large emissions from the forestry sector, energy and other sectors contribute only 0.557 Giga tonnes of CO<sub>2</sub>-equivalent which is still below the global average.

Regulations regarding environmental protection of the Republic of Indonesia mandate the government to promote and adopt policies that maintain its environment for the benefit of present and future populations. Addressing the challenge of climate change is one of the Indonesian Government's highest priorities, which has committed to a national emission reduction target and is implementing a comprehensive response to climate change in achieving this target, including adaptation to unavoidable climate change. Through its various government agencies and in partnership with the private sector and non-governmental organizations, Indonesia seeks to develop and adopt pre-emptive and corrective actions and activities to address the predicted and actual impacts of climate change.

Indonesia's Second National Communication, to be submitted to the UNFCCC, documents recent emissions, the impacts of Climate Change, as

well as Indonesia's efforts in dealing with Climate Change, especially through mitigation and adaptation measures.

Addressing the issue of climate change in Indonesia - with a wide range of complications and territories- requires the attention of a diverse group of experts and key government and non-governmental agencies, coordinated by the Directorate of Climate Change Impact Management under the Deputy Minister of the Ministry of Environment.

This formal document submitted to the UNFCCC was only made possible with financial support from Annex I countries of UNFCCC through GEF (Global Environment Facility).

Finally, on behalf of the Government of the Republic of Indonesia, I wish to acknowledge with gratitude to all these organizations, institutions and individuals who participated in the process of the preparation of the Second National Communication. Your active participation in the deliberations was vital to the success of this project.

Thank you very much.



**Prof. Dr. Gusti Muhammad Hatta**  
Minister for the Environment

---

# Table of Contents

|   |        |
|---|--------|
| <b>EXECUTIVE SUMMARY</b>  | ix     |
| <b>LIST OF FIGURE</b>   | xxiii  |
| <b>LIST OF TABLE</b>  | xxix   |
| <b>GLOSSARY OF ABBREVIATION</b>   | xxxiii |
| <br>  |        |
| <b>CHAPTER I</b>  |        |
| <b>NATIONAL CIRCUMSTANCES</b>   | I-1    |
| 1.1 Geography and Climate   | I-1    |
| 1.2 Extreme Climate Events  | I-3    |
| 1.3 Economic and Social Development                                       | I-4    |
| 1.4 Sectoral Conditions   | I-7    |
| 1.4.1 Energy Sector   | I-6    |
| 1.4.2 Industrial Sector   | I-8    |
| 1.4.3 Forestry Sector   | I-9    |
| 1.4.4 Agriculture Sector  | I-11   |
| 1.4.5 Water Sector  | I-13   |
| 1.4.6 Coastal and Marine Sector   | I-14   |
| <br>  |        |
| <b>CHAPTER II</b>   |        |
| <b>NATIONAL GHG INVENTORY</b>   | II-1   |
| 2.1 Introduction  | II-1   |
| 2.2 Institutional Arrangements  | II-1   |
| 2.3 Description of the Process of Preparing the Inventory                 | II-2   |
| 2.4 Overview of Source and Sink Category Emission Estimates for year 2000 | II-3   |
| 2.5 Sectoral Emissions  | II-9   |
| 2.5.1 Energy  | II-9   |
| 2.5.2 Industry  | II-13  |
| 2.5.3 Agriculture   | II-14  |
| 2.5.3.1 Livestock   | II-15  |
| 2.5.3.2 Rice Cultivation  | II-18  |
| 2.5.3.3 Agricultural Soils  | II-19  |
| 2.5.3.4 Biomass Burning   | II-21  |
| 2.5.4 Land Use Change and Forestry (LUCF)                                 | II-22  |
| 2.5.5 Waste   | II-26  |
| 2.6 Emissions Trends  | II-27  |
| 2.7 Key Category Analysis   | II-28  |
| 2.8 Uncertainty Analysis  | II-29  |
| 2.9 Quality Assurance/Quality Control (QA/QC)                             | II-30  |
| 2.10 Plans for Improvement  | II-31  |

|  |       |
|--|-------|
| <b>CHAPTER III</b>   |       |
| <b>GENERAL DESCRIPTION OF STEPS TAKEN OR ENVISAGED TO IMPLEMENT THE CONVENTION</b> | III-1 |
| 3.1 Introduction   | III-1 |
| 3.2 Process of Integrating Climate Change into the National Development Plan       | III-1 |
| 3.3 Financial Institutions   | III-4 |
| <b>CHAPTER IV</b>  |       |
| <b>MEASURES TO FACILITATE ADEQUATE ADAPTATION TO CLIMATE CHANGE</b>                | IV-1  |
| 4.1 Introduction   | IV-1  |
| 4.2 Historical Climate Change  | IV-1  |
| 4.3 Future Climate Change  | IV-4  |
| 4.4 Climate Change Impacts and Adaptation Assessment                               | IV-10 |
| 4.4.1 Agriculture  | IV-10 |
| 4.4.2 Water Resources  | IV-19 |
| 4.4.3 Forestry   | IV-24 |
| 4.4.4 Coastal and Marine   | IV-29 |
| 4.4.5 Health   | IV-45 |
| <b>CHAPTER V</b>   |       |
| <b>MEASURES TO MITIGATE CLIMATE CHANGE</b>   | V-1   |
| 5.1 Energy Sector  | V-1   |
| 5.1.1 Policy Framework   | V-1   |
| 5.1.2 Emissions Projections and Mitigation Measures                                | V-4   |
| 5.1.3 Mitigation Costs   | V-14  |
| 5.2 Industry Sector  | V-18  |
| 5.2.1 Policies Framework   | V-18  |
| 5.2.2 Emissions Projections and Mitigation Measures                                | V-18  |
| 5.3 Forestry Sector  | V-20  |
| 5.3.1 Policy Framework   | V-20  |
| 5.3.2 Emissions Projections and Mitigation Measures                                | V-22  |
| 5.3.3 Mitigation Costs   | V-25  |
| 5.4 Peatland   | V-26  |
| 5.4.1 Policy Framework   | V-26  |
| 5.4.2 Emissions Projections  | V-27  |
| 5.5 Agriculture Sector   | V-27  |
| 5.5.1 Policy Framework   | V-27  |
| 5.5.2 Emissions Projections and Mitigation Measures                                | V-28  |
| 5.6 Waste  | V-32  |



**CHAPTER VI****OTHER INFORMATION**

|     |  |      |
|-----|--|------|
| 6.1 | Transfer of Technology   | VI-1 |
| 6.2 | Research   | VI-2 |
| 6.3 | Systematic Observations  | VI-3 |
| 6.4 | Information on Education, Training, Public Awareness and Capacity Building | VI-6 |
| 6.5 | Efforts to Promote Information Sharing                                     | VI-7 |

**CHAPTER VII****OBSTACLES, SHORTAGES AND NEEDS RELATED TO FUNDING,  
TECHNOLOGY AND CAPACITY**

|     |   |       |
|-----|---|-------|
| 7.1 | Climate Change Funding                                  | VII-1 |
| 7.2 | Technology  | VII-2 |
| 7.3 | Awareness, Education, Empowerment and Capacity Building | VII-3 |

**REFERENCES**



---

# Executive Summary

Indonesia continues its efforts and actions towards the implementation of the commitments as a Non-Annex I Party to the United Nations Framework Convention on Climate Change (UNFCCC). Indonesia presented its First National Communication to the UNFCCC in 1999. One of the most important parts of the content was the first National Greenhouse Gases Emissions Inventory (NGHGEI) for the year 1990 and also the results of the first studies on the country's vulnerability to climate change. The Second National Communication (SNC) presents National Greenhouse Gases Emissions Inventory (NGHGEI) for the years 2000 to 2005.

This Communication was supported by the Global Environmental Fund (GEF) through the United Nations Development Programme (UNDP), along with further funding from Government of Indonesia. The funds received allowed the improvement of the emissions inventory, and supported the development of studies on climate change mitigation and adaptation in Indonesia. On this occasion, the process of planning the National Communication content included consultations with academics and representatives from government institutions, private sector and non-governmental organizations, in order to capture their opinion and points of view about what elements of the previous communication should be improved in this second assessment. A public consultation was also held for the same purpose.

As requested, Indonesia's Second National Communication was prepared in accordance with the UNFCCC Reporting guidelines on National Communications.

## National Circumstances

Indonesia is located between 6°08' North and 11°15' South latitude, and from 94°45' to 141°05' East longitude. The country covers 1,910,931 Km<sup>2</sup>. It has five large islands (Sumatra, Java, Kalimantan, Sulawesi and Irian Jaya) and about 13,667 small islands, of which over half (56%) of which are nameless, and only 7% permanently inhabited (Figure 1-1). Extensive coastal plains and mountain areas of 1,000 metres above sea level are characteristics of Sumatra, Kalimantan and Irian Jaya. Of the 200 million ha of land territory, about 50 million ha is devoted to various agricultural activities. There is nearly 20 million ha of arable land, of which about 40% is wetland (rice fields), 40% is dry land, and 15% is shifting cultivation. Since 2005, the Republic of Indonesia has been divided administratively into 33 provinces (BPS, 2008)

The country's population is the fourth most populous nation in the world, following China, India and the United States. The population grew from 119 million in 1971 to 219 million in

2005. While the growth rate is slowing down from 1.49% (1990–2000) to become 1.34% (2001–2005), it is projected that Indonesia's population will reach 300 million in 2030.

Poverty remains a challenge, while unemployment and underemployment are still relatively high. In the Mid-Term National Development Plan (RPJMN) 2004–2009, it was targeted that poverty and unemployment would be about 5.1% and 8.2% respectively. Recent report from BPS stated that the number of unemployment by February 2008 reached 8.46% or about 9.43 million people. However, based on the projection of the Institute for Development Economics and Finance (Indef), due to current economic crisis the number of poor and unemployed in 2008 may increase to 9.5% and 16.3% respectively (Depsos, 2009).

Indonesia's GDP is approximately US\$ 175 billion, in which trade (16.7%), manufacturing (28%), agriculture (15.4%), and services (10.17%) are the main contributors. Earnings from exports were about US\$ 69 billion (primarily oil and gas, textiles, appliances, coal, copper), while imports accounted for about US\$ 44.8 billions (primarily food, chemicals, capitals and consumer goods). GDP growth has increased steadily since 1998, reaching 6.3% in 2007 thus bringing per capita GDP to around US\$ 2000.

The role of sectors other than oil and gas has become much more significant in the Indonesian economy since Asian Financial Crisis (AFC), with the oil and gas sectors experiencing negative growth rates each year since with the exception of 2000. Among the non-oil and gas sectors, agriculture has continued to play an important role. It is a sector that was significantly less affected by the AFC, and indeed helped the recovery of Indonesian economy after the crisis through a substantial increase in exports and its absorption of unemployed workers in its role as an "employer of last resort" for many jobseekers (Siregar, 2008).

## National GHG inventory

The National Greenhouse Gases Inventory (NGHGI) was estimated using Tier 1 and Tier 2 of the 2006 IPCC Reporting Guidelines and the IPCC GPG for LULUCF. In 2000, total GHG emissions for the three main greenhouse gases (CO<sub>2</sub>, CH<sub>4</sub>, N<sub>2</sub>O, PFC) without LULUCF (LUCF and peat fires) reached **556,728.78** Gg CO<sub>2</sub>e. With the inclusion of LULUCF, total net GHG emissions from Indonesia increase significantly to about 1,377,982.95 Gg CO<sub>2</sub>e (Table 1a and 1b). The GHG emissions (in CO<sub>2</sub> equivalent) were distributed unevenly between the three gases recorded, i.e. net CO<sub>2</sub> totalled 1,112,878.82 Gg, representing 80.8% of the National GHG emissions; methane (CH<sub>4</sub>) totalled 236,617.97 Gg (CO<sub>2</sub>e) or 17.2%; and nitrous oxide (N<sub>2</sub>O) totalled 28,341.02 Gg (CO<sub>2</sub>e) or 2.0%. The main contributing sectors were Land Use Change and Forestry, followed by energy, peat fire related emissions, waste, agricultural and industry.

**Table 1a. Summary of 2000 GHG emission and removal (in Gg CO<sub>2</sub>e)**

|                    | CO <sub>2</sub> | CH <sub>4</sub> | N <sub>2</sub> O | PFC    | Total        |
|--------------------|-----------------|-----------------|------------------|--------|--------------|
| Energy             | 247,522.25      | 30,174.69       | 3,240.64         | NO     | 280,937.58   |
| Industrial Process | 40,342.41       | 2,422.73        | 133.22           | 145.15 | 43,043.52    |
| Agriculture        | 2,178.30        | 50,800.18       | 22,441.25        | NO     | 75,419.73    |
| LUCF <sup>1</sup>  | 821,173.35      | 56.35           | 24.47            | NO     | 821,254.17   |
| Waste              | 1,662.49        | 153,164.02      | 2,501.45         | NO     | 157,327.96   |
| Total              | 1,112,878.82    | 236,617.97      | 28,341.02        | 145.15 | 1,377,982.95 |

<sup>1</sup>Note: Emission from peat fire was included

**Table 1b. Summary of GHG emissions from 2000-2005 from all sectors (in Gg CO<sub>2</sub>e)**

| Source                         | 2000         | 2001         | 2002         | 2003         | 2004         | 2005         |
|--------------------------------|--------------|--------------|--------------|--------------|--------------|--------------|
| Energy                         | 280,937.58   | 306,774.25   | 327,910.62   | 333,950.21   | 372,123.28   | 369,799.88   |
| Industrial Process             | 43,043.52    | 49,810.15    | 43,716.26    | 47,901.63    | 47,985.20    | 48,733.38    |
| Agriculture                    | 75,419.73    | 77,500.80    | 77,029.94    | 79,828.80    | 77,862.54    | 80,179.31    |
| LUCF                           | 649,254.17   | 560,546.00   | 1,287,494.79 | 345,489.33   | 617,423.23   | 674,828.00   |
| Peat Fire                      | 172,000.00   | 194,000.00   | 678,000.00   | 246,000.00   | 440,000.00   | 451,000.00   |
| Waste                          | 157,327.96   | 160,817.76   | 162,800.37   | 164,073.89   | 165,798.82   | 166,831.32   |
| Total With LUCF & peat fire    | 1,377,982.95 | 1,349,448.96 | 2,576,951.98 | 1,217,243.86 | 1,721,193.07 | 1,791,371.89 |
| Total Without LUCF & peat fire | 556,728.78   | 594,902.96   | 611,457.19   | 625,754.53   | 663,769.84   | 665,543.89   |

<sup>1</sup>Note: Emission from peat fire was taken from van der Werf et al (2008). <sup>2</sup>Estimated based from MoF (2009) and Bappenas (2009a),

<sup>3</sup>Emission 2001 was not included in determining the trend

The emission estimates in the SNC are lower than those reported by a 2007 PEACE, World Bank and DFID study, which suggested Indonesia to be the 3rd largest emitter country. The study estimated that the total emission from Indonesia was about 3,014,000 Gg CO<sub>2</sub> where LUCF contributed about 85% or about 2,563,000 Gg CO<sub>2</sub> (twice the SNC estimate above). Indeed, a further study from the World Bank (2008) suggested that the mean annual CO<sub>2</sub> emissions from LUCF reached up to 2,398,000 Gg, assuming 53% from peat fire, 20% from peat drainage (peat oxidation), 22% from deforestation and 5% from palm oil and timber plantation establishment. Between the SNC and World Bank estimates, a recent study from DNPI (2009) suggested that the total GHG emissions from LUCF in 2005 reached 1,880,000 Gg CO<sub>2</sub>e where about 55% was from peat emissions. These large differences in Indonesian emissions estimates thus appear to be mainly due to differences in estimates of LUCF emissions, particularly from peat.

Inter-annual variation of emissions from peat fires is also very high. High emissions normally occurred in El Nino years (1997, 2002, and 2006). The highest estimate was from Hooijer *et al* (2006), which estimated the emission based on a Borneo hot-spot count and a carbon calculation method used by Page *et al.* (2002), who estimated emissions from peat fires in 1997 El-Nino. This approach may give an overestimate as hotspot counts in peat lands are not fully proportional to CO<sub>2</sub> emissions, which are governed by further factors such as the depth and area of burning. Therefore, this relationship may not also be generally applicable for the

whole of Indonesia, and so the extrapolation of emission estimates over Indonesia based on limited ground checks in Kalimantan may lead to overestimation.

Van der Werf *et al.* (2007) attempted to improve emission estimates from peat fires using several sources of satellite data with biogeochemical and atmospheric modeling to better understand and constrain peat fire emissions from Indonesia. This resulted in far lower estimates. The NGHGI adopted the study of the Van der Werf *et al.* (2007). In 2000, their estimated emissions from peat fires were calculated at about 172,000 Gg CO<sub>2</sub>e, while the average emission of peat fire (2000-2006) was about 466,000 Gg CO<sub>2</sub>e.

Indonesia plans to improve emission estimates, particularly from LULUCF and peat lands in the next National GHG Inventory. At present, activities undertaken by the Ministry of Forestry and Ministry of Agriculture to improve emission estimates from peat lands are the following:


- a. The Ministry of Agriculture (through the National Research Consortium for Climate Variability and Climate Change) is conducting studies to develop emissions factors from peat lands under different usage scenarios in Central Kalimantan. The Ministry of Agriculture will also expand this study to other provinces and request support from international agencies. Additional surveys to improve data on peat depth (particularly in Papua) are also being planned. It is expected that the funding allocated for the 3<sup>rd</sup> National Communication can provide additional support for the studies.
- b. The Ministry of Forestry is improving the emission sink factors from forests and emission factors from fire (both in mineral soils and peat land). The programme is being undertaken through the INCAS (Indonesian National Carbon Accounting System) and other relevant research programmes under the MoF and partners.
- c. The state Ministry of Environment is conducting a pilot study on Peatland Management, including calculation of GHG emissions from peat lands in West Kalimantan and Riau Province.
- d. A detailed analysis commissioned by Indonesia's National Development Planning Agency (BAPPENAS) is also commissioning a detail analysis on peat land emission and projection as well as mitigation strategies. The analysis is undertaken by a multi-disciplinary team of Indonesian scientists, economists and legal specialists.

## Steps planned to implement the Convention

To effectively implement the UNFCCC, the Government of Indonesia has made a number of significant steps in mainstreaming climate change issues with other national development priorities. The first of these was the issuance of the National Action Plan on Climate Change (MoE, 2007a), which describes appropriate actions to reduce GHG emissions and adaptation activities in Indonesia. This was then followed by a further document, "National Development Planning: Indonesia responses to climate change", called as Yellow Book. The National Development Planning Agency (Bappenas) subsequently developed a Climate Change Sectoral

Roadmap (ICCSR), meant to bridge the National Action Plan on Climate Change into the 5 year Mid-Term Development Plan (RPJM) 2010-2014, and to provide inputs for the subsequent RPJMN until 2030. The process of developing the Climate Change Roadmap is shown in Figure 1.

To support and accelerate the implementation of climate change programs, the Government of Indonesia established a number of innovative ways to link international financial resources with national investment strategies. One new source of funds to support the implementation of climate change programs in Indonesia is the Indonesian Climate Change Trust Fund (ICCTF), described briefly in the following section. This Indonesian Climate Change Trust Fund (ICCTF; Bappenas, 2009b) aims to be a showcase of alternative financing for climate change mitigation and adaptation programs. At this stage, the ICCTF has five specific objectives namely (i) to facilitate and accelerate investment in renewable energy and efficiency and simultaneously reduce GHG emissions from the energy sector, (ii) to reduce emission from deforestation and forest degradation and stabilizing carbon stock through sustainable forest and peat land management, (iii) to reduce vulnerability in coastal zones, agriculture and water sectors, (iv) to bridge the financial gaps necessary to address climate change mitigation and adaptation and (v) to increase the effectiveness and impact of external finance for climate change programs.



**Figure 1. Process of Development of Climate Change Sectoral Roadmap (ICCSR)**

Following commitment of the GOI to actively participate in reducing its emissions through National Appropriate Mitigation Action (NAMA), in 2009, the President of the Republic of Indonesia at the G-20 meeting in Pittsburgh and at COP15 in Copenhagen committed to an ambitious, world-leading target of 26% reduction in carbon emissions from Business As Usual (BAU) by 2020. Further emissions reductions of 41% are expected with international support. With this commitment, Indonesia will follow a low carbon development path. As such, BAPPENAS is coordinating the sectors to develop the National Action Plan for Reducing Greenhouse Gas Emissions (RAN-GRK). The GOI lists three principles in the RAN-GRK, stating that

the mitigation actions (i) should not hinder economic growth, and should prioritize people’s welfare, especially in with regard to energy resilience and food security, (ii) support protection of poor and vulnerable communities, including environment conservation in the framework of sustainable development and (iii) consist of core activities to reduce emissions and supporting activities to strengthen the policy framework. Figure 2 shows the interconnection between policies, guidelines, roadmaps and investment funds towards low carbon development.




Figure 2. Process of mainstreaming climate change into national agenda: Policies, guidelines, roadmaps & investment funds (BAPPENAS, 2009b). Note: M is mitigation and A is adaptation.

## Measures to Facilitate Adequate Adaptation to Climate Change

Due to its geographical location, topography and socioeconomic aspects, Indonesia is vulnerable to the impacts of climate variability and climate change. The El Niño and La Niña phenomena (ENSO), as well as extreme meteorological conditions, have historically resulted in serious damage that affects a wide range of different socioeconomic sectors. A number of studies suggested that El Niño events have become more frequent as the global temperature anomalies associated with each El Niño continue to increase (Hansen *et al.*, 2006; Timmerman *et al.* 1999). This suggests that the increasingly high temperatures are exacerbating the extreme regional weather and climate anomalies associated with El Niño.

By assessing historical natural hazard data from 1907-2007 (OFDA/CRED International Disaster Database 2007), it is clear that the first climate-natural hazard categorized as global hazard occurred for the first time in early 1950s; by the 1980s they were occurring more frequently. Over this entire record, the most frequent hazard has been flooding, followed by landslides and water or vector borne diseases, wind storms, forest fires, drought, and high tide/storm surge.



Furthermore, the top 20 natural hazards, causing huge economic loss and adverse human impacts, mostly occurred after the 1980s, suggesting increasing trend in hazards' intensity (Boer and Perdinan, 2008).

With the increasing trend of climate hazards frequency and intensity, the most affected group will be the poorest sections of society, which are most exposed and least resilient. The capacity of this group to adapt to the extreme climate events and future climate change is limited due by their limited resource availability and access to climate information and technologies. It is likely, therefore, that their reliance on national and local public assistance from government will increase in the future. During the 2006/07 El-Nino, many farmers in the Timor district of East Nusa Tenggara Province—one of the most vulnerable districts to drought—experienced crop failure due to drought. As a consequence, the major source of income for the poorest came from government aid (*'bantuan pemerintah'*). In addition to aid, many of farmers often have to sell their domestic livestock (*ternak*) or engage in informal laboring to generate additional income. At Indramayu, the drought occurrence associated with the 2002/03 El-Nino caused huge rice production loss. In 2003, the number of household that could not meet their food basic needs increased by 14% compare to the normal years (Boer *et al.*, 2006).

It is clear, therefore, that improving access to a diversified set of incomes and resources is a key method for improving climate resilience. This is closely linked with rapid poverty reduction, which is essential to help poor and vulnerable communities improve their resilience both to natural climate variability and to the greater stresses of human-induced climate change in the future.


*Observed Climatic Changes and Predictions.* Analysis of long historical climate data suggests that maximum and minimum temperature have increased consistently (MoE, 2007). Significant decreases and/or increases in rainfall have also detected in many part of Indonesian region, with different significant trends in different areas. Based on data over 300 stations, with length of records between 20 and 50 years, a significantly decreasing trend in Dec-Jan rainfall was observed in small part of Java and Papua, and Sumatra and large part of Kalimantan islands, whereas significant increasing trend were observed in most of Java and Eastern Indonesia such as Bali, NTB and NTT (Figure 3). For Jun-Aug rainfall, significantly decreasing trends were observed in most of Indonesian region with exception in Pandeglang (West Java), Makasar (South Sulawesi), Monokwari, Sorong (Irian Jaya) and Maluku (Figure 3).

Monsoon onset has also changed in many part of Indonesia. Based on analysis of data from 92 stations spread over Indonesia, monsoon onset has increasingly been delayed in some part of Indonesia, particularly in Java; similarly, the length of the wet season has tended to shorten, particularly in South Sumatra, Java and Kalimantan. Other studies conducted in East Java also suggest that the number of extreme dry months in the Brantas Catchment area has increased in the last five decades, particularly in areas near to the coast (Aldrian and Djamil 2006). In such coastal areas, the number of extreme dry months increased to 4 months in the last ten years and in 2002 it reached 8 months, a level considered as the longest dry season for the whole five

decades. In mountain areas, the number of dry months is about 1-2 months over the last ten years, with maximum number of 4 months.


Using 14 General Circulation Models (GCMs), under increasing GHG emission scenarios (SRESA2), most of models are in agreement that in 2025 the wet seasonal rainfall (DJF) in Java, Bali, NTB, NTT and Papua will increase, while in other parts will decrease. By 2050 and 2080, most of Indonesian regions will experience higher rainfall than current condition, with exception in northern part of Sumatra and Kalimantan (Figure 4). Furthermore, dry season rainfall in most parts of Java might decrease in 2025, increase again in 2050, and then decrease in 2080, particularly in West Java and South Sumatra (Figure 4). Under low emission scenarios (SRESB1), the pattern of change is similar to that of high emission scenarios but the magnitude of change is slightly lower.

The monsoon onset more generally in Java and Bali may also delay under warming atmosphere. It is clear that a 30-day delay in monsoon onset is very likely to occur more frequently in 2050 than it does today and the length of the rainy season will shorten (Naylor *et al.*, 2007).



**Figure 3. Trend of seasonal rainfall for Dec-Feb (above) and June-Aug (below) in Indonesia (Source: Boer *et al.*, 2009a)**

Global warming is also very likely to increase sea levels. Historical data shows increasing trends in mean sea level (MSL) in a number of locations. However, the rate of increase varies with locations (Sofian, 2009). The relative sea level rise accelerates significantly in areas where coastal erosion is removing material and where the land border has been subsiding. The increase in MSL has also increased the problem of saltwater intrusion and salinity.



**Figure 4. Trend of wet seasonal (December-February) and dry seasonal (June-August) rainfall under high emission scenarios (SRESA2). Note: Dark Red (indicator 1) means that all GCM model are in agreement to suggest the seasonal rainfall will decrease and dark blue (Indicator 0) means that all models are in agreement to suggest the seasonal rainfall will increase (Boer *et al.*, 2009a).**

*Sector-Wise impacts and Adaptation Measures.* Changes in spatial rainfall patterns, the length of the wet season and inter-seasonal variability will have serious implications for many sectors. In the agriculture sector, the current cropping pattern may no longer be the most effective food production system. At present, the pattern used in most of the rice growing areas of Indonesia is rice-rice. The second planting depends heavily on irrigation water. Under extreme drought years, the availability of irrigation water is very limited, usually leading to major rice production losses. Under a changing climate, the occurrence of extreme climate events (drought) will be more frequent than the current climate and there is a possibility that the dry season will persist for longer periods. Therefore, keeping this cropping pattern in the future may expose Indonesian farmers to more frequent crop failures. Thus, in areas where the pattern of rainfall changes in this direction, farmers should consider alter their cropping pattern from rice-rice to rice-non rice. If the rice-rice pattern is maintained, improvement of water storage and irrigation facilities will be required for compensating the decreased in JJA rainfall. More efforts to create new short maturing rice varieties should also be in place to anticipate the shorter wet season. The increase in temperature and the changes in rainfall pattern and length of seasons may also trigger the development of crop pests and diseases. Field observations in a number Java districts, such as Indramayu, Magelang, Semarang, Boyolali, Kulonprogo, and Ciamis, provide the evidence of this phenomena (Wiyono, 2007).

Furthermore, analysis of climate change impacts in rice production in Java suggested that production between 2025 and 2050 is likely to decrease by about 1.8 and 3.6 million ton from the current production level respectively (Boer *et al.*, 2009b). By including the impact of rice field conversion to non-agriculture lands in Java, i.e. taking 0.77% of land out of production per annum, the production loss in 2025 and 2050 will increase to 5.2 and 13.0 million ton respectively. Thus, the impact of rice field conversion will be much higher than the impact of climate change. To compensate this loss, new rice areas of around 1.5 and 3.5 million ha will be needed in 2025 and 2050 respectively outside Java. Alternatively, the loss can be compensated by increasing rice productivity by about 20% and 50% from the current levels. To achieve this, breeding technology would have to be able to increase rice productivity by about 1 t/ha and 2.5 t/ha from the current productivity (~5 t/ha).

A rapid assessment conducted by Parry *et al.* (1992) in a number of locations in Indonesia suggested that sea level rise due to global warming will also reduce local rice supply in Krawang and Subang districts by about 300,000 tonnes. Similarly, maize output would likely be reduced by 10,000 tons--about half of this due to inundation. Sea-level rise would also be likely to affect fish and prawn production. The loss is estimated at over 7000 tonnes and 4000 tons respectively (valued at over US\$0.5m). In the lower Citarum Basin sea-level rise could result in the inundation of about 26,000 ha of ponds and 10,000 ha of crop land. This could result in the loss of 15 000 tons of fish, shrimp and prawns and about 940 000 ton of rice. Parry *et al.* stated that the socio-economic implications of this transition in Subang District alone could be the loss of employment for about 43 000 farm laborers. In addition more than 81 000 farmers would have to look for other sources of income due to the inundation of their rice fields or prawn and fish farms due to sea-level rise.

For water resources, the present condition of the water balance suggests that in most of Java and the eastern islands of Indonesia is already in deficit for most of year. With such conditions, increasing planting in these islands is not possible, further restricting options for increasing rice production outlined above. Under changing a climate, more districts will have water scarcity problems (Herienschah *et al.*, 2009). A key need is the development of new initiatives to anticipate the scarcity of water due to climate change and increases on water demand, especially in urban areas where populations are increasing and industrial activities are taking place. Inter-basin transfer of water may be one of the potential options to anticipate the scarcity of water in the future. In Indonesia many basins have surplus water resources, even in the ultimate stage of development, while others face serious shortages, especially during extreme drought years. Creation of storages and inter-basin transfer of water from surplus to deficit regions (such as in West Nusa Tenggara) could therefore be an option for achieving more equitable distribution of resources and their optimal utilization.

For coastal and Outer Islands, the increase in sea level by about 25 to 50 cm in 2050 and 2100 as projected by many models will inundated many parts of the coastal cities of Indonesia. Land subsidence will exacerbate this, increasing the total area that will be inundated permanently. Between 25% and 50% of area in a number of sub-districts in coastal cities such as Semarang,

Surabaya, Jakarta and Medan will be under water permanently. The increase of sea level rise may also inundate the outer islands of the country, and this will affect the area of Indonesian territory. The analysis suggests that an increase of sea level of up to 50 cm will not inundate the outer islands of Indonesia permanently. However, in combination with tidal patterns in the region, about five outer islands will temporarily inundate. These islands include Alor (next to Timor Leste), Pelampong (next to Singapura), Senua (next to Malaysia), Simuk and Sinyaunyau (next to India).

The increase in sea temperature will also caused serious problems for the coral ecosystems. Wetland International (Burke *et al.*, 2002) reported that the 1997 El-Niño damaged about 18% of the coral ecosystems in South East Asia. In Indonesia, coral bleaching was observed in many places such as in the eastern part of Sumatra, Java, Bali, and Lombok. In 'thousands islands' (north of the Jakarta coast), about 90-95% of the corals located 25 m below sea surface have been bleached.

For forests sector, the decreasing dry season rainfall and shortening length of wet season will increase the risk of forest fire. Two islands which are very prone to fire are Sumatra and Kalimantan. Based on hotspot density patterns, two provinces that have very high hot spot density are Riau Province and Central Kalimantan. Hotspot densities in these two islands increased rapidly when dry season rainfall decreases or length of dry season extends, particularly during El Nino years. It was revealed that the hot spot density increase rapidly as the monthly rainfall in dry season by more than 50 mm below normal (Ardiansyah and Boer, 2010).


For health, extreme weather related to ENSO also contributes to the outbreak of human diseases such as malaria, dengue, diarrhea, cholera and other vector borne diseases. Dengue cases were found to increase significantly in La-Niña years when seasonal rainfall levels increased above average. A significant upward trend in the number of dengue cases was also observed in Java, especially in large cities. In the future, under changing climate it is likely that the risk of transmission of dengue and malaria may well increase (Hidayati *et al.*, 2009).

All sectors have developed a number of measures to increase the capacity to address and to increase resilience to current and future climate risks respectively. The programs are described in the Yellow Book and the sectoral roadmap to address climate change (Bappenas, 2009a and 2009 b).

## Measures to Mitigate Climate Change

The impact of implementation of mitigation options on the level of GHG emission is evaluated by comparing the emission level under Business as Usual (BAU) with emission level under two mitigation scenarios. Based on the emissions projection analysis, net emissions under BAU in 2020 are expected to be approximately 2.95 Gt CO<sub>2</sub> – more than double the emissions in year 2000. It was found that by 2020, the rate of net emissions under mitigation scenario 1 reached

about 31% of that of the BAU and under mitigation scenario 2, it could reduce up to about 48% (Figure 5).



**Figure 5. Projection of emissions of all sectors under BAU and mitigation scenarios (MoE, 2009). Note: Contribution of transportation sector to the total energy emission is approximately 20% (based on emissions data from 2000-2006).**

In energy sector, reducing of the emission from the BAU will be done through additional efforts related to energy conservation and new and renewable energy development (i.e. enhance geothermal program, micro-hydro, biofuel, biomass waste to energy, solar PV, wind energy and coal bed methane [CBM]). In industry sector, emission reduction will be done by increasing the efficiency in production processes, introduction of new technologies, or by changing the raw materials (i.e., using waste as an alternative material in the cement industry). The efforts can be achieved through the clean development mechanism (CDM) scheme and private sector participation. Public funding is still needed for the dissemination of programs related to CO<sub>2</sub> emissions reductions in industrial processes.

In Agriculture sector, the key mitigation program will be through (i) implementing no-burning technology for land clearing, specifically in the horticulture and agriculture plantation sub-sectors, (ii) developing a Fire Early Warning System particularly in peat areas, (iii) introducing low methane emitting rice varieties, (iv) increasing the use of agriculture waste for bio-energy and composting, (v) introducing biogas technology, (vi) improving feeding quality and supplementation for livestock and (vi) developing new agriculture areas which will be directed to unproductive land such as grassland and abandoned land through revitalization of agriculture spatial plan program. In forestry sector, the mitigation measures will be done through (i) combating illegal logging and its associated trade, (ii) revitalizing forestry

industries, (iii) conserving and rehabilitating forest resources, (iv) empowering the economy of the community within and surrounding the forest area, and (v) stabilizing forest area for promoting and strengthening sustainable forest management. In waste sector, the emission reduction from solid waste will be done through converting the open dumping practices to a more managed solid waste handling system (i.e., a sanitary landfill equipped with gas flaring or utilization systems), and handling liquid domestic waste using sewerage systems.

Following commitment of the GOI to actively participate in reducing its emissions through National Appropriate Mitigation Action (NAMA), in 2009, the President of the Republic of Indonesia at the G-20 meeting in Pittsburgh and at COP15 in Copenhagen committed to an ambitious, world-leading target of 26% voluntary reduction in carbon emissions from Business As Usual (BAU) by 2020. Further emissions reductions of 41% are expected with international support. With this commitment, Indonesia will follow a low carbon development path. As such, BAPPENAS is coordinating the sectors to develop the National Action Plan for Reducing Greenhouse Gas Emissions (RAN-GRK). The GOI lists three principles in the RAN-GRK, stating that the mitigation actions (i) should not hinder economic growth, and should prioritize people's welfare, especially in with regard to energy resilience and food security, (ii) support protection of poor and vulnerable communities, including environment conservation in the framework of sustainable development and (iii) consist of core activities to reduce emissions and supporting activities to strengthen the policy framework

## Other Information

Other information considered relevant to the achievement of the objectives of the Convention includes (i) technology transfer, (ii) Research, (iii) systematic observations, (iv) information on education, training, public awareness and capacity-building and (v) efforts to promote information sharing. The role of NGOs in the implementation of adaptation and mitigation activities with local communities is quite significant. In general, NGOs play their roles as implementer and facilitator of CDM projects as well as voluntary carbon projects.

To accelerate the development of mitigation and adaptation technologies, the Government of Indonesia under the coordination of the Ministry of Agriculture has established the Research Consortium on Climate Variability and Climate Change. To monitor climate and sea level, the Government of Indonesia has installed rainfall stations in all Indonesia regions, a number of automatic weather stations (AWS) and radar in a number of places particularly in areas prone to weather/climate hazards, one Global Atmospheric Watch (GAW) in Bukittinggi, West Sumatra as part of the global atmospheric observation under World Weather Watch.

In forestry sector, development of systematic forest and land use monitoring systems is also crucial to support the implementation of programs for reducing emission from deforestation and forest degradation. The Government of Indonesia is in the process of establishing Indonesia's National Carbon Accounting System (INCAS), based on Australia's system but tailored to

Indonesia's (INCAS) unique circumstances. In addition, Indonesia is developing Indonesia's Forest Resource Information System (FRIS), a comprehensive and transparent information management system to support effective planning and forest management decision making for forest lands in Indonesia.

Capacity building activities related to climate change were also quite significant but many of the activities were conducted at national level. To promote information sharing related to climate, many agencies have established a Web-based climate information system.

### **Barriers, and Related Financial, Technical and Capacity Needs**

As developing country, Indonesia has several limitations in funding in the frame of climate change anticipation as well as developing and disseminating environmentally friendly technology. Therefore, the problems of technology transfer and funding are two issues that should receive support from international community. Additional important activities, which will ensure success of mitigation and adaptation program, are awareness, education, empowerment and capacity building.

To meet financial need for addressing climate change problem, the GOI Indonesia has created various funding schemes, from domestic sources as well as from bilateral and multilateral sources. One of important funding sources is Indonesian Climate Change Trust Fund (ICCTF). At the initial phase, the ICCTF will be created as an "Innovative Fund," which involves grant funding from development partners that will help overcome barriers for early program deployment. At the later stages, the ICCTF may advance by establishing a "Transformation Fund" mechanism, which would involve all available funding (i.e., public-private partnerships, loan and world capital market sources)

The immediate need to mitigation technology for Indonesia is the energy efficient technology particularly for sectors that consume energy such as power plants, industry, transportation, as well as household and commercial sectors. Real time monitoring and the use of sophisticated fire fighting technology and peat management is also required to control emission from forest/peat fire and peat decomposition. To increase the resilience to current and future climate risk, the development and implementation of a comprehensive communications strategy to increase the capacity in using climate information is also needed as well as institutionalizing the use of climate information. The development and promotion of tools for adaptation planning tailored to user's requirements that include (i) decision-support tools such as methods for assessing the costs and benefits of adaptation strategies, and guides for risk management, (ii) methods for understanding social impacts, and (iii) a national 'one stop shop' website where decision-makers and their advisers can access information about climate projections, likely climate change impacts, tools, guides and approaches to adaptation planning will also be required.



# List of Figure

|             |   |       |
|-------------|---|-------|
| Figure 1.1  | Map and overview of Indonesia, 2007   | I-1   |
| Figure 1.2  | Percent decrease in seasonal rainfall from normal during two strong El Niño years, 1997 and 1982 (Irawan, 2002)   | I-3   |
| Figure 1.3  | Number of climate hazards by type (left) and by year (right).<br>Source: Based on data from OFDA/CRED International Disaster Database (2007)  | I-4   |
| Figure 1.4  | Final energy consumption in Indonesia, 2000 – 2007 [MEMR, 2008]   | I-7   |
| Figure 1.5  | Fraction of forest loss in forest area and non-forest area  | I-10  |
| Figure 1.6  | Rate of deforestation in Indonesia (MoF, 2007)  | I-11  |
| Figure 1.7  | Development of the main agricultural systems from 1986 to 2006<br>(Data taken from Central Bureau of Statistics)  | I-12  |
| Figure 1.8  | Development of the main plantation areas from 1986 to 2006<br>(Data taken from Central Bureau of Statistics)  | I-13  |
| Figure 2.1  | Structure of the institutional arrangements for developing<br>NGHGI 2000-2005   | II-2  |
| Figure 2.2  | National emissions contributions by sector  | II-7  |
| Figure 2.3  | Estimate of emissions from peat fires from various studies. Note:<br>Emissions estimates from years beyond the publication date of the reports were made by Aldrian (2008) and Wibowo and Boer (2009) | II-8  |
| Figure 2.4  | CO <sub>2</sub> emissions by sectors in 2000  | II-8  |
| Figure 2.5  | Methane and nitrous oxide emissions in 2000 by sector   | II-9  |
| Figure 2.6  | Source of GHG emissions from fuel combustion activities   | II-10 |
| Figure 2.7  | Source categories of fugitive emissions from fuels  | II-11 |
| Figure 2.8  | CO <sub>2</sub> emissions estimates from the energy sector using reference<br>and sectoral approaches   | II-11 |
| Figure 2.9  | CO <sub>2</sub> emissions from the energy sector by source category   | II-12 |
| Figure 2.10 | Annual Growth rate of CO <sub>2</sub> emission and GDP of main emitter<br>countries (World Bank)  | II-13 |
| Figure 2.11 | GHG emissions estimates from production process activities in<br>the industrial sector  | II-14 |
| Figure 2.12 | GHG emissions from the agriculture sector from 2000-2005 by<br>source category  | II-15 |
| Figure 2.13 | Distribution of methane emissions from enteric fermentation<br>(Gg CH <sub>4</sub> ) by species type  | II-16 |
| Figure 2.14 | Methane emissions from livestock in Indonesia from 2000-2006  | II-17 |
| Figure 2.15 | Methane emissions from manure management (Gg CH <sub>4</sub> )  | II-17 |
| Figure 2.16 | Methane emissions from rice from 2000-2005.   | II-18 |

|             |   |       |
|-------------|---|-------|
| Figure 2.17 | CO <sub>2</sub> emissions from liming in agriculture.   | II-19 |
| Figure 2.18 | CO <sub>2</sub> emissions from urea fertilization from 2000-2005  | II-19 |
| Figure 2.19 | Direct N <sub>2</sub> O emissions from N synthetic fertilizer and manure management applied to managed soil and flooded rice  | II-20 |
| Figure 2.20 | Indirect N <sub>2</sub> O emissions and projections from N synthetic fertilizer and animal manure applied to managed soils  | II-21 |
| Figure 2.21 | GHG emissions from LUCF sector from 2000-2005 by source category  | II-24 |
| Figure 2.22 | Categories of GHG emissions sources within the waste sector   | II-26 |
| Figure 2.23 | GHG emissions from the waste sector from 2000-2005 by source category   | II-27 |
| Figure 2.24 | Emissions trends by sector  | II-28 |
| Figure 3.1  | Linkages between climate change-related documents, development planning and the budgeting process   | III-2 |
| Figure 3.2  | Development of the climate change roadmap   | III-3 |
| Figure 3.3  | Process of mainstreaming climate change into national agenda: Policies, guidelines, roadmaps & investment funds (BAPPENAS, 2009b).<br>Note: M is mitigation and A is adaptation.  | III-4 |
| Figure 3.4  | ICCTF development (BAPPENAS, 2009b)   | III-5 |
| Figure 3.5  | Coordination mechanism of ICCTF (BAPPENAS, 2009b)   | III-6 |
| Figure 4.1  | (a) Anomaly of mean global temperature of sea-land and (b) 2001-2005 mean surface temperature relative to 1951-1980 measured at meteorological stations and ship and satellite SST measurements (Hansen <i>et al.</i> , 2006)   | IV-2  |
| Figure 4.2  | Sea level rise trends based on altimeter data from January 1993 to December 2008 (Sofian, 2010)   | IV-3  |
| Figure 4.3  | Seasonal rainfall trends (mm/year) for Indonesia (Boer <i>et al.</i> , 2009)  | IV-5  |
| Figure 4.4a | Distribution of maximum monthly rainfall during the wet and dry seasons in Sibolga-North Sumatra, Maulaboh-West Sumatra and Jatiwangi-West Java (Boer <i>et al.</i> , 2010)   | IV-6  |
| Figure 4.4b | Distribution of maximum monthly rainfall (mm) during the wet and dry seasons in MuaraTewe-Central Kalimantan and Sorong-Papua (Boer <i>et al.</i> , 2009)   | IV-7  |
| Figure 4.5  | Level of probability for seasonal rainfall to change in the future (Note: rainfall in areas colored dark red will decrease in the future, and rainfall in areas colored dark blue will increase in the future with relatively high level of confidence (Boer <i>et al.</i> , 2009)  | IV-8  |
| Figure 4.6  | (a) Summed precipitation for AMJ and JAS for the present climate (dashed line) and for the future predicted climate, using the A2 scenario, and (b) Likelihood of exceeding the 30-day monsoon threshold in 2050. The thick rectangle shows the middle tercile, and the horizontal lines on either side show the lower and upper terciles. The arrows indicate the mean future probability for all GCMs. The vertical |       |

|             |  |       |
|-------------|--|-------|
|             | lines show the observed probability for 1983–2004 (Naylor <i>et al.</i> , 2007).<br>Note: The analysis was done using 15 different GCMs and 3EDMs  | IV-9  |
| Figure 4.7  | Change in cumulative cropping area in the wet season (MH) and dry season (MK) during ENSO years (El Niño and La Niña) compared to normal years (Boer, 2009)  | IV-10 |
| Figure 4.8  | Mean area of dry season rice affected by drought from 1989-2006 (above), during El Niño years (middle) and La Niña years (bottom). Source: Boer (2009)   | IV-11 |
| Figure 4.9  | The relationship between comfort index and length of pregnancy period (left) and comfort index and interval between two births (right). Source: Rohman and Boer, 2001  | IV-13 |
| Figure 4.10 | Changes in rice production in Java by 2025 and 2050 compared to current rice production due to increasing in temperatures and CO <sub>2</sub> concentrations for SRESB1 and SRESA2 using various cultivated area values and planting indices (Boer <i>et al.</i> , 2008)   | IV-15 |
| Figure 4.11 | Increasing percentage of irrigated rice fields from a dam water source by increasing IP to 2.5 for West Java, 2.3 for Central Java and 2.2 for East Java (Boer <i>et al.</i> , 2008)   | IV-16 |
| Figure 4.12 | Expanding irrigated area needed to compensate for loss of rice production in Java due to climate change and land conversation where no change in IP is feasible (Scenario 2). Source: Boer <i>et al.</i> (2008)  | IV-18 |
| Figure 4.13 | Relationship between fraction of unutilized rainfall (f) and forest cover (Herienschah <i>et al.</i> , 2009)   | IV-20 |
| Figure 4.14 | Relationship between HDI and industrial areas in Indonesia (Herienschah <i>et al.</i> , 2009)  | IV-21 |
| Figure 4.15 | Status of water balance under current and future climate (Herienschah <i>et al.</i> , 2009)  | IV-22 |
| Figure 4.16 | Number of districts without surplus water by length under current and future climate scenarios (Herienschah <i>et al.</i> , 2009). Note: Both the bar graph (top) and table (below) show the same data.  | IV-23 |
| Figure 4.17 | Classification of regions at risk to fire based hotspot density patterns – Sumatra (left) and Kalimantan (right) (Ardiansyah and Boer, 2009)   | IV-25 |
| Figure 4.18 | The relationship between hotspot density and rainfall anomaly in Kalimantan and Sumatra for (a) Pattern 1, (b) Pattern 3, (c) Pattern 4, (d) Pattern 5 and (e) Pattern 6   | IV-26 |
| Figure 4.19 | Median value of hot spot density for pattern 5 and pattern 6 regions under current and future climate (Ardiansyah and Boer, 2009)  | IV-27 |
| Figure 4.20 | Keetch and Byram Drought Index for baseline period of 1960-1990 (top), for A2 projection period of 2070-2100 (middle) and for B2 projection period of 2070-2100 (bottom) forecast Kalimantan, Indonesia. The fire danger index rating classes are: low (0-999); moderate (1000-1449); high (1500-1749); extreme (1750 -2000). Source: CIFOR (2008) | IV-27 |

|             |  |       |
|-------------|--|-------|
| Figure 4.21 | The number of surviving and dead trees per hectare after forest fires in 1982/83 and 1997/98 (Ngakan, 2002)  | IV-28 |
| Figure 4.22 | Composition of species per hectare in plots which were lightly, moderately and severely damaged by fires in Kutai National Park, East Kalimantan   | IV-28 |
| Figure 4.23 | Distribution of floods based on the scenario of sea level rise + high tide: 0.25+2.69 m (A) and 1.0+2.69 m (B). Source: Hariati <i>et al.</i> , 2009   | IV-31 |
| Figure 4.24 | Distribution breadth and depth of inundation due to sea level rise and high tide of Medan and surrounding areas (Hariati <i>et al.</i> , 2009)   | IV-32 |
| Figure 4.25 | Distribution of low elevation ( $\leq 0$ meter dpl) areas in DKI Jakarta   | IV-33 |
| Figure 4.26 | Distribution of floods based on sea level rise of 0.25 m (A), 1.0 m (B), sea level rise + high tide of 2.28 m (C) and sea level rise + high tide of 3.03 m (D). Source: Hariati <i>et al.</i> , 2009 | IV-34 |
| Figure 4.27 | Area of flooding and depth of flood due to sea level rise and high tide in the Jakarta region. Source: Hariati <i>et al.</i> , 2009  | IV-35 |
| Figure 4.28 | Number of residents affected by flooding due to sea level rise and high tide in Jakarta. Source: Hariati <i>et al.</i> , 2009  | IV-35 |
| Figure 4.29 | (A) Growth of cone impairment (cone depression) in the groundwater aquifer from 1984 until 2000, and (B) land subsidence rates in Semarang City. Source: Georisk Project, 2008                       | IV-37 |
| Figure 4.30 | Distribution of floods based a sea level rise of 0.25 m (A), 1.0 m (B), and sea level rise + high tide equal to 2.28 m (C) and 3.03 m (D) in Semarang. Source: Hariati <i>et al.</i> , 2009          | IV-38 |
| Figure 4.31 | Flood area and flood depth due to sea level rise and high tide in Semarang. Source: Hariati <i>et al.</i> , 2009   | IV-39 |
| Figure 4.32 | Distribution of floods based on land subsidence and sea level rise of 0.25 m (A), 0.50 m (B), 0.75 m (C) and 1.0 m (D). Source: Hariati <i>et al.</i> , 2009   | IV-39 |
| Figure 4.33 | Flood area and flood depth due to sea level rise and land subsidence in Semarang area. Source: Hariati <i>et al.</i> , 2009  | IV-40 |
| Figure 4.34 | The number of residents affected by flooding due to sea level rise and high tide. Source: Hariati <i>et al.</i> , 2009   | IV-41 |
| Figure 4.35 | Distribution of floods based sea level rise of 0.25 m (A), 1.0 m (B), and sea level rise + high tide equal to 3.16 m (C) and 3.91 m (D) in Surabaya. Source: Hariati <i>et al.</i> , 2009            | IV-42 |
| Figure 4.36 | Flood area and flood depth due to sea level rise and high tide in Surabaya. Source: Hariati <i>et al.</i> , 2009   | IV-43 |
| Figure 4.37 | The number of residents affected by flooding due to sea level rise and high tide in Surabaya. Source: Hariati <i>et al.</i> , 2009   | IV-43 |
| Figure 4.38 | Construction of a dike to reduce flooding from high tides and extreme waves in East Harbour of Muara Bary, Jakarta. Source: Media Indonesia, 2007  | IV-44 |

|             |  |       |
|-------------|--|-------|
| Figure 4.39 | Incidence rate of dengue fever and the number of affected cities and districts in Note: 1973, 1988 and 1998 were La Niña years. Source: Ministry of Health.  | IV-46 |
| Figure 4.40 | Annual trend of dengue incidence rates in districts in Java in period between 1992-2005 (cases per 100,000 people per year). Source: MoE, 2007b.   | IV-47 |
| Figure 4.41 | Transmission potential of dengue fever under current and future climate (Hidayati <i>et al.</i> , 2009). Note: Red represents the highest risk and blue represents the lowest risk   | IV-49 |
| Figure 4.42 | Transmission potential of malaria under current and future climate (Hidayati <i>et al.</i> , 2009). Note: Red represents the highest risk and blue represents the lowest risk  | IV-50 |
| Figure 5.1  | Energy projections by type of energy under the BAU scenario, 2010-2030   | V-5   |
| Figure 5.2  | Power plant projections by fuel type under the BAU scenario, 2010-2030   | V-6   |
| Figure 5.3  | Projection of GHG emissions generation by fuel type under the BAU scenario, 2010-2030  | V-7   |
| Figure 5.4  | Projection of GHG emissions generation by sector under the BAU scenario, 2010-2030   | V-7   |
| Figure 5.5  | GHG emissions projections by fuel type under BAU, CLIMATE and PERPRES scenarios  | V-11  |
| Figure 5.6  | Total GHG emissions projections under BAU, CLIMATE and PERPRES scenarios   | V-11  |
| Figure 5.7  | GHG emissions reduction potential under BAU, CLIMATE 1, CLIMATE 2, and PERPRES scenarios   | V-12  |
| Figure 5.8  | Total projected demand of biofuel in the transportation and industrial sectors and estimated investment cost of the biofuel production system  | V-14  |
| Figure 5.9  | Rate of carbon sequestration and emissions under the BAU scenario and the two mitigation scenarios (Ministry of Forestry)  | V-23  |
| Figure 5.10 | The projection of methane emissions from rice paddies under the baseline and mitigation scenarios (Setyanto <i>et al.</i> , 2009)  | V-28  |
| Figure 5.11 | Projected methane emissions from livestock (Suryahadi <i>et al.</i> , 2009)  | V-31  |
| Figure 5.12 | Projection of emissions of all sectors under BAU and mitigation scenarios (MoE, 2009). Note: Contribution of transportation sector to the total energy emission is approximately 20% (based on emissions data from 2000-2006). | V-33  |
| Figure 6.1  | Number of NGOs working in climate change mitigation activities on energy and waste sectors (Boer <i>et al.</i> , 2008)   | VI-1  |
| Figure 6.2  | Existing operational Sea Level Monitoring Stations in Indonesia (MoE, 2008)  | VI-4  |
| Figure 6.3  | Number of agroclimatic stations by province (Surmaini <i>et al.</i> , 2010)  | VI-5  |

## List of Figure

|            |  |       |
|------------|--|-------|
| Figure 6.4 | Donor countries/agencies who contribute to the implementation of capacity building activities on climate change (Based on data collected from the Ministry of Environment) | VI-7  |
| Figure 6.5 | Types of capacity building activities on climate change by year (Based on data collected from the Ministry of Environment)   | VI-8  |
| Figure 6.6 | Development of web-based climate information system at four regional offices (Provided by the BMKG)  | VI-8  |
| Figure 7.1 | Climate Change Financing Scheme (BAPPENAS, 2009)   | VII-2 |

# List of Table

|            |  |       |
|------------|--|-------|
| Table 1.1  | Per capita GDP and growth rates of the economy, 1997-2007  | I-5   |
| Table 1.2  | Labor force and open unemployment in Indonesia, 2000-2008  | I-6   |
| Table 1.3  | Type of final energy consumption by sector, 2007 MMBOE   | I-7   |
| Table 1.4  | Growth and production of industrial products (000 ton) in the period 2000-2005   | I-9   |
| Table 1.5  | Forested and non-forested land in forest and non-forest area in 2007<br>(thousand ha)  | I-10  |
| Table 2.1  | Summary of national GHG emissions in 2000 (in Gg CO <sub>2</sub> e)  | II-4  |
| Table 2.2  | Energy sector emissions from 2000 to 2005 by gas (Gg CO <sub>2</sub> e)  | II-12 |
| Table 2.3  | CO <sub>2</sub> emissions from fossil consumption and fugitive emission from 2000-2005 (Gg CO <sub>2</sub> e)  | II-12 |
| Table 2.4  | GHG emissions from the industrial sector from 2000 to 2005 by gas<br>(in Gg CO <sub>2</sub> e)   | II-13 |
| Table 2.5  | GHG emissions from the industrial sector from 2000 to 2005 by source<br>category (in Gg CO <sub>2</sub> e)   | II-14 |
| Table 2.6  | GHG emissions from the agriculture sector from 2000 to 2005 by gas<br>(in Gg CO <sub>2</sub> e)  | II-15 |
| Table 2.7  | Livestock population in Indonesia from 2000-2006   | II-16 |
| Table 2.8  | Direct and indirect N <sub>2</sub> O emissions from manure management  | II-18 |
| Table 2.9  | Distribution of GHG emissions from grassland burning from 2000-2005<br>(Gg CO <sub>2</sub> e)  | II-21 |
| Table 2.10 | Distribution of GHG emissions from cropland burning from 2000-2005<br>(Gg CO <sub>2</sub> e)   | II-22 |
| Table 2.11 | Adjustment of Indonesia's land category by Ministry of Forestry into 2006<br>IPCC GL categories  | II-22 |
| Table 2.12 | GHG emissions and removal from LUCF (in Gg CO <sub>2</sub> e)  | II-23 |
| Table 2.13 | Emissions from peat fires according to relevant studies  | II-25 |
| Table 2.14 | Impact of selecting different emissions estimates from peat fire on national<br>figures and the relative position of Indonesia as an emitter country   | II-25 |
| Table 2.15 | GHG emissions and removal from waste (in Gg CO <sub>2</sub> e)   | II-27 |
| Table 2.16 | Key Category Analysis  | II-28 |
| Table 2.17 | Estimates of uncertainty level for activity data and emissions factors   | II-30 |
| Table 2.18 | Level of current uncertainty of Indonesian National GHG Inventory for<br>2000 and 2004 and trend and potential reduced uncertainty with the<br>incorporation of necessary improvement for the first three key source<br>categories | II-30 |

|            |  |       |
|------------|--|-------|
| Table 4.1  | Decrease in rice production in Java compared to current production as increasing temperature and CO <sub>2</sub> concentration in Scenario1 (Boer <i>et al.</i> , 2009)            | IV-14 |
| Table 4.2  | Total rice field area (ha) in three provinces in Java for 2025 and 2050 using a rice field conversion rate of 0.77% per year (Scenario 2)  | IV-15 |
| Table 4.3  | Cropping index scenario for rice in three provinces in Java for 2025 and 2050 (Boer <i>et al.</i> , 2008)  | IV-16 |
| Table 4.4  | Increased productivity levels needed to maintain current production levels in 2025 in 2050 under Scenario 5 (IP increased and land conversion occurred; Boer <i>et al.</i> , 2008) | IV-17 |
| Table 4.5  | Increased productivity levels needed maintain current production in 2025 and 2050 under Scenario 6 (IP constant and land conversion occurred)                                      | IV-17 |
| Table 4.6  | Number of districts that have no months without water surplus (Herienschah <i>et al.</i> , 2009)   | IV-23 |
| Table 4.7  | Area affected by forest fire in 1997/1998 (expressed in ha)  | IV-24 |
| Table 4.8  | Total national economic loss due to fires during the 1997 El Niño year (expressed in million USD)  | IV-25 |
| Table 4.9  | Economic loss per ha of affected land use type   | IV-31 |
| Table 4.10 | Impact of sea level rise and high tide on population and land use in Medan (Hariati <i>et al.</i> , 2009)  | IV-32 |
| Table 4.11 | Percent distribution area of low elevation (<0 meters below sea level in DKI Jakarta)  | IV-33 |
| Table 4.12 | Impact of sea level rise and high tide on the population and land use in Jakarta Source: Hariati <i>et al.</i> , 2009  | IV-36 |
| Table 4.13 | Shoreline retreat due to sea level rise and land subsidence (Hariarti <i>et al.</i> , 2009)  | IV-40 |
| Table 4.14 | Impact of sea level rise, high tide and land subsidence on the population and land use in Semarang. Source: Hariarti <i>et al.</i> , 2009  | IV-41 |
| Table 4.15 | Impact of sea level rise and high tide on population and land uses in Surabaya Source: Hariati <i>et al.</i> , 2009  | IV-43 |
| Table 4.16 | The ten main of health problems recorded by hospitals in Indonesia in 2006 (Source: Ditjen Pelayanan Medik - Ministry of Health, 2006)   | IV-45 |
| Table 4.17 | Percent change in transmission potential for malaria and dengue fever (Hidayati <i>et al.</i> , 2009)  | IV-48 |
| Table 4.18 | General adaptation options to reduce the potential health impacts of climate change  | IV-51 |
| Table 5.1  | Projection of energy supply mix under BAU, CLIMATE and PERPRES scenarios (million BOE)   | V-9   |
| Table 5.2  | Projection of sectoral energy demand under BAU, CLIMATE and PERPRES scenarios (million BOE)  | V-10  |
| Table 5.3  | Projection of installed capacity demand of power generation under BAU, CLIMATE 1 and CLIMATE 2 scenarios (in MW)   | V-10  |



|            |  |      |
|------------|--|------|
| Table 5.4  | Selected mitigation options  | V-13 |
| Table 5.5  | Additional demand of biofuel and GHG emissions reduction potential and cost  | V-15 |
| Table 5.6  | New investments for advanced (more efficient) coal power plants under the BAU, CLIMATE1 and CLIMATE 2 scenarios                | V-15 |
| Table 5.7  | The installed capacity, additional investment and GHG emissions of natural gas power plants from 2010-2030                     | V-16 |
| Table 5.8  | The installed capacity and the required investment for renewable power plants under the BAU, CLIMATE 1 and CLIMATE 2 scenarios | V-17 |
| Table 5.9  | Energy efficiency improvements by sector, 2020-2030  | V-18 |
| Table 5.10 | Projected emissions from the industrial sector under BAU and mitigation scenarios (Ggram CO <sub>2</sub> e)                    | V-19 |
| Table 5.11 | Rate of planting in the Strategic Plan of the Ministry of Forestry (MoF, 2009)   | V-23 |
| Table 5.12 | Program support from international agencies for forestry mitigation projects in Indonesia                                      | V-24 |
| Table 5.13 | Mitigation potential and cost for forestry sector  | V-26 |
| Table 5.14 | Mitigation scenarios for reducing methane emissions from rice paddies  | V-29 |
| Table 5.15 | Projected mitigation technology options adopted for livestock  | V-31 |
| Table 5.16 | Projection of GHG emissions from waste sector under BAU and alternative scenarios (Dewi <i>et al.</i> , 2009)                  | V-33 |
| Table 6.1  | Rain gauges density in Indonesia   | VI-4 |



# Glossary of Abbreviations

|                        |   |                       |  |
|------------------------|---|-----------------------|--|
| <b>EDM</b>             | = Empirical Downscaling Model   | <b>CWPB</b>           | = Centre-Worked Prebake                                      |
| <b>AD</b>              | = Activity Data   | <b>Depsos</b>         | = Ministry of Social   |
| <b>ADB</b>             | = Asian Development Bank  | <b>DFID</b>           | = United Kingdom Department for International Development    |
| <b>ADPC</b>            | = Asian Disaster Preparedness Centre  | <b>DJF</b>            | = December, January, February                                |
| <b>ADRC</b>            | = Asian Disaster Reduction Center   | <b>DNM</b>            | = Directorate for Nature and Management                      |
| <b>AFC</b>             | = Asian Financial Crisis  | <b>DNPI</b>           | = National Committee on Climate Change                       |
| <b>Al</b>              | = Aluminum  | <b>DNS</b>            | = Debt for Nature Swaps                                      |
| <b>AMJ</b>             | = April, May, June  | <b>DOC</b>            | = Degradable Organic Content                                 |
| <b>APL</b>             | = Area Penggunaan Lain (Other land uses)  | <b>EE</b>             | = Energy Efficiency  |
| <b>APPI</b>            | = Association of Fertilizer Producers   | <b>EF/RF</b>          | = Emissions Factor/Removal Factor                            |
| <b>AR</b>              | = Afforestation and Reforestation   | <b>EFBs</b>           | = Empty Fruit Bunches  |
| <b>AS</b>              | = Ammonium Sulfate  | <b>ENSO</b>           | = El Niño Southern Oscillation                               |
| <b>AWS</b>             | = Automatic Weather Stations  | <b>f</b>              | = Fraction of rainfall that cannot be utilized               |
| <b>BALITKILIMAT</b>    | = Research Agency on Agro-climatology and Hydrology                               | <b>FBS</b>            | = Feed Supplement Blocks                                     |
| <b>BAPLAN</b>          | = Directorate General of Planology  | <b>FCCP</b>           | = Forestry-Climate Change Project                            |
| <b>BAPPENAS</b>        | = National Agency for Planning and Development                                    | <b>FCPF</b>           | = Forest Carbon Partnership Facility                         |
| <b>BAU</b>             | = Business as Usual   | <b>FMU</b>            | = Forest Management Unit                                     |
| <b>BBSLDA</b>          | = Center for Agriculture Land Resources and Development                           | <b>FMUs</b>           | = Forest Management Units                                    |
| <b>BI</b>              | = Births  | <b>GAW</b>            | = Global Atmospheric Watch                                   |
| <b>BMKG</b>            | = Meteorology, Climatology, and Geophysical Agency                                | <b>GCMs</b>           | = General Circulation Models                                 |
| <b>BOD</b>             | = Biological Oxygen Demand  | <b>GDP</b>            | = Gross Domestic Product                                     |
| <b>bpd</b>             | = barrels per day   | <b>GEF</b>            | = Global Environment Facility                                |
| <b>BPPT</b>            | = Agency for Technology Assessment and Application                                | <b>GEF</b>            | = Global Environmental Fund                                  |
| <b>BPS</b>             | = Badan Pusat Statitistik (Indonesian Bureau of Statistics)                       | <b>GFC</b>            | = Global Financial Crisis                                    |
| <b>BPSDA</b>           | = Balai Pengelolaan Sumberdaya Air  | <b>GFF</b>            | = Global Forest Fund   |
| <b>C</b>               | = Confidential  | <b>GHGs</b>           | = Greenhouse Gases   |
| <b>CBM</b>             | = Coal Bed Methane  | <b>GL</b>             | = Guidelines   |
| <b>CCROM-IPB</b>       | = Centre for Climate Risk and Opportunity Management-Bogor Agriculture University | <b>GL</b>             | = Grass Land   |
| <b>CCS</b>             | = Carbon Capture and Storage  | <b>GOI</b>            | = Government of Indonesia                                    |
| <b>CDM</b>             | = Clean Development Mechanism   | <b>GPG</b>            | = Good Practice Guidance                                     |
| <b>CFBC</b>            | = Coal Fluidized Bed Combustion   | <b>NFI</b>            | = National Forest Inventory                                  |
| <b>CH<sub>4</sub></b>  | = Methane   | <b>H<sub>2</sub>S</b> | = Hydrogen Sulfide   |
| <b>CI</b>              | = Cropping Index  | <b>ha</b>             | = Hectare  |
| <b>CIFOR</b>           | = for International Forestry Research   | <b>HDI</b>            | = Human Development Index                                    |
| <b>CL</b>              | = Crops Land  | <b>HK</b>             | = Conservation Forest  |
| <b>CO<sub>2</sub></b>  | = Carbon Dioxide  | <b>HL</b>             | = Protection Forest  |
| <b>CO<sub>2</sub>e</b> | = Carbon Dioxide Equivalent   | <b>HoB</b>            | = Heart of Borneo  |
| <b>COP</b>             | = Conference of Parties   | <b>HP</b>             | = Production Forest  |
| <b>COP13</b>           | = 13th Conference of Parties  | <b>HPK</b>            | = Convertible Production Forest                              |
| <b>COP15</b>           | = 15th Conference of Parties  | <b>IAARD</b>          | = Indonesian Agency for Agriculture Research and Development |
| <b>COREMAP</b>         | = Coral Reef Rehabilitation and Management Program                                | <b>ICCTF</b>          | = Indonesian Climate Change Trust Fund                       |
| <b>CRED</b>            | = Center for Research on the Epidemiology of Disaster                             | <b>IDR</b>            | = Indonesia Rupiah   |
|                        |   | <b>IE</b>             | = Including Elsewhere  |
|                        |   | <b>IFCA</b>           | = Indonesian Forest Climate Alliance                         |
|                        |   | <b>IGCC</b>           | = Integrated Gasification Combined Cycle                     |
|                        |   | <b>INCAS</b>          | = Indonesian National Carbon Accounting System               |
|                        |   | <b>INDEF</b>          | = Institute for Development Economics and Finance            |
|                        |   | <b>IndTEWS</b>        | = Indonesia Tsunami Early Warning System                     |

## Glossary of Abbreviations

|                     |   |                     |  |
|---------------------|---|---------------------|--|
| <b>Inpres</b>       | = Presidential Instruction  | <b>NTT</b>          | = East Nusa Tenggara   |
| <b>IPO</b>          | = Current Planting Index  | <b>ODA</b>          | = Official Development Assistance                                |
| <b>IPB</b>          | = Bogor Agriculture University / Institut Pertanian Bogor                     | <b>ODA</b>          | = Overseas Development Assistance                                |
| <b>IPCC</b>         | = Intergovernmental Panel on Climate Change                                   | <b>OFDA</b>         | = Office of U.S. Foreign Disaster Assistance                     |
| <b>IRI</b>          | = International Research Institute for Climate and Society                    | <b>OM</b>           | = Organic Matter   |
| <b>ITB</b>          | = Bandung Institute of Technology   | <b>P</b>            | = Population Density   |
| <b>ITCZ</b>         | = Inter-tropical Convergence Zone   | <b>PDO</b>          | = Pacific Decadal Oscillation                                    |
| <b>JAS</b>          | = July, August, September   | <b>PEACE</b>        | = Pelangi Energi Abdi Citra Enviro                               |
| <b>JJA</b>          | = June, July, August  | <b>Permenhut</b>    | = Ministry of Forestry Regulation                                |
| <b>KAfarm</b>       | = Water demand for agriculture  | <b>PERPRES</b>      | = Presidential Decree  |
| <b>KAindustry</b>   | = Water demand for industrial sectors   | <b>PFCs</b>         | = Perfluorocarbons   |
| <b>KApopulation</b> | = Water demand for domestic use   | <b>PLN</b>          | = National Electricity Enterprise                                |
| <b>kc</b>           | = Crop Coefficient  | <b>PP</b>           | = Pregnancy Period   |
| <b>NTB</b>          | = West Nusa Tenggara  | <b>PP</b>           | = Polypropylene  |
| <b>KEN</b>          | = National Energy Policy  | <b>PP</b>           | = Government Regulation  |
| <b>Kep. MENLH</b>   | = Decree of the State Minister of Environment                                 | <b>PU</b>           | = Ministry of Public Works                                       |
| <b>Kepmen Ekuin</b> | = Ministerial Decree  | <b>QA/QC</b>        | = Quality Assurance/Quality Control                              |
| <b>Km</b>           | = Kilometer   | <b>RAN-GRK</b>      | = National Action Plan for Reducing Greenhouse Gas Emissions     |
| <b>LAPAN</b>        | = National Agency for Aviation and Space                                      | <b>REDD</b>         | = Reducing Emissions from Deforestation and Forest Degradation   |
| <b>LFG</b>          | = Landfill Gas  | <b>REL</b>          | = Reference Emission Level                                       |
| <b>LIPI</b>         | = The Center for Oceanography Research of the Indonesian Institute of Science | <b>RKP</b>          | = Rencana Kerja Pemerintah                                       |
| <b>LPG</b>          | = Liquefied Petroleum Gas   | <b>Rmonthly</b>     | = Monthly Rainfall   |
| <b>LPMH</b>         | = Harvested Area During the Wet Season  | <b>RPJMN</b>        | = The Mid-Term National Development Plan                         |
| <b>LPMK</b>         | = Harvested Area During the Dry Season  | <b>RUKN</b>         | = General Plan of National Electricity                           |
| <b>LPTP</b>         | = Lembaga Pengembangan Teknologi Pedesaan                                     | <b>RUPTL</b>        | = Business Plan of Electricity Supply                            |
| <b>LUCF</b>         | = Land Use Change and Forestry  | <b>SCCF</b>         | = Special Climate Change Fund                                    |
| <b>LULUCF</b>       | = Land Use, Land Use Change and Forestry                                      | <b>SGP-GEF UNDP</b> | = Small Grant Program-Global Environmental Facility UNDP         |
| <b>MAM</b>          | = March, April, May   | <b>SIGN</b>         | = Sistem Inventarisasi Gas Rumah Kaca Nasional                   |
| <b>MDGs</b>         | = UN Millennium Development Goals   | <b>SLR</b>          | = Sea Level Rise   |
| <b>MEMR</b>         | = Ministry of Energy and Mineral Resources                                    | <b>SNC</b>          | = Second National Communication                                  |
| <b>MEWS</b>         | = Meteorological Early Warning System   | <b>SNI</b>          | = National Standard  |
| <b>MH</b>           | = Wet Season  | <b>SON</b>          | = September, October, November                                   |
| <b>MK</b>           | = Dry Season  | <b>SSTs</b>         | = Sea Surface Temperatures                                       |
| <b>MMBOE</b>        | = Million Barrel Oil Equivalent   | <b>SWDS</b>         | = Solid Waste Disposal Site                                      |
| <b>MoA</b>          | = Ministry of Agriculture   | <b>SWS</b>          | = Satuan Wilayah Sungai  |
| <b>MoF</b>          | = Ministry of Forestry  | <b>TC</b>           | = Technical Cooperation  |
| <b>MRT</b>          | = Mass Rapid Transport  | <b>TERANGI</b>      | = Indonesian Coral Reef Foundation                               |
| <b>MRV</b>          | = Reporting and Verification  | <b>TOW</b>          | = Total Organic Waste  |
| <b>MSE</b>          | = Ministry of State of Environment  | <b>TP</b>           | = Potential Transmissions  |
| <b>MSW</b>          | = Municipal Solid Waste   | <b>UNDP</b>         | = United Nations Development Programme                           |
| <b>MW</b>           | = Megawatts   | <b>UNEP</b>         | = United Nations Environment Programme                           |
| <b>N</b>            | = Nitrogen  | <b>UNFCCC</b>       | = United Nations Framework Convention on Climate Change          |
| <b>N2O</b>          | = Nitrous Oxide   | <b>UN-OCHA</b>      | = United Nations Office for Coordination of Humanitarian Affairs |
| <b>NA</b>           | = Not Applicable  | <b>USD</b>          | = United State Dollar  |
| <b>NAD</b>          | = Nanggroe Aceh Darusalam   | <b>USDA</b>         | = United State Department of Agriculture                         |
| <b>NAMA</b>         | = National Appropriate Mitigation Action                                      | <b>WHO</b>          | = World Health Organization                                      |
| <b>NAO</b>          | = North Atlantic Oscillation  | <b>WRI</b>          | = World Resources Institute                                      |
| <b>NE</b>           | = Note Estimated  | <b>WWF</b>          | = World Wildlife Fund  |
| <b>NGHGI</b>        | = The Indonesian National Greenhouse Gas Inventory                            | <b>WWT</b>          | = Wastewater Treatment   |
| <b>NGOs</b>         | = Environmental non-Governmental Organizations                                |                     |  |
| <b>NO</b>           | = Not Occurring   |                     |  |
| <b>NPK</b>          | = Nitrogen, Phosphorus And Potassium  |                     |  |
| <b>NTFP</b>         | = Non Timber Forest Product   |                     |  |

## CHAPTER I

# National Circumstances

### 1.1 Geography and Climate

Indonesia is located between 6°08' North and 11°15' South latitude, and from 94°45' to 141°05' East longitude. The country covers 790 million hectares (ha), with a total coastline of 81,000 km and land territory of about 200 million ha. It has five large islands (Sumatra, Java, Kalimantan, Sulawesi and Irian Jaya) and approximately 13,667 small islands, of which more than half (56%) are nameless and only 7% are permanently inhabited (Figure 1.1). Extensive coastal plains and mountain areas of 1,000 metres (m) above sea level are characteristics of Sumatra, Kalimantan and Irian Jaya. Of the 200 million ha of land territory, about 50 million ha are devoted to various agricultural activities. There is nearly 20 million ha of arable land, of which about 40% is wetland (e.g., rice fields), 40% is dry land, and 15% is shifting cultivation. Since 2005, the Republic of Indonesia has been divided administratively into 33 provinces.



Figure 1.1 Map and overview of Indonesia, 2007

Indonesia's climate is dominated by monsoons, which give a degree of homogeneity across the region. Indonesia lies across the range of the Inter-tropical Convergence Zone (ITCZ) where the northeast and southeast trade winds penetrate the doldrums. Strong ascending motion, overcast skies, strong squalls, heavy rainfall and severe local thunderstorms with variable intensities are characteristics of this zone.

The rainfall pattern in Indonesia consists of three types (Boerema, 1938). The first type is a monsoon rainfall with a monthly rainfall peak in December. The second is a more localised rainfall pattern in the eastern equatorial part of the country with a monthly rainfall peak in July-August. The third rainfall type is an equatorial rainfall characterized by two monthly rainfall peaks, in March and in October.

The monsoon rainfall can be further categorized into two groups that are prevalent in southern Indonesia. While both groups indicate a clear distinction between a dry and a wet season throughout the year, the first shows a longer dry period (i.e., in the eastern Nusa Tenggara islands) and is drier overall compared to the second group (i.e., the western islands of Java, South Sumatra and South Sulawesi). The regions in the first group experience severe drought more frequently. Also, in general, rainfall variation is larger during the dry season (April to September) compared to the wet season (October to March).


The more localised rainfall pattern located mainly in the eastern equatorial part of Indonesia (e.g., Maluku and Sorong) results in wet seasons between April and September and dry seasons between October and March. The dry periods of this type of rainfall are not as severe as those of monsoonal types; as a result, the total rainfall in a given year is relatively high.

The third type of rainfall, equatorial, can be divided into two groups. The first group covers the west coast of North Sumatra with no pronounced dry season and with slightly increased rainfall around March and October. The second group covers the west coast of South Sumatra with uniform distribution of rainfall throughout the year.

Overall, these three types of rainfall result in a wet season that varies in length from as long as 280 to 300 days or as short as 10 to 110 days, with the rainfall during that season varying from 4,115 mm to as low as 640 mm. Rainfall variability in Indonesia is strongly affected by phenomena associated with ENSO (El Niño Southern Oscillation). The impact of ENSO on rainfall is more pronounced during the dry season than the wet season (USDA, 1984; Las *et al.*, 1999). Various studies on ENSO influences on inter-annual rainfall variability in Indonesia reveal the following seasonal patterns (USDA, 1984; ADPC, 2000; Yoshino *et al.*, 2000; and Kirono and Partridge, 2002): (i) the end of the dry season occurs later than normal during El Niño and earlier during La Niña years, (ii) the onset of the wet season is delayed during El Niño and advanced during La Niña years, (iii) a significant reduction of dry season rainfall could be expected during El Niño and a significant increase during La Niña years, (iv) long dry spells occur during the monsoon period, particularly in Eastern Indonesia. Moreover, as well as causing rainfall variation, the effect of ENSO itself is varied. ENSO effects are particularly strong in regions that are strongly influenced by monsoon systems, weaker in regions that are dominated by equatorial systems, and not traceable in regions driven by local weather systems (Tjasyono, 1997).

## 1.2 Extreme Climate Events


Extreme climate events in Indonesia are normally associated with ENSO. El Niño events are associated with drought and La Niña event are associated with heavy rainfall and flooding. Because the ENSO signal is very strong in the regions dominated by monsoonal rainfall (i.e., Java, Bali and Nusa Tenggara), extreme events are most pronounced in these areas. In these islands, for example, the decrease in dry season rainfall could be twice that of the other islands (Figure 1.2). Over recent years, El Niño events have become more frequent as the global temperature anomalies continue to increase (Hansen *et al.*, 2006). This indicates that the extreme regional weather and climate anomalies associated with El Niño are being exacerbated by increasingly higher temperatures. These warmer conditions have been linked to higher concentrations of atmospheric greenhouse gases (GHGs) (Timmermann *et al.* 1999; Collins 2000, b; Cubasch *et al.* 2001; Chen *et al.* 2005; Guilyardi 2005).



**Figure 1.2 Percent decrease in seasonal rainfall from normal during two strong El Niño years, 1997 and 1982 (Irawan, 2002)**

With the increasing frequency and intensity of extreme climate events, the most affected group will be the poor, as their adaptive capacity is most limited. If more vulnerable groups are exposed to repeated climate hazards, resilience will be more quickly compromised, making it increasingly more difficult to recover. In many cases, this is leading to high levels of unsecured and informal household debt. The Government of Indonesia also spends significant amounts of public funding in an effort to reduce vulnerability and improve resilience of poor populations. Based on data collected by the Office of U.S. Foreign Disaster Assistance/Center for Research on the Epidemiology of Disaster (OFDA/CRED) International Disaster Database (2007) from 1907 to 2007, it is clear that climate-related hazards are becoming increasingly more dominant in

the current decade (2000-2010). From the record, the number of natural hazards categorized as global hazards was 345 events; about 60% were climate-related hazards. The first climate hazards occurred in 1953 and they became more frequent after 1980s (Figure 1.3). The most frequent hazard was flooding, followed by landslides and water or vector-borne diseases, wind storms, forest fires, drought and high tide/storm surge.



**Figure 1.3 Number of climate hazards by type (left) and by year (right).**  
Source: Based on data from OFDA/CRED International Disaster Database (2007)

### 1.3 Economic and Social Development

Indonesia is the fourth most populous nation in the world following China, India and the United States. According to national statistics (Indonesian Bureau of Statistics – BPS 2005/2006), the total population of Indonesia in 1971 was 119 million, increasing to 230.3 millions in 2008. Currently, the population growth rate appears to be slowing from 1.49% (1990–2000) to 1.34% (2001–2005). Nevertheless, it is projected that Indonesia’s population will exceed 300 million by 2030.

Life expectancy has improved from 41 years in 1990 to 65 years in 1997. However, adult illiteracy dropped 40% in just 37 years. Based on BPS data from 2005, the number of people living in urban areas totals approximately 44 million people (20% of the total population), while people living on coastal areas reaches roughly 20 million (9% of the total population). The majority of people living near the coast are poor, particularly in North Sumatra, parts of West and Central Java, and most of the islands in the eastern parts of Indonesia and Papua.

Indonesia had been experiencing marked improvements in its poverty rates until the economic crisis in 1999. Prior to 1999, Indonesia had been successful in its programme to alleviate poverty – in 1970, 60% of the population (70 million people) was living in absolute poverty; by 1990, the



number had dropped to 27 million or 15% of the population and continued to improve to 1997, when the figure decreased to 20 million. However, in 1999, for the first time in years, Indonesia experienced a severe 18-month drop in the country's social condition, resulting in over 100 million people living below the poverty line.

As such, poverty remains a challenge. Unemployment and underemployment are also still relatively high. The Mid-Term National Development Plan (RPJMN) 2004–2009, its target for poverty and unemployment was 5.1% and 8.2%, respectively. A recent report from BPS stated that unemployment by February 2008 reached 8.46% or about 9.43 million people. However, based on the projection of the Institute for Development Economics and Finance (INDEF), due to current economic crisis, the number of poor and unemployed may have increased to 9.5% and 16.3%, respectively, in 2008 (Depsos, 2009).

Indonesia's gross domestic product (GDP) is approximately US\$175 billion, with trade (16.7%), manufacturing (28%), agriculture (15.4%) and services (10.17%) as the main contributors. Earnings from exports (oil and gas, textiles, appliances, coal, copper) were approximately US\$69 billion, while imports (food, chemicals, capitals and consumer goods) generated about US\$44.8 billion. The GDP growth has increased steadily since 1998 and reached 6.3% in 2007 (Table 1.1). This has brought per capita GDP to around US\$2000, which is considerably higher than the pre-crisis level (around US\$1,000; Siregar, 2008). The role of non-oil and gas sectors has become more significant since the Asian Financial Crisis (AFC); the oil and gas sectors' contribution to the economy decreased, as indicated by the negative growth rates for the whole period, except in the year 2000. Among the non-oil and gas sectors, agriculture has played an important role. It is the sector least affected by the AFC and in fact helped the recovery of Indonesian economy after the crisis. This was due to its role as "the employer of last resort" and its substantial increases in exports (Siregar, 2008).

**Table 1.1 Per capita GDP and growth rates of the economy, 1997-2007**

| Economic parameters                    | 1997 | 1998  | 1999 | 2000 | 2001 | 2002 | 2003 | 2004 | 2005 | 2006 | 2007 |
|--|------|-------|------|------|------|------|------|------|------|------|------|
| Per capita GDP (USD)                   | 1093 | 478   | 689  | 804  | 772  | 922  | 1098 | 1186 | 1318 | 1663 | 1946 |
| GDP Growth                             | 4.7  | -13.1 | 0.8  | 4.9  | 3.6  | 4.5  | 4.8  | 5    | 5.7  | 5.5  | 6.3  |
| Non-Oil & Gas GDP Growth (% per annum) | 5.2  | -14.2 | 1    | 5.3  | 4.9  | 5.2  | 5.7  | 6    | 6.6  | 6.1  | 6.9  |
| Oil-Gas GDP Growth (% per annum)       | -1   | -0.5  | -1.3 | 1    | -5.3 | -1.3 | -2.9 | -3.5 | -3.1 | -1.3 | -0.8 |
| Manufacturing GDP Growth (% per annum) | 5.3  | -11.4 | 3.9  | 6    | 3.3  | 5.3  | 5.3  | 6.4  | 4.6  | 4.6  | 4.7  |
| Agriculture GDP Growth (% per annum)   | 1    | -1.3  | 2.2  | 1.9  | 3.3  | 3.4  | 3.8  | 2.8  | 2.7  | 3.4  | 3.5  |

Source: (Based on BPS data; Siregar, 2008)

Success in pursuing macroeconomic stability over the past few years has been the primary factor in reducing poverty in Indonesia. However, economic growth alone is not sufficient for improving the well-being of the poor; additional efforts are still needed to address their needs. Indeed, the global financial crisis triggered by the U.S. sub-prime mortgage crisis caused some doubt about the continuation of post-AFC Indonesian economic growth. Gradual increases in commodity prices, especially in the agriculture sector, followed by abrupt decreases in 2009 have exacerbated uncertainties (Siregar, 2008). This may negatively affect efforts to further develop the commodities and related industries.

The current global financial crisis (GFC) is distinct from the AFC in 1997/98. In the case of the AFC, the significant rupiah depreciation was followed by increases in Indonesian exports, especially of agriculture, shortening the recession period. The current depreciation of rupiah following the GFC seems unlikely to be followed by similar increases in exports because the main export destinations around the world have also been adversely affected; the recession may thus last considerably longer (Siregar, 2008).

Despite positive macroeconomic achievements, the country still faces major challenges in a number of key areas, such as employment creation, improvement of the investment climate and ensuring that Indonesia fully benefits from its integration into the international trading system (European Community, 2005). Between the onset of the AFC in 1997 until 2006, unemployment had continuously increased in absolute as well as in percentage terms, reaching 11.1 million (10.4 percent) (Table 1.2). Although this has been reduced over the course of the last two years, the figures are still reasonably high, with 9.4 million people (8.5%) unemployed in February 2008. Due to the current GFC, the number of unemployed may continue to increase if the main sources of recovery and growth are capital-intensive sectors (Siregar, 2008).

**Table 1.2 Labor force and open unemployment in Indonesia, 2000-2008**

| Year   | Labor Force<br>(Million people) | Open Unemployment<br>(Million People) | Open Unemployment<br>(% of total) |
|--------|---------------------------------|---------------------------------------|-----------------------------------|
| 2000   | 95.65                           | 5.81                                  | 6.08                              |
| 2001   | 98.81                           | 8.01                                  | 8.10                              |
| 2002   | 100.78                          | 9.13                                  | 9.06                              |
| 2003   | 102.63                          | 9.82                                  | 9.57                              |
| Aug-04 | 104.00                          | 10.3                                  | 9.90                              |
| Feb-05 | 105.80                          | 10.9                                  | 10.30                             |
| Feb-06 | 106.30                          | 11.1                                  | 10.44                             |
| Feb-07 | 108.13                          | 10.55                                 | 9.76                              |
| Feb-08 | 111.48                          | 9.43                                  | 8.46                              |

Source: (Based on BPS data; Siregar, 2008)

## 1.4 Sectoral Conditions

### 1.4.1 Energy Sector

Indonesian final energy consumption has been growing in line with the country's economic and population growth. According to the latest official published data from Data and Information Center (PUSDATIN) – MEMR (2009a): Indonesia Energy-Economic Handbook 2009, the domestic consumption of final energy (excluding biomass) in Indonesia grew from 495.45 MMBOE (million barrel oil equivalent) in 2000 to 564.94 MMBOE in 2005 (at average growth rate of 3.1 % per year). The majority of this consumption is accounted by industry sector, followed by transportation, residential, commercial, and ACM (agriculture, construction, and mining) sectors (see Figure 1.4). In 2008, the final consumption is about 658.36 MMBOE. The share of total energy consumption by sector in 2008 is industrial sector 48%, transportation 31%, household 13%, commercial 4%, and ACM 5% (Table 1.3).




Figure 1.4 Final energy consumption in Indonesia, 2000 – 2007 [MEMR, 2009]

Table 1.3 Type of final energy consumption by sector, 2008 MMBOE [MEMR, 2009]

| Sector         | Oil Products | LPG | Natural Gas | Coal | Electricity | Total | %   |
|----------------|--------------|-----|-------------|------|-------------|-------|-----|
| Industri       | 49           | 1   | 89          | 160  | 29          | 328   | 48  |
| Transportation | 191          | -   | 0           | -    | 0           | 191   | 31  |
| Household      | 40           | 14  | 0           | -    | 31          | 85    | 13  |
| Commercial     | 7            | 1   | 0           | -    | 19          | 28    | 4   |
| Others         | 25           | -   | -           | -    | -           | 25    | 5   |
| Total          | 312          | 16  | 90          | 160  | 79          | 657   | 100 |

Source: Indonesian Energy Outlook 2009, Data and Information Center, MEMR

Indonesia is endowed with various types of energy resources, fossil energy resources as well as new and renewable energy resources. As an overview, among of the fossil resources, coal has the longest reserves to production ratio, i.e. 82 years, while new and renewable resources are also potential. The utilization level of the new and renewable resources, however, is still low. This indicates that these resources could be further developed to supply future energy demand. The distribution of location of each resource and the corresponding infrastructure of the resources (mostly outside Java) are mismatched with energy demand center (mostly Java). Therefore, the development of energy resources in Indonesia considers this situation.

To supply the domestic energy demand, Indonesia uses different types primary energy sources, i.e. oil, natural gas, coal, hydro, geothermal, and biomass. Among these primary energy sources, the dominant energy source is fossil fuels. As an overview, the primary energy supply mix (excluding biomass) in 2005: oil (48.5%), natural gas (27.1%), coal (18.7%), hydro (3.4%) and geothermal (2.3%). In terms of annual average growth during 2000 - 2005, coal outpaced other fuels: coal 14.3%, oil 2.6%, and gas 0.6%. According to the Indonesian National Energy Policy (Presidential Regulation No.5/2006), it is projected that coal, natural gas, and renewable energy will replace oil in the future so that in 2025 the primary energy supply mix will become: coal 33%, natural gas 30%, oil 20%, and renewable and other new energy 17%. It should be noted that the primary consideration in targeting the energy supply mix is energy security concerns.

Concerning the biomass energy, it is primarily consumed in rural residential and agro industries. During 2000 – 2005, the biomass consumption is relatively constant, in the order of 270 MMBOE per annum. A relatively small fraction ( $\pm 15\%$ ) of the biomass is used in industrial captive power and heat generations, i.e. in pulp paper industry, wood processing, and palm oil mills. With regard to power sector, the primary energy supply is dominated by coal, followed by natural gas and oil. As an overview, energy mix in the power sector in 2005: coal (40.4%), oil (26.2%), hydro (13.3%), natural gas (11.2%), geothermal (8.9%).

Elasticity of energy in Indonesia is relatively high compared to other countries (1.84 compared to around 1.0). This implies that more energy is needed by Indonesia to increase per unit of GDP. One of national energy policy objectives is to reduce elasticity to less than 1 by 2025. The objective can largely be achieved through energy efficiency and the national policy of energy efficiency (EE) is already established. This policy is also supported by EE programmes and measures.

### **1.4.2 Industrial Sector**

The growth of Industrial sector, particularly for mineral, chemical and metal products in the period of 2000-2005 was quite slow and even negative for some industries (Table 1.4). Since 1998, after the global economic crisis, investment on these types of industries has been very limited. It is estimated that until 2025 these types of industries may not grow.

**Table 1.4. Growth and production of industrial products (000 ton) in the period 2000-2005**

| Year                         | 2000      | 2001      | 2002      | 2003      | 2004      | 2005      | Growth (%/yr) |
|------------------------------|-----------|-----------|-----------|-----------|-----------|-----------|---------------|
| <b>Mineral Products</b>      |           |           |           |           |           |           |               |
| Cement production            | 27,800.00 | 31,100.00 | 31,360.00 | 30,600.00 | 33,000.00 | 33,920.00 | 3.26          |
| Clinker Export               | 3,552.00  | 3,707.00  | 3,791.00  | 4,270.00  | 4,680.00  | 3,407.24  | -0.62         |
| Lime production              | 4,917.53  | 9,382.15  | 2,770.10  | 2,744.89  | 2,820.01  | 2,827.93  | -8.21         |
| Glass production             | 1,700.00  | 1,301.71  | 1,435.26  | 1,342.25  | 1,460.50  | 1,499.85  | -2.29         |
| Ceramic Production           | 13.32     | 2.16      | 2.16      | 4.43      | 1.31      | 0.50      | -53.68        |
| <b>Chemical products</b>     |           |           |           |           |           |           |               |
| Ammonia production           | 4,785.00  | 4,406.80  | 4,771.42  | 4,860.41  | 4,546.26  | 5,125.31  | 1.19          |
| HN03 production              | 23.04     | 13.77     | 14.99     | 14.99     | 14.99     | 14.99     | -8.32         |
| Silicon Carbide              | 27.40     | 18.13     | 19.73     | 19.73     | 19.73     | 19.73     | -6.16         |
| Calcium Carbide              | 22.45     | 70.17     | 76.40     | 76.40     | 20.23     | 20.23     | -0.78         |
| CaC2 in Asetilena Production | 19.02     | 19.02     | 12.10     | 11.64     | 12.89     | 12.83     | -7.07         |
| Methanol                     | 794.47    | 931.36    | 785.03    | 792.33    | 787.91    | 845.54    | 1.03          |
| Ethylene                     | 499.32    | 397.58    | 428.27    | 476.28    | 465.05    | 487.22    | -0.44         |
| Ethylene Dichloride          | 760.95    | 787.89    | 782.03    | 798.91    | 799.99    | 727.99    | -0.71         |
| Carbon Black                 | 94.52     | 98.18     | 90.83     | 95.65     | 110.00    | 123.00    | 4.65          |
| <b>Metal Products</b>        |           |           |           |           |           |           |               |
| Basic Oxygen Furnace         | 27.40     | 18.13     | 19.73     | 19.73     | 19.73     | 19.73     | -6.16         |
| Pig Iron                     | 0.29      | 0.23      | 2.40      | 3.21      | 2.63      | 33.03     | 78.36         |
| Sinter                       | 1,355.69  | 1,917.86  | 859.42    | 2,211.91  | 2,425.27  | 2,269.83  | 8.28          |
| Al production                | 240.00    | 240.00    | 240.00    | 240.00    | 240.00    | 240.00    | 0.00          |
| PFC                          | 0.07      | 0.07      | 0.07      | 0.07      | 0.07      | 0.07      | 0.00          |
| Lead Production              | 27.40     | 18.13     | 19.73     | 19.73     | 19.73     | 19.73     | -6.16         |
| Zinc Production              | 71.87     | 98.74     | 55.28     | 56.62     | 40.97     | 61.82     | -2.61         |

Source: BPS (2007)

### 1.4.3 Forestry Sector

In Indonesia, the state land is divided into two areas: forest area and non-forest area. Forest area is not always covered by forest. Similarly, non-forest area does not necessarily mean it has no forest. Non-forest area is commonly referred to as "area for other uses" (i.e., non-forest activities such as for agriculture plantation, settlement, road etc.; abbreviated as APL). Total forest area is approximately 133.6 million ha and non-forest area is 54.3 million ha (Baplan, 2008). Based on its function, about 15% of forest area is categorized as conservation forest (HK), 22% as protection forest (HL), 46% as production forest (HP) and 17% as convertible production forest (HPK). This convertible production forest can be released as APL in the future. Based on 2007 satellite assessments, forest area covered by forest was about 85.9 million ha (64%), 39.1 million ha were without forest and 8.5 million ha were unidentified due to cloud cover (BAPLAN, 2008).


In non-forest area (APL), the forested land was estimated to be about 7.96 million ha (15%; Table 1.5).

**Table 1.5 Forested and non-forested land in forest and non-forest area in 2007 (thousand ha)**

|                   | Forest Area (x 000 ha) |               |               |               | APL           |
|-------------------|------------------------|---------------|---------------|---------------|---------------|
|                   | HK                     | HL            | HP            | HPK           |               |
| Forested land     | 14,365                 | 22,102        | 38,805        | 10,693        | 7,960         |
| Non-forested land | 4,009                  | 5,622         | 18,404        | 11,057        | 44,163        |
| Unidentified      | 1,502                  | 2,328         | 3,706         | 981           | 2,216         |
| <b>Total</b>      | <b>19,876</b>          | <b>30,052</b> | <b>60,915</b> | <b>22,732</b> | <b>54,339</b> |

Source: Baplan (2008)

From satellite interpretation, rate of forest loss between 2000 and 2005 reached about 1.1 million ha per year (BAPLAN, 2008). Most of deforestation mainly occurred in HP and the followed by APL, HPK, HK and HL (Figure 1.5; MoF 2008). The rate of deforestation is expected to increase in the future, particularly in open access forest areas. Open access area is defined as forest areas that have not been granted to concessionaires and have no on-site agencies responsible for managing the areas. At present only conservation forests have on-site management agency (called as National Park Agencies). In one of its priority policies, the Ministry of Forestry will establish a Forest Management Unit (FMU) in all forest areas. Thus, all forest areas in the country will have on-site management unit. It is estimated that the number of FMU needed for all over Indonesia is about 700 units. With the current budget, the Government of Indonesia is only able to establish about 210 FMUs in the next 15 years. The establishment of FMU will therefore be prioritized in regions with high deforestation risk.



**Figure 1.5 Fraction of forest loss in forest area and non-forest area**

There are many factors causing deforestation and forest degradation in Indonesia. They include (i) forest fire, (ii) illegal logging, (iii) forest encroachment, (iv) forest conversion for establishment of agriculture plantation, transmigration areas, and establishment of new districts, (v) development of new rice fields, and (vi) large-scale mining activities. The first three factors are defined as unplanned deforestation and the remaining factors are defined as planned deforestation. To reduce the rate of unplanned deforestation and forest degradation, and reduce the area of degraded land and forest, the In Government of Indonesia has established five priority policies as follows: (i) combating illegal logging and forest fire, (ii) restructuring forestry sector industries, including enhancement of plantation development, (iii) forest rehabilitation and conservation, (iv) promoting sustainable forest area, and (v) strengthening economies of local communities. Implementation of these policies has reduced the rate of deforestation quite significantly (Figure 1.6). However, since a decentralized governance system is implemented, the desire to bloom and to form new autonomous regions is nearly unstoppable. From 2007 to 2008, approximately 43 new districts have been formed under the law (<http://www.setneg.go.id/>). In addition, several district-level governments have requested the to release forest area for expanding the area of their districts. This planned deforestation may become a big threat for the future of Indonesian forest areas.




Figure 1.6 Rate of deforestation in Indonesia (MoF, 2007)


#### 1.4.4 Agriculture Sector

During the period from 2005-2009, agricultural development continued to record varying success. One notable success was that Indonesia achieved rice self-sufficiency since 2008, and self-sufficiency in maize consumption of households occurred in 2008. Stable production of rice, which is the main food is helping to stabilize domestic food prices, so that Indonesia can avoid a food crisis that struck many countries during this period. However, the long-term trend indicates a steady decline in sawah (paddy field) area, as much as 14,000 ha annually (Figure 1.7). This is primarily due to conversions of paddy field to urban and settlement developments (Agus *et al.* 2006). If this continues, rice self-sufficiency will not sustain. On the other hand, plantation

area has drastically increased at a rate of 571,000 ha annually. The areas of annual upland and idle lands (including Imperata grass and bush lands) fluctuate, but there is an increasing trend for these two land use systems over the last six years. The increase in plantation, upland farming and idle lands postulates a decreasing forest area.


Rapid increase in the area of agriculture plantation is mainly due to the high growth of palm oil plantation. The area of palm oil increased exponentially over the last two decades (Figure 1.10). Between 1994 and 2008, the average increase was 12.3% annually from about 0.6 to 6.6 million ha. The rapid increase in the palm oil plantations is driven by the demand increase in the domestic and international markets, including the demand for bio-diesel. Areas of cacao and coffee plantation have also increased, but not as drastically as that for palm oil.

To increase the production of main agriculture commodities, particularly rice and palm oil, Indonesia has targeted to expand the palm oil plantation as much as 1.375 million ha between 2010 and 2015 (DG of Estate Crop, 2008). The target provinces for the establishment of palm oil plantations are East, West and Central Kalimantan, Jambi and South Sumatra. These five provinces account for 78% of the total area. To secure rice production in the future, Indonesia also plans to have 15 million ha of land permanently allocated for cropland. At present, the total area of paddy field (sawah) is approximately 7.8 million ha; thus, an additional 7.2 million ha is needed to meet this target. Based on Ministry of Agriculture estimate (IAARD, 2007), total suitable area for sawah is 8.3 million ha in which 3 million ha is swamp land (including mineral wetland and peatland) and the other 5.3 million ha is convertible upland. However, the area of potential suitable land changes rapidly because of competition in land uses for housing and urban developments. Thus, the currently available lands for sawah may be less (Agus *et al.* 2006).



**Figure 1.7 Development of the main agricultural systems from 1986 to 2006 (Data taken from Central Bureau of Statistics)**





**Figure 1.8 Development of the main plantation areas from 1986 to 2006**  
(Data taken from Central Bureau of Statistics)

### 1.4.5 Water Sector

In general, the Indonesian territory has 6% of the world's fresh water reserve or approximately 21% of water reserve in the Asia-Pacific region. The availability of water in Indonesia is quite high as indicated by the high level of rainfall and the potential availability of surface water and groundwater. However, many areas in Indonesia have suffered from difficulties of usable water availability in recent years. Java (with its high population and high industrial activities), Bali and East Nusa Tenggara already have water deficits (Directorate General of Water Resource, 2007). The total water demands of the country are currently for supporting irrigation, domestic, and municipal and industrial usages of 1,074 m<sup>3</sup>/sec; however, the low flow available during a normal climatic year are only about 790 m<sup>3</sup>/sec or approximately 76% of the total water demand. This deficit will increase as population and economic activities increase.

There are a number of factors causing the reduction of water quantity and quality in Indonesia. The first factor is the degradation of the carrying capacity of upstream areas of water catchments as a result of the uncontrolled clearing of forests. This has been indicated by the increase in the number of critical catchments areas of river basins from 22 river basins in 1984 to 39 river basins in 1992 and an increase to 62 catchment areas in 1998 (MoE, 2007). The second factor is uncontrolled land clearing within flooding areas, water catchment areas and riverbanks that has resulted in reduced infiltration capacity, changes in river morphology, and reduced carrying capacity of the streams, thus expanding the risk and increasing the frequency of flooding. The third factor is the uncontrolled abstraction of water that also causes increased saltwater intrusion and land subsidence. The fourth factor is the degradation of riverbeds in Java, Bali and West Nusa Tenggara due to exploitation of sand that causes infrastructure and structural damage along rivers. The fifth and final factor is the increased sedimentation of riverbeds resulting from household solid waste disposal and mining.

Efforts for managing water resources have been carried out by the Government through instrumentation of regulation and programs executed by related sectors. In general, the efforts focused into two activities: the conservation of water and control of water pollution.

#### 1.4.6 Coastal and Marine Sector

Indonesia is an archipelago state that has approximately 5.8 million km<sup>2</sup> of ocean area and 81,000 km coastline (the longest coastline in the world), with more than 17,500 islands. Coastal areas, small islands, marine life and fisheries play an important role in supplying food energy, supporting natural cycles, and regulating the global climate. From an economic standpoint, Indonesia's fishery resources equate to 6.65 million ton per annum that consist of 4.35 million ton in territorial and 2.3 ton in the Executive Economic Zone (MoE, 2007). About 140 million people or 60% of the total population lives within 50 km of the shoreline, spread across 42 cities and 182 districts.

Indonesia is also regarded as one of the world's mega centers of biodiversity for its extraordinarily high levels of diversity; each coral reef area containing at least 2,500 species of fish, although, at present, but only 30% of these reefs are in good or excellent condition. Due to the richness of these reef areas, the coastal zone policy and management decisions made by the country will have a major impact on the global heritage of coral reef diversity (Bryant *et al.*, 1998).

However, in several areas the oceans have been damaged due to non-environmentally friendly practices. The coral reef damage level in Indonesia reached 40% - 24% of all reefs are considered to be in damaged and medium damaged condition with only 6% remaining in very good condition. It has been estimated that 90% of Indonesia's coral reefs have been damaged by a combination of non-environmentally friendly fish-catching practices, over-fishing, sedimentation and land-based pollution, and coral mining (MoE, 2007).

Similar conditions are also observed in Indonesia's mangrove ecosystems. In 1993, Indonesia's mangrove forest was noted to have an area of 3.7 million ha, dispersed throughout the Indonesian coast, starting from the Island of Sumatra, Kalimantan, Java, Sulawesi, West Nusa Tenggara, East Nusa Tenggara, all of Maluku Islands, and in Papua. However, in 2005, the remaining mangrove forest was only 1.5 million ha. The mangrove forest has significantly decreased between 2% (the lowest) in East Java and 100% (the highest) in East Nusa Tenggara. The causes of disappearance of mangrove forests are the illegal cutting of mangrove wood, area clearing for shrimp breeding and also wet and dry farms (MoE, 2007).

Sea grasses are common throughout the tropical and temperate coastal waters of the Asia and Pacific Region. The Center for Oceanography Research of the Indonesian Institute of Science (LIPI) reported that 12 species of seagrass lived in Indonesia's waters in 2006 (MoE, 2007). However, 20-25% of seagrass areas in Indonesia have been damaged by a combination of coastal development, elevated sedimentation levels, destructive fishing methods and land-based pollution, thermal discharge, petroleum product spills, and dredge and fill operations.

Many initiatives are already carried by the Government and environmental non-governmental organizations (NGOs) related to coastal and marine management such as the Coral Reef Rehabilitation and Management Program (COREMAP) program, including mapping priority areas for rehabilitation and utilization in marine and coastal areas, and mangrove rehabilitation management as well as the Coral Triangle Initiative. This initiative conserves a marine region that spans those parts of Indonesia, Malaysia, New Guinea, the Philippines, the Solomon Islands, and Timor Leste with at least 500 species of reef-building corals; provides incomes and food security more than 120 million people living in this area.



## CHAPTER II

# National GHG Inventory

## 2.1 Introduction

The Indonesian National Greenhouse Gas Inventory (NGHGI) includes estimates of emissions by source and sink for the period 2000-2005. It was drawn up in line with Articles 4 and 12 of the United Nations Framework Convention on Climate Change (UNFCCC) and the Guidelines for National Communications of non-Annex I Parties of the UNFCCC, adopted in decision 17/CP.8, which state that non-Annex I Parties should include information in a national inventory of anthropogenic emissions by source and absorption by sinks of all greenhouse gases (GHGs) not controlled by the Montreal Protocol, within the limits of their possibilities, using the methodologies promoted and approved by the Conference of Parties (COP).


The calculations of GHG emissions reported in the Second National Communication (SNC) were made for the six emissions categories defined by the Intergovernmental Panel on Climate Change (IPCC): Energy, Industrial Processes, Solvents, Agriculture, Land Use Change and Forestry and Waste. The NGHGI 2000-2005 reports on the three main GHGs included in Appendix A of the Kyoto Protocol: carbon dioxide (CO<sub>2</sub>), methane (CH<sub>4</sub>), and nitrous oxide (N<sub>2</sub>O). Some information is also included on three other minor gases (CO, NO<sub>x</sub> and perfluorocarbons [PFC]). The emissions estimates for each emissions category were created thanks to the dedicated work of specialists from the Ministry of Energy and Mineral Resources, the Ministry of Transportation, the Ministry of Forestry, the Ministry of Agriculture, the Ministry of Environment, Bandung Institute of Technology and Bogor Agriculture University.

## 2.2 Institutional Arrangements

The development of the NGHGI was conducted by the Working Group on Inventory assisted by experts and a technical team from two universities (Bandung Institute of Technology and Bogor Agriculture University), as well as two research agencies (Forest Research and Development Agency and the Indonesian Agency for Agriculture Research and Development). The members of the working group came from various government institutions concerned with energy and mining, industry, transportation, waste, agriculture and forestry. The working group was officially recognized through the Environment Minister Decree and worked under the coordination of the Deputy of Nature Conservation Enhancement & Environmental Destruction Control of the Ministry of Environment (Figure 2.1).

## 2.3 Description of the Process of Preparing the Inventory

Preparation of the NGHGI 2000-2005 was carried out in five phases: (1) beginning, (2) development, 3) sectoral consultation, 4) writing up the report and external review and 5) revision and publication, as outlined below:



**Figure 2.1 Structure of the institutional arrangements for developing NGHGI 2000-2005**

1. **Beginning:** In this phase, a meeting of the working group and inventory experts was called to discuss the methodologies and good practices to follow in the preparation of the Inventory. During this phase, the type of activity data and emissions factors required for the Inventory were identified. The Coordinator of the Working Group through the Ministry of Environment officially requested the activity data for each sector and the experts collected this data, as well as the emissions/removal factors from various publications and research agencies.
2. **Development:** After activity data and emissions/removal factors were gathered for each emissions category or sector, the data and information obtained were recorded on spreadsheets. Emissions were then estimated and a report was prepared by the experts; this included the emissions estimates, as well as the analysis carried out and the sources of information consulted. During this phase, coordinators of the working group followed up with requests for information made by the experts and were given support in obtaining additional data that was necessary for further estimates.

3. **Sectoral Consultation.** After the experts completed the Inventory, the results were discussed with the members of the working group to check the estimates, activity data consistency and emissions/removal factors. Through discussion with external experts from different sectors, the levels of uncertainty in activity data and emission factors were determined. Emissions trends were estimated by emissions category and by gas, and both general and category uncertainties were estimated.
4. **Writing up the Report and External Review.** After the consultation with sectors, the final inventory report was made up according to the format by the inventory experts. International experts through the United Nations Development Programme (UNDP) Country Office reviewed the reports, along with all spreadsheets of the inventory for all sectors. Inputs and comments from the external reviewers were used for the improvement of the inventory.
5. **Revision and Publication.** After the completion, the Ministry of Environment submitted the report to relevant ministries for final approval. The final adjustments and corrections were made to the document if errors were found.

## 2.4 Overview of Source and Sink Category Emission Estimates for year 2000

**Methodology.** From the Beginning Phase, the working group and the experts decided to adopt the 2006 IPCC Guidelines (GL) using Tier 1 and Tier 2 even though the adoption of this methodology was not in line with the Guidelines for National Communications of non-Annex I Parties of the UNFCCC, adopted in decision 17/ CP.8. The UNFCCC Guidelines stated that the non-Annex 1 countries used the revised 1996 IPCC GL for developing their inventory. The adoption of the 2006 IPCC GL was considered for some sectors as it covers some sources which are not included in the revised 1996 IPCC GL. Most sectors used Tier 1, and to some extent sectors adopted Tier 2 where local emissions/removal factors were used.

**National Emissions.** In 2000, total GHG emissions for the three main greenhouse gases (CO<sub>2</sub>, CH<sub>4</sub> and N<sub>2</sub>O) without land use, land use change and forestry (LULUCF) and peat fires reached 556,499 Gg CO<sub>2</sub>e. With the inclusion of LULUCF, total GHG emissions from Indonesia increase significantly to about 1,205,753 Gg CO<sub>2</sub>e (Table 2.1). The GHG emissions (in CO<sub>2</sub> equivalent) were distributed unevenly between the three gases recorded: CO<sub>2</sub> totalled 940,879 Gg, representing 78% of the total; methane (CH<sub>4</sub>) totalled 236,388 Gg (CO<sub>2</sub>e), or 20% of the total; and nitrous oxide (N<sub>2</sub>O) totalled 28,341 Gg (CO<sub>2</sub>e), or 2% of the total. The main contributing sectors were Land Use Change and Forestry, followed by energy, peat fire-related emissions, waste, agriculture and industry (Figure 2.2).

The emissions estimates in the SNC are lower than those reported by a 2007 study by Pelangi Energi Abdi Citra Enviro (PEACE), World Bank and the United Kingdom Department for International Development (DFID), which suggested Indonesia to be the third largest

emitting country globally. The study estimated that the total emissions from Indonesia were approximately 3,014,000 Gg CO<sub>2</sub>, where LUCF contributed about 85% or 2,563,000 Gg CO<sub>2</sub> (twice the SNC estimate above). Indeed, a later study from the World Bank in 2008 suggested that the mean annual CO<sub>2</sub> emissions from LUCF reached up to 2,398,000 Gg, assuming 53% from peat fire, 20% from peat drainage (peat oxidation), 22% from deforestation and 5% from palm oil and timber plantation establishment. Between the SNC and World Bank estimates, a recent study from the National Committee on Climate Change in 2009 suggested that the total GHG emissions from LUCF in 2005 reached 1,880,000 Gg CO<sub>2</sub>e, where approximately 55% was from peat emissions (DNPI, 2009). These large differences in Indonesian emissions estimates thus appear to be mainly due to differences in estimates of LUCF emissions, particularly from peat.

**Table 2.1a Summary of national GHG emissions in 2000**

| No   | Source and sink Categories                                | CO <sub>2</sub> removal (Gg) | CO <sub>2</sub> emission (Gg) | CH <sub>4</sub> (Gg) | N <sub>2</sub> O (Gg) | CO (Gg)         | NO <sub>x</sub> (Gg) | NM VOC (Gg) | SO <sub>x</sub> (Gg) |
|--|---|------------------------------|-------------------------------|----------------------|-----------------------|-----------------|----------------------|-------------|----------------------|
| <b>Total National Emission and Removals</b>    |   | <b>-296,794.38</b>           | <b>1,352,471.68</b>           | <b>11,256.59</b>     | <b>91.42</b>          | <b>2,335.71</b> | <b>85.66</b>         | <b>NE</b>   | <b>NE</b>            |
| <b>1. Energy (without Biomass)</b>             |   |                              | <b>247,522.25</b>             | <b>1,436.89</b>      | <b>10.45</b>          | <b>NE</b>       | <b>NE</b>            | <b>NE</b>   | <b>NE</b>            |
| <b>A</b>                                       | <b>Fuel Combustion Activity</b>                           |                              | <b>240,876.89</b>             | <b>455.51</b>        | <b>10.40</b>          | <b>NE</b>       | <b>NE</b>            | <b>NE</b>   | <b>NE</b>            |
| 1  | Energy production (electricity, heat, oil & gas refining) |                              | 84,011.42                     | 1.89                 | 0.64                  | NE              | NE                   | NE          | NE                   |
| 2  | Manufacturing Industries and Construction                 |                              | 63,032.48                     | 7.39                 | 1.10                  | NE              | NE                   | NE          | NE                   |
| 3  | Transportation  |                              | 55,689.23                     | 14.32                | 2.68                  | NE              | NE                   | NE          | NE                   |
| 4  | Commercial/Intstitutional                                 |                              | 3,320.84                      | 2.14                 | 0.03                  | NE              | NE                   | NE          | NE                   |
| 5  | Residential   |                              | 23,878.82                     | 428.26               | 5.86                  | NE              | NE                   | NE          | NE                   |
| 6  | Non Specified   |                              | 10,944.09                     | 1.50                 | 0.09                  | NE              | NE                   | NE          | NE                   |
| <b>B.</b>                                      | <b>Fugitive Emissions</b>                                 |                              | <b>6,645.36</b>               | <b>981.38</b>        | <b>0.05</b>           | <b>NE</b>       | <b>NE</b>            | <b>NE</b>   | <b>NE</b>            |
| 1  | Solid Fuels   |                              |                               | 17.22                |                       | NE              | NE                   | NE          | NE                   |
| 2  | Oil and Natural Gas                                       |                              | 6,645.36                      | 964.17               | 0.05                  | NE              | NE                   | NE          | NE                   |
| <b>2. Industrial Processes and Product Use</b> |   |                              | <b>40,342.41</b>              | <b>104.44</b>        | <b>0.43</b>           | <b>NE</b>       | <b>NE</b>            | <b>NE</b>   | <b>NE</b>            |
| <b>A.</b>                                      | <b>Mineral</b>  |                              | <b>28,923.82</b>              |                      |                       | <b>NE</b>       | <b>NE</b>            | <b>NE</b>   | <b>NE</b>            |
| 1  | Cement Production   |                              | 25,006.12                     |                      |                       | NE              | NE                   | NE          | NE                   |
| 2  | Lime Production   |                              | 3,688.15                      |                      |                       | NE              | NE                   | NE          | NE                   |
| 3  | Glass Production  |                              | 170.00                        |                      |                       | NE              | NE                   | NE          | NE                   |
| 4  | Other Process Uses of Carbonates                          |                              | 59.55                         |                      |                       | NE              | NE                   | NE          | NE                   |
| <b>B.</b>                                      | <b>Chemical</b>   |                              | <b>9,938.97</b>               | <b>9.54</b>          | <b>0.43</b>           | <b>NE</b>       | <b>NE</b>            | <b>NE</b>   | <b>NE</b>            |
| 1  | Ammonia Production  |                              | 8,092.32                      | NE                   | NE                    | NE              | NE                   | NE          | NE                   |
| 2  | Nitric Acid Production                                    |                              | NE                            | NE                   | 0.21                  | NE              | NE                   | NE          | NE                   |
| 3  | Caprolactam, Glyoxal and Glyoxylic Acid                   |                              | NE                            | NE                   | 0.22                  | NE              | NE                   | NE          | NE                   |
| 4  | Carbide Production  |                              | 21.02                         | 0.32                 |                       | NE              | NE                   | NE          | NE                   |
| 5  | Petrochemical and Carbon Black Production                 |                              |                               |                      |                       |                 |                      |             |                      |
|  | - Methanol  |                              | 305.87                        | 1.83                 | NE                    | NE              | NE                   | NE          | NE                   |
|  | - Ethylene  |                              | 1,122.97                      | 1.50                 | NE                    | NE              | NE                   | NE          | NE                   |
|  | - Ethylene Dichloride                                     |                              | 149.15                        | 0.23                 | NE                    | NE              | NE                   | NE          | NE                   |
|  | - Carbon Black  |                              | 247.64                        | 5.67                 | NE                    | NE              | NE                   | NE          | NE                   |
| <b>C.</b>                                      | <b>Metal</b>  |                              | <b>1,152.98</b>               | <b>94.90</b>         | <b>NE</b>             | <b>NE</b>       | <b>NE</b>            | <b>NE</b>   | <b>NE</b>            |
| 1  | Iron and Steel Production                                 |                              | 623.91                        | 94.90                | NE                    | NE              | NE                   | NE          | NE                   |
| 2  | Ferroalloys Production                                    |                              | 2.39                          | NE                   | NE                    | NE              | NE                   | NE          | NE                   |
| 3  | Aluminium Production                                      |                              | 384.00                        | NE                   | NE                    | NE              | NE                   | NE          | NE                   |
| 4  | Lead Production   |                              | 19.05                         | NE                   | NE                    | NE              | NE                   | NE          | NE                   |
| 5  | Zinc Production   |                              | 123.62                        | NE                   | NE                    | NE              | NE                   | NE          | NE                   |
| <b>D.</b>                                      | <b>Others</b>   |                              | <b>326.65</b>                 | <b>NE</b>            | <b>NE</b>             | <b>NE</b>       | <b>NE</b>            | <b>NE</b>   | <b>NE</b>            |
| 1  | Lubricant Use   |                              | 307.07                        | NE                   | NE                    | NE              | NE                   | NE          | NE                   |
| 2  | Paraffin Wax Use  |                              | 19.57                         | NE                   | NE                    | NE              | NE                   | NE          | NE                   |
| <b>3. Solvent and Other Product Use</b>        |   |                              | <b>NE</b>                     | <b>NE</b>            | <b>NE</b>             | <b>NE</b>       | <b>NE</b>            | <b>NE</b>   | <b>NE</b>            |
| <b>4. Agriculture</b>                          |   |                              | <b>2,178.30</b>               | <b>2,419.06</b>      | <b>72.39</b>          | <b>2,294.68</b> | <b>84.67</b>         |             |                      |
| A  | Enteric Fermentation                                      |                              |                               | 607.89               |                       |                 |                      |             |                      |
| B  | Manure Management   |                              |                               | 79.69                |                       |                 |                      |             |                      |



| No                                     | Source and sink Categories   | CO <sub>2</sub> removal (Gg) | CO <sub>2</sub> emission (Gg) | CH <sub>4</sub> (Gg) | N <sub>2</sub> O (Gg) | CO (Gg)      | NO <sub>x</sub> (Gg) | NM VOC (Gg) | SO <sub>x</sub> (Gg) |
|--|--|------------------------------|-------------------------------|----------------------|-----------------------|--------------|----------------------|-------------|----------------------|
| C                                      | Rice Cultivation   |                              |                               | 1,660.03             | NE                    |              |                      |             |                      |
| D                                      | Agriculture Soils  |                              |                               | NE                   |                       |              |                      |             |                      |
| 1                                      | Direct N <sub>2</sub> O Soils  |                              |                               | NE                   | 50.11                 |              |                      |             |                      |
| 2                                      | Indirect N <sub>2</sub> O Soils  |                              |                               | NE                   | 18.34                 |              |                      |             |                      |
| 3                                      | Direct N <sub>2</sub> O from manure  |                              |                               | NE                   | 0.40                  |              |                      |             |                      |
| 4                                      | Indirect N <sub>2</sub> O from manure  |                              |                               | NE                   | 0.11                  |              |                      |             |                      |
| E                                      | Prescribed Burning of Savanna/grassland  |                              |                               | 24.05                | 2.20                  | 679.78       | 40.79                |             |                      |
| F                                      | Prescribed Burning of Agriculture Residues                                     |                              |                               | 47.39                | 1.23                  | 1,614.90     | 43.88                |             |                      |
| G                                      | Others   |                              |                               |                      |                       |              |                      |             |                      |
| 1                                      | Liming   |                              | 218.02                        | NO                   | NE                    | NO           | NO                   | NO          | NO                   |
| 2                                      | Urea Fertilization   |                              | 1,960.28                      | NO                   | IE                    | NO           | NO                   | NO          | NO                   |
| <b>5. Land Use Change and Forestry</b> |  | <b>-296,794.38</b>           | <b>1,232,766.22</b>           | <b>2.68</b>          | <b>0.08</b>           | <b>41.04</b> | <b>0.99</b>          |             |                      |
| A                                      | Changes in forest and other woody biomass stocks                               | -215,154.48                  |                               |                      |                       |              |                      |             |                      |
| B                                      | Forest and grassland conversion  |                              | 729,655.23                    | NE                   | NE                    | NE           | NE                   |             |                      |
| C                                      | Abandonment of croplands, pastures, plantation forests, or other managed lands | -81,639.89                   |                               |                      |                       |              |                      |             |                      |
| D                                      | CO <sub>2</sub> emissions and removals from soils                              | NE                           | 216,312.51                    |                      |                       |              |                      |             |                      |
| E                                      | Others:  |                              |                               |                      |                       |              |                      |             |                      |
|  | - Forest Burning   |                              |                               | 2.68                 | 0.08                  | 41.04        | 0.99                 |             |                      |
|  | - Peat Fire*   |                              | 172,000.00                    | NE                   | NE                    | NE           | NE                   |             |                      |
| <b>6. Waste</b>                        |  |                              | <b>1,662.49</b>               | <b>7,293.52</b>      | <b>8.07</b>           | <b>NE</b>    | <b>NE</b>            |             |                      |
| A1                                     | Unmanaged Waste Disposal Sites   |                              | NE                            | 612.83               | NE                    | NE           | NE                   | NE          |                      |
| A2                                     | Unmanaged Dumpsite   |                              | NE                            | 249.72               | NE                    | NE           | NE                   | NE          |                      |
| B                                      | Biological Treatment of Solid Waste  |                              | NE                            | 4.42                 | 0.33                  | NE           | NE                   | NE          | NE                   |
| C                                      | Open Burning Waste   |                              | 1,662.49                      | 62.34                | 1.44                  | NE           | NE                   | NE          | NE                   |
| D1                                     | Domestic Wastewater Treatment and Discharge                                    |                              |                               | 459.59               | 6.30                  | NE           | NE                   | NE          |                      |
| D2                                     | Industrial Wastewater Treatment and Discharge                                  |                              |                               | 5,904.63             | NE                    | NE           | NE                   | NE          |                      |
| <b>7. Others</b>                       |  |                              | <b>176,765.69</b>             | <b>NE</b>            | <b>NE</b>             | <b>NE</b>    | <b>NE</b>            | <b>NE</b>   | <b>NE</b>            |
|  | International bunkers  |                              | NE                            | NE                   | NE                    | NE           | NE                   | NE          | NE                   |
|  | Aviation   |                              | NE                            | NE                   | NE                    | NE           | NE                   | NE          | NE                   |
|  | Marine   |                              | NE                            | NE                   | NE                    | NE           | NE                   | NE          | NE                   |
|  | Biomass  |                              | 176,765.69                    |                      |                       |              |                      |             |                      |


Note: \* Emissions from peat fire are from van der Warf *et al.* (2007). Shaded cells do not require entries. NO: Not Occurring; NE: Not estimated; NA: Not Applicable; IE: Including Elsewhere; C: Confidential

**Table 2.1b. Summary of national GHG emissions in 2000 (in Gg)**

|           |   | HFCs   |         |         |         |        | PFCs |        | SF6 |
|-----------|---|--------|---------|---------|---------|--------|------|--------|-----|
|           |   | HFC-32 | HFC-125 | HFC 124 | HFC 143 | Others | CF4  | Others |     |
| <b>1</b>  | <b>Energy</b>   |        |         |         |         |        |      |        |     |
| <b>A</b>  | <b>Fuel Combustion Activity</b>                           |        |         |         |         |        |      |        |     |
| 1 A 1     | Energy production (electricity, heat, oil & gas refining) |        |         |         |         |        |      |        | NE  |
| 1 A 2     | Manufacturing Industries and Construction                 |        |         |         |         |        |      |        |     |
| 1 A 3     | Transportation  |        |         |         |         |        |      |        |     |
| 1 A 4 a   | Commercial/Institutional                                  |        |         |         |         |        |      |        |     |
| 1 A 4 b   | Residential   |        |         |         |         |        |      |        |     |
| 1 A 5     | Non Specified   |        |         |         |         |        |      |        |     |
| <b>B.</b> | <b>Fugitive Emissions</b>                                 |        |         |         |         |        |      |        |     |
| 1 B 1     | Solid Fuels   |        |         |         |         |        |      |        |     |
| 1 B 2     | Oil and Natural Gas                                       |        |         |         |         |        |      |        |     |
| <b>2</b>  | <b>Industrial Processes and Product Use</b>               |        |         |         |         |        |      |        |     |
| <b>A.</b> | <b>Mineral</b>  |        |         |         |         |        |      |        |     |
| 2 A 1     | Cement Production   |        |         |         |         |        |      |        |     |
| 2 A 2     | Lime Production   |        |         |         |         |        |      |        |     |
| 2 A 3     | Glass Production  |        |         |         |         |        |      |        |     |
| 2 A 4     | Other Process Uses of Carbonates                          |        |         |         |         |        |      |        |     |

|           |   | HFCs   |         |         |         |        | PFCs |        | SF6 |
|-----------|---|--------|---------|---------|---------|--------|------|--------|-----|
|           |   | HFC-32 | HFC-125 | HFC 124 | HFC 143 | Others | CF4  | Others |     |
| <b>B.</b> | <b>Chemical</b>                                   |        |         |         |         |        |      |        |     |
| 2 B 1     | Ammonia Production                                |        |         |         |         |        |      |        |     |
| 2 B 2     | Nitric Acid Production                            |        |         |         |         |        |      |        |     |
| 2 B 4     | Caprolactam, Glyoxal dan Glyoxylic Acid           |        |         |         |         |        |      |        |     |
| 2 B 5     | Carbide Production                                |        |         |         |         |        |      |        |     |
| 2 B 8     | Petrochemical and Carbon Black Production         |        |         |         |         |        |      |        |     |
|           | - Methanol  |        |         |         |         |        |      |        |     |
|           | - Ethylene  |        |         |         |         |        |      |        |     |
|           | - Ethylene Dichloride                             |        |         |         |         |        |      |        |     |
|           | - Carbon Black                                    |        |         |         |         |        |      |        |     |
| <b>C.</b> | <b>Metal</b>                                      |        |         |         |         |        | NE   | 0.020  |     |
| 2 C 1     | Iron and Steel Production                         |        |         |         |         |        | NE   | NE     |     |
| 2 C 2     | Ferroalloys Production                            |        |         |         |         |        | NE   | NE     |     |
| 2 C 3     | Aluminium Production                              |        |         |         |         |        | IE   | 0.020  |     |
| 2 C 5     | Lead Production                                   |        |         |         |         |        | NE   | NE     |     |
| 2 C 6     | Zinc Production                                   |        |         |         |         |        | NE   | NE     |     |
| <b>D.</b> | <b>Others</b>                                     |        |         |         |         |        |      |        |     |
| 2 D 1     | Lubricant Use                                     |        |         |         |         |        |      |        |     |
| 2 D 2     | Paraffin Wax Use                                  |        |         |         |         |        |      |        |     |
| <b>3</b>  | <b>Solvent and Other Product Use</b>              |        |         |         |         |        |      |        |     |
| <b>4</b>  | <b>Agriculture</b>                                |        |         |         |         |        |      |        |     |
| 4 A 1     | Enteric Fermentation                              |        |         |         |         |        |      |        |     |
| 4 A 2     | Manure Management                                 |        |         |         |         |        |      |        |     |
| 4 C 1 b   | Biomass Burning CL                                |        |         |         |         |        |      |        |     |
| 4 C 1 c   | Biomass Burning GL                                |        |         |         |         |        |      |        |     |
| 4 C 2     | Liming  |        |         |         |         |        |      |        |     |
| 4 C 3     | Urea Fertilization                                |        |         |         |         |        |      |        |     |
| 4 C 4     | Direct N <sub>2</sub> O Soils                     |        |         |         |         |        |      |        |     |
| 4 C 5     | Indirect N <sub>2</sub> O Soils                   |        |         |         |         |        |      |        |     |
| 4 C 6 a   | Direct N <sub>2</sub> O from manure               |        |         |         |         |        |      |        |     |
| 4 C 6 b   | Indirect N <sub>2</sub> O from manure             |        |         |         |         |        |      |        |     |
| 4 C 7     | Rice Cultivation                                  |        |         |         |         |        |      |        |     |
| <b>5</b>  | <b>Land Use Change and Forestry</b>               |        |         |         |         |        |      |        |     |
| A         | Changes in forest and other woody biomass stocks  |        |         |         |         |        |      |        |     |
| B         | Forest and grassland conversion                   |        |         |         |         |        |      |        |     |
| C         | Abandonment of managed lands                      |        |         |         |         |        |      |        |     |
| D         | CO <sub>2</sub> emissions and removals from soils |        |         |         |         |        |      |        |     |
| E         | Others:   |        |         |         |         |        |      |        |     |
|           | - Forest Burning                                  |        |         |         |         |        |      |        |     |
|           | - Peat Fire*                                      |        |         |         |         |        |      |        |     |
| <b>6</b>  | <b>Waste</b>                                      |        |         |         |         |        |      |        |     |
| 6 A 2 a   | Unmanaged Waste Disposal Sites                    |        |         |         |         |        |      |        |     |
| 6 A 2 b   | Unmanaged Dumpsite                                |        |         |         |         |        |      |        |     |
| 6 B       | Biological Treatment of Solid Waste               |        |         |         |         |        |      |        |     |
| 6 C 2     | Open Burning Waste                                |        |         |         |         |        |      |        |     |
| 6 D 1     | Domestic WWT and Discharge                        |        |         |         |         |        |      |        |     |
| 6 D 2     | Industrial WWT and Discharge                      |        |         |         |         |        |      |        |     |
| 7         | Others (please specify)                           |        |         |         |         |        |      |        |     |
|           | Memo Items  |        |         |         |         |        |      |        |     |
|           | International Bunkers                             |        |         |         |         |        |      |        |     |
|           | Aviation  |        |         |         |         |        |      |        |     |
|           | Marine  |        |         |         |         |        |      |        |     |
|           | CO <sub>2</sub> emissions from biomass            |        |         |         |         |        |      |        |     |

Note: Shaded cells do not require entries. CL: Crops Land. GL: Grass Land. WWT: Wastewater Treatment; NO: Not Occurring; NE: Not Estimated; NA: Not Applicable; IE: Including Elsewhere; C: Confidential



**Figure 2.2 National emissions contributions by sector in 2000**

Inter-annual variation of emissions from peat fires is also very high (Figure 2.3). High emissions normally occurred in El Niño years (1997, 2002, and 2006). The highest estimate was from Hooijer *et al* (2006), which estimated the emissions based on a Borneo hot-spot count and a carbon calculation method used by Page *et al.* (2002), who estimated emissions from peat fires in 1997 El Niño. This approach may give an overestimate, as hot-spot counts in peat lands are not fully proportional to CO<sub>2</sub> emissions, which are governed by additional factors such as the depth and area of burning. Therefore, this relationship may not be generally applicable for the whole of Indonesia and the extrapolation of emission estimates over Indonesia based on limited ground checks in Kalimantan may lead to overestimation.

Van der Werf *et al.* (2008) attempted to improve emission estimates from peat fires using several sources of satellite data with biogeochemical and atmospheric modelling to better understand and constrain peat fire emissions from Indonesia. This resulted in far lower estimates. The NGHGI adopted the study of the van der Werf *et al.* (2007). In 2000, their estimated emissions from peat fires were calculated at approximately 172,000 Gg CO<sub>2</sub>e, while the ten-year average emissions from peat fire was about 466,000 Gg CO<sub>2</sub>e from 1997-2007.




Figure 2.3 Estimate of emissions from peat fires from various studies. Note: Emissions estimates from years beyond the publication date of the reports were made by Aldrian (2008) and Wibowo and Boer (2009)

Emissions by Gas. Without LULUCF, the major source of CO<sub>2</sub> emissions is from fuel combustion, followed by industrial processes, agriculture and waste (Figure 2.4). Total emissions from these four sources was 556,500 Gg (Figure 2.4). With LULUCF and peat fire, the total CO<sub>2</sub> emissions increased to 1,112,879 Gg. As such, LULUCF and peat fire contributed to about 74% of the total CO<sub>2</sub> emissions (Figure 2.4).




Figure 2.4 CO<sub>2</sub> emissions by sectors in 2000

The main source of methane emissions was the waste sector (65%), followed by the agriculture (22%) and energy (13%) sectors. The total methane emissions from all sectors was 236,388 Gg CO<sub>2</sub>e. For nitrous oxide, the main source of emissions was the agriculture sector (79%), followed by the waste (9%) and energy (11%) sectors (Figure 2.5). Total N<sub>2</sub>O emissions was 28,341 Gg CO<sub>2</sub>e.




Figure 2.5 Methane and nitrous oxide emissions in 2000 by sector

## 2.5 Sectoral Emissions

### 2.5.1 Energy

In the 2006 IPCC Guidelines, emissions sources from energy systems are classified in three main categories, namely (i) fuel combustion emissions, (ii) fugitive emission generated from energy production systems (coal mining, oil and gas production, refinery, fuel transportation, etc), and (iii) emissions from transport, injection, and storage of CO<sub>2</sub> (related to carbon capture and storage [CCS]). Only the first two sources were included in the SNC. The source categories being analyzed in the SNC are presented in Figures 2.6 and 2.7.

The calculation of CO<sub>2</sub> emissions applied both sectoral and reference approaches. The two approaches often have different results because the reference approach is a top-down approach using a country's energy supply data and has no detailed information on how the individual fuels are used in each sector. In the case of Indonesia, CO<sub>2</sub> emissions from fuel consumption estimated using the reference approach were slightly higher than those using the sectoral approach in 2004. They differed between 4.3% and 4.7% (Figure 2.8). This can be at least partially attributed to the fact that the energy consumption figure used in the sectoral approach turns out to be less than the consumption figure used in the reference approach.




Figure 2.6 Source of GHG emissions from fuel combustion activities




Figure 2.7 Source categories of fugitive emissions from fuels




Figure 2.8 CO<sub>2</sub> emissions estimates from the energy sector using reference and sectoral approaches

Based on the sectoral approach, CO<sub>2</sub> was the main GHG emitted from the energy sector. It contributes approximately 89% of the total emissions from this category. Methane only contributed approximately 10% and N<sub>2</sub>O contributed less than 1% (Table 2.2).


**Table 2.2 Energy sector emissions from 2000 to 2005 by gas (Gg CO<sub>2</sub>e)**

| Type of Gases               | 2000    | 2001    | 2002    | 2003    | 2004    | 2005    |
|-----------------------------|---------|---------|---------|---------|---------|---------|
| CO <sub>2</sub>             | 247,522 | 274,145 | 296,080 | 303,086 | 341,536 | 339,426 |
| CH <sub>4</sub>             | 30,175  | 29,263  | 28,366  | 27,315  | 26,662  | 26,711  |
| N <sub>2</sub> O            | 3,241   | 3,367   | 3,464   | 3,549   | 3,924   | 3,663   |
| Total (GgCO <sub>2</sub> e) | 280,938 | 306,774 | 327,911 | 333,950 | 372,123 | 369,800 |

Emissions of CO<sub>2</sub> from the energy sector were mainly from fuel combustion, while the contribution of fugitive emissions was almost negligible (Table 2.3). Within the fuel combustion category, the main sources were electricity, petroleum and gas refining (35%), followed by transportation (23%), residential (9%) and manufacturing industries and construction (27%). In the period 2000-2005, CO<sub>2</sub> emissions from fuel combustion showed an increase from 240,877 Gg in 2000 to 333,438 Gg in 2005 (Figure 2.9), which represents a 38% increase, or 6.4% per year.

**Table 2.3 CO<sub>2</sub> emissions from fossil consumption and fugitive emission from 2000-2005 (Gg CO<sub>2</sub>e)**

| Activity   | 2000    | 2001    | 2002    | 2003    | 2004    | 2005    |
|--|---------|---------|---------|---------|---------|---------|
| Energy (Gg CO <sub>2</sub> )                                     | 247,522 | 274,145 | 296,080 | 303,086 | 341,536 | 339,426 |
| Fossil fuel consumption (fuel combustion, (Gg CO <sub>2</sub> )) | 240,877 | 267,782 | 289,615 | 296,688 | 335,546 | 333,438 |
| Fugitive emissions from fuel (Gg CO <sub>2</sub> )               | 6,645   | 6,363   | 6,466   | 6,398   | 5,991   | 5,988   |



**Figure 2.9 CO<sub>2</sub> emissions from the energy sector by source category**



The annual GHG emissions growth rate from fuel consumption in Indonesia was higher than the Indonesian economic growth rate. This differs from other countries where gross domestic product (GDP) is higher than the emissions growth rate (Figure 2.10). Nevertheless, the CO<sub>2</sub> intensity of the country has tended to sharply increase after 1998 and the Asian Financial Crisis (World Bank, 2008).




Figure 2.10 Annual Growth rate of CO<sub>2</sub> emission and GDP of main emitter countries (World Bank)

### 2.5.2 Industry

GHG emissions from the industrial sector are comprised of CO<sub>2</sub>, CH<sub>4</sub>, N<sub>2</sub>O, and PFC. PFC is emitted from the production of aluminum (Al) from a single company. The technology used in the Al smelter is Centre-Worked Prebake (CWPB). Total GHG emissions in the form of CO<sub>2</sub>e were 43,044 Gg in 2000 and increased to 48,733 Gg in 2005 (Table 2.4) – an increase of about 13.2% (2.2% per year). Indeed, there was a significant increase in emissions in 2001.


Table 2.4 GHG emissions from the industrial sector from 2000 to 2005 by gas (in Gg CO<sub>2</sub>e)

| GHG                          | 2000   | 2001   | 2002   | 2003   | 2004   | 2005   |
|------------------------------|--------|--------|--------|--------|--------|--------|
| CO <sub>2</sub>              | 40,342 | 46,469 | 41,928 | 44,073 | 44,054 | 44,838 |
| CH <sub>4</sub>              | 2,423  | 3,022  | 1,452  | 3,450  | 3,787  | 3,572  |
| N <sub>2</sub> O             | 133    | 174    | 190    | 233    | 0      | 178    |
| PFC                          | 145    | 145    | 145    | 145    | 145    | 145    |
| Total (Gg CO <sub>2</sub> e) | 43,044 | 49,810 | 43,716 | 47,902 | 47,985 | 48,733 |

Major GHG emissions from production processes in the industrial sector are mainly from the production of mineral products (~67%), followed by chemical products (22%), metal products (10%) and others (1%) (Table 2.5). By sub-sector, the dominant sources of emissions are from the cement and ammonium industries (Figure 2.11). As evidenced in Figure 2.11, the two industries account for around 60-80% of the total CO<sub>2</sub> eq emissions in Indonesia.

**Table 2.5 GHG emissions from the industrial sector from 2000 to 2005 by source category (in Gg CO<sub>2</sub>e)**

| Source Category | 2000  | 2001  | 2002  | 2003  | 2004  | 2005  |
|-----------------|-------|-------|-------|-------|-------|-------|
| Mineral         | 29153 | 34978 | 30310 | 30105 | 32425 | 32052 |
| Chemical        | 10273 | 9514  | 10118 | 10447 | 9702  | 10982 |
| Metal           | 3291  | 5001  | 3019  | 7029  | 5526  | 5347  |
| Others          | 327   | 318   | 269   | 321   | 332   | 353   |
| Total           | 43044 | 49810 | 43716 | 47902 | 47985 | 48733 |



**Figure 2.11 GHG emissions estimates from production process activities in the industrial sector**

### 2.5.3 Agriculture


Unlike in other sectors, estimates of national emissions from the agriculture sector were based on emissions from districts and provinces. For rice cultivation, data were from the district level, while for livestock, data were from the provincial level. Because of this, variation in biophysical conditions between districts and provinces was taken into account in defining emissions factors.

From the analysis it was found that in 2000, total emissions of the three main GHGs (CO<sub>2</sub>, CH<sub>4</sub> and N<sub>2</sub>O) from this sector reached 75,420 Gg; by 2005, they had increased by 6.3% to 80,179 Gg CO<sub>2</sub>e (Table 2.6). Methane contributed to approximately 65% of the total emissions, while N<sub>2</sub>O contributed 31% and CO<sub>2</sub> contributed 4%.

**Table 2.6 GHG emissions from the agriculture sector from 2000 to 2005 by gas (in Gg CO<sub>2</sub>e)**

| Gas                             | 2000   | 2001   | 2002   | 2003   | 2004   | 2005   |
|---------------------------------|--------|--------|--------|--------|--------|--------|
| CO <sub>2</sub>                 | 2,178  | 3,232  | 3,215  | 3,457  | 3,692  | 3,837  |
| CH <sub>4</sub>                 | 50,800 | 50,677 | 50,833 | 52,547 | 49,342 | 50,670 |
| N <sub>2</sub> O                | 22,441 | 23,592 | 22,982 | 23,825 | 24,828 | 25,672 |
| Total (in Gg CO <sub>2</sub> e) | 75,420 | 77,501 | 77,030 | 79,829 | 77,863 | 80,179 |

By source category, the major emissions were from rice cultivation, direct N<sub>2</sub>O emissions and enteric fermentation from ruminants. These three sources accounted for approximately 83% of the total GHG emissions from this sector (Figure 2.12).




**Figure 2.12 GHG emissions from the agriculture sector from 2000-2005 by source category**

### 2.5.3.1 Livestock

*Enteric Fermentation.* Livestock populations in Indonesia are mostly concentrated in Java, Sumatra and Sulawesi. The highest number of ruminant animals is located in the East Java Province, followed by Nanggroe Aceh Darusalam (NAD; Table 2.7). Enteric fermentation is the most important source of methane emissions from livestock; these emissions are largely from beef cattle (62.8%), buffalo (16.0%) and goat herds (10.4%; Figure 2.13).

**Table 2.7 Livestock population in Indonesia from 2000-2006**

| No. | Species Type             | 2000       | 2001       | 2002       | 2003       | 2004       | 2005       | 2006       |
|-----|--------------------------|------------|------------|------------|------------|------------|------------|------------|
| 1   | Beef Cattle (AU)         | 8,121,691  | 8,069,525  | 8,183,187  | 7,546,157  | 7,547,990  | 7,561,691  | 7,799,932  |
| 2   | Dairy Cattle (AU)        | 265,744    | 260,249    | 268,873    | 280,388    | 273,098    | 542,078    | 276,803    |
| 3   | Buffalo (AU)             | 1,766,248  | 1,710,212  | 1,754,202  | 1,796,835  | 1,751,355  | 1,550,100  | 1,575,649  |
| 4   | Sheep (head)             | 7,414,965  | 7,367,776  | 7,603,892  | 7,774,294  | 8,037,667  | 8,290,477  | 8,947,878  |
| 5   | Goat (head)              | 12,613,108 | 12,198,587 | 12,292,318 | 12,465,046 | 12,592,730 | 13,203,098 | 13,567,972 |
| 6   | Swine (head)             | 5,247,200  | 4,871,896  | 5,237,758  | 5,429,766  | 5,548,999  | 6,043,706  | 5,378,041  |
| 7   | Horse (head)             | 412,919    | 404,162    | 396,697    | 388,599    | 376,378    | 365,478    | 373,981    |
| 8   | Local Chicken (000 head) | 261,132    | 268,786    | 276,023    | 278,068    | 278,966    | 281,431    | 292,122    |
| 9   | Broiler (000 head)       | 534,811    | 624,690    | 849,644    | 855,669    | 785,547    | 820,501    | 808,510    |
| 10  | Layer (000 head)         | 69,703     | 70,127     | 78,024     | 79,168     | 93,497     | 84,688     | 100,236    |
| 11  | Duck (000 head)          | 29,674     | 32,591     | 46,624     | 34,547     | 33,255     | 33,102     | 33,195     |



**Figure 2.13 Distribution of methane emissions from enteric fermentation by species type**

Methane Emissions from Manure Management. Methane emissions from manure management in 2000 were 79.69 Gg. Methane emissions from manure management were dominated by pig farms, which contributed 46.1%, followed by dairy cattle (10.3%) and beef cattle, which contributed the third largest emissions (10.2%) (Figure 2.14). East Java Province was the largest contributor, producing 9.72 Gg (12.2%), followed by Central Java at 8 Gg (12.2%) and North Sumatra at 7.7 Gg (1.9%). Methane emissions from manure management were small compared to enteric fermentation, comprising only 13% of the total emissions from livestock (Figure 2.15).




Figure 2.14 Distribution of methane emissions from manure management by species type




Figure 2.15 Methane emissions from livestock in Indonesia from 2000-2006

N<sub>2</sub>O Emissions from Manure Management. Direct and indirect N<sub>2</sub>O emissions from manure management are presented in Table 2.8. The annual direct N<sub>2</sub>O emissions were 0.399 Gg, while the annual indirect N<sub>2</sub>O emissions were 0.134 Gg. The total N<sub>2</sub>O emissions from manure management was therefore 0.5130 Gg.

**Table 2.8 Direct and indirect N<sub>2</sub>O emissions from manure management**

| Manure Management System | Species Type        | Number of Animal | Annual direct N <sub>2</sub> O emissions (Gg/Year) | Annual indirect N <sub>2</sub> O emissions (Gg/Year) |
|--------------------------|---------------------|------------------|--|--|
| Pasture*                 | Beef Cattle**       | 2,436,507        | -  | -  |
| Daily Spread             | Dairy Cattle        | 265,744          | 0  | 0.000914   |
| Dry Lot                  | Beef Cattle***      | 5,685,184        | 0.155217   | 0.023283   |
|                          | Buffalo             | 1,766,248        | 0.042792   | 0.008558   |
|                          | Sheep               | 7,414,965        | 0.055731   | 0.011146   |
|                          | Goats               | 12,613,108       | 0.118935   | 0.023787   |
|                          | Swine               | 2,623,600        | 0.008427   | 0.001896   |
|                          | Horses              | 412,919          | 0.010372   | 0.002334   |
|                          | Poultry With Litter | Broiler          | 534,810,990  | 0.004555   |
|                          | Layer               | 69,702,890       | 0.000885   | 0.004426   |
| Poultry Without Litter   | Native Chicken      | 261,132,020      | 0.002211   | 0.012159   |
|                          | Duck                | 29,674,120       | 0.000424   | 0.002331   |
| <b>Total</b>             |                     |                  | <b>0.399549</b>                                    | <b>0.113610</b>                                      |


\* The emissions from pasture is calculated in the section called "N<sub>2</sub>O emissions from managed soil"

\*\* Beef cattle in pasture management system is 30% of population

\*\*\* Beef cattle in dry lot management system is 70% of population

### 2.5.3.2 Rice Cultivation

Activity data used to calculate the emissions from rice fields were based on data on the area of rice fields and planting intensity from BPS 2005. Scaling factors for different soil types and correction factors for rice farms based on different water regimes were compiled from various pieces of research in Indonesia. Total CH<sub>4</sub> emissions from Indonesian rice fields in 2000 and 2005 were 1,660 and 1,649 Gg, respectively (Figure 2.16). Methane emissions were calculated based on the proportion of rice field area in each of the years 2000, 2001, 2002, 2003, 2004 and 2005. An increase of 3.6% in 2003 from emissions in year 2000 can be attributed to the growth of rice harvested area, especially in the outer islands. Reduction of total CH<sub>4</sub> emissions by as much as 3.1% in 2004 was due to an El Niño event in the same year, which influenced almost all harvested rice areas in Indonesia.



**Figure 2.16 Methane emissions from rice cultivation from 2000-2005.**

### 2.5.3.3 Agricultural Soils

Liming. Lime consumption data was not available during the data collection period for the SNC. Therefore, the CO<sub>2</sub> emissions from liming were determined from the recommended dose of lime for palm oil plantations in acid sulfate and peat soil (Note: liming on food crops is rarely applied by farmers). Using this method, the CO<sub>2</sub> emissions from liming from 2000 to 2005 are shown in Figure 2.17. Lime consumption in Indonesia increased, which is consistent with the expansion of peatlands for palm oil plantation after the year 2000.




Figure 2.17 CO<sub>2</sub> emissions from liming in agriculture.

**Urea Fertilization.** Adding urea to soils during fertilization leads to a loss of CO<sub>2</sub> that was previously fixed in fertilizer during the industrial production process that manufactured it. Urea (CO(NH<sub>2</sub>)<sub>2</sub>) is converted into ammonium (NH<sub>4</sub><sup>+</sup>), a hydroxyl ion (OH<sup>-</sup>) and bicarbonate (HCO<sub>3</sub><sup>-</sup>), in the presence of water and urease enzymes. Similar to the soil reaction following the addition of lime, the bicarbonate that is formed evolves into CO<sub>2</sub> and water.

Data on urea consumption for years 2000-2005 were derived from the Indonesia's Association of Fertilizer Producers (APPI). CO<sub>2</sub> emissions from urea application in the agricultural sector are shown in Figure 2.18. These emissions are currently increasing, as consumption of urea is increasing at an average rate of 4.5% per year since 2001.




Figure 2.18 CO<sub>2</sub> emissions from urea fertilization from 2000-2005

N<sub>2</sub>O Emissions from Managed Soils – Direct N<sub>2</sub>O Emissions. Urea, ammonium sulphate (AS) and nitrogen, phosphorus and potassium (NPK) are the general types of inorganic nitrogen (N) fertilizers most commonly used in the agricultural sector of Indonesia. Urea and AS are the most common nitrogen-based inorganic fertilizers used in large plantations (APPI, 2008). This type of fertilizer is normally applied for fruits, vegetables and other perennial crops that have high economic value. The concentration of N in urea and AS is 46% and 21%, respectively (Petrokimia Gresik, 2008). However, data on NPK fertilizer consumption is not available. Instead, direct N<sub>2</sub>O emissions on managed soil and flooded rice were calculated from the levels of application of N fertilizer and animal manure. The N fertilizer consumption data are only provided for urea and ammonium sulphate. N<sub>2</sub>O emissions from flooded rice paddies were calculated based on harvested area of rice and from managed soil (i.e., maize, soybean and palm oil). Direct N<sub>2</sub>O emissions from 2000-2005 from managed soil are shown in Figure 2.19. Fluctuation of N<sub>2</sub>O direct emissions from managed soil, as shown in Figure 2.19, can be attributed to the consumption of urea and AS for agriculture in Indonesia. Urea is the main N source used in agriculture, mostly for food crops; however, AS is normally applied in plantation crops such as sugarcane. Some AS is used in rice fields to improve the number of tillers of rice plant.




Figure 2.19 Direct N<sub>2</sub>O emissions from N synthetic fertilizer and manure management applied to soils

**N<sub>2</sub>O Emissions from Managed Soils – Indirect N<sub>2</sub>O Emissions.** Indirect N<sub>2</sub>O emissions from soil were also calculated based on the consumption of N fertilizer (urea, AS and animal manure) in agriculture. The indirect N<sub>2</sub>O emissions from N fertilizer for perennial and estate crops were based on the N fertilizer consumption data from 2000 and the use of animal manure used in agricultural cropping. The resulting N<sub>2</sub>O emissions calculated was approximately 18.34 Gg of N<sub>2</sub>O/year (Figure 2.20).






Figure 2.20 Indirect N<sub>2</sub>O emissions from N synthetic fertilizer and animal manure applied to soils

### 2.5.3.4 Biomass Burning

**Prescribed Savannah Burning (Grassland Burning).** Emissions from grassland burning were calculated based on the 2000-2005 harvested area of upland rice data derived from BPS. Assumptions from expert judgment suggested that 80% of upland rice outside the island of Java was burned and that these areas conducted shifting cultivation; within Java, however, judgment suggested that burning of grassland is not common practice. The emissions of CH<sub>4</sub>, CO, N<sub>2</sub>O and NO<sub>x</sub> from cropland burning are shown in Table 2.9. Results showed that there were not significant differences in emissions from grassland burning between years.

Table 2.9 Distribution of GHG emissions from grassland burning from 2000-2005 (Gg CO<sub>2</sub>e)

| GHG's emission    | 2000     | 2001     | 2002     | 2003     | 2004     | 2005     |
|-------------------|----------|----------|----------|----------|----------|----------|
| CO <sub>2</sub>   | -        | -        | -        | -        | -        | -        |
| CH <sub>4</sub>   | 24.05    | 21.29    | 21.31    | 21.86    | 22.70    | 22.51    |
| N <sub>2</sub> O  | 2.20     | 1.94     | 1.95     | 2.00     | 2.07     | 2.06     |
| CO                | 679.78   | 601.61   | 602.23   | 617.90   | 641.44   | 636.17   |
| NO <sub>x</sub>   | 40.79    | 36.10    | 36.13    | 37.07    | 38.49    | 38.37    |
| CO <sub>2</sub> e | 1,187.05 | 1,048.49 | 1,052.01 | 1,079.06 | 1,118.40 | 1,111.31 |

**Field Burning of Agriculture Residue (Cropland burning).** Emissions from cropland burning were calculated based on the 2000-2005 rice production data derived from BPS. Assumptions from expert judgment mentioned that the proportion (harvest index) for yield and biomass is

1:1, and the fraction of biomass that is burned differs between regions. The emissions of CH<sub>4</sub>, CO, N<sub>2</sub>O and NO<sub>x</sub> from cropland burning are shown in Table 2.10.

**Table 2.10 Distribution of GHG emissions from cropland burning from 2000-2005 (Gg)**

| GHG's emission    | 2000     | 2001     | 2002     | 2003     | 2004     | 2005     |
|-------------------|----------|----------|----------|----------|----------|----------|
| CO <sub>2</sub>   | -        | -        | -        | -        | -        | -        |
| CH <sub>4</sub>   | 47.39    | 43.86    | 45.45    | 45.79    | 47.63    | 47.44    |
| N <sub>2</sub> O  | 1.23     | 1.14     | 1.18     | 1.19     | 1.23     | 1.23     |
| CO                | 1,614.90 | 1,494.66 | 1,548.68 | 1,560.22 | 1,623.03 | 1,616.36 |
| NO <sub>x</sub>   | 43.88    | 40.62    | 42.08    | 42.40    | 44.10    | 43.92    |
| CO <sub>2</sub> e | 1,376.17 | 1,274.46 | 1,319.74 | 1,330.47 | 1,381.53 | 1,377.48 |

#### 2.5.4 Land Use Change and Forestry (LUCF)

Application of the 2006 IPCC GL for LUCF requires consistent land categories to estimate emissions and removal of GHGs for all land categories, relevant carbon pools, and non-CO<sub>2</sub> gases (based on key source/sink category analysis). This has implications for the activity data required and emissions factors for all land categories, carbon pools and related non-CO<sub>2</sub> gases. Since land categories in the 2006 IPCC GL are different from current forest classes applied by Directorate Forestry of Forestry Planning (Ministry of Forestry), the forest categories by Ministry of Forestry were grouped following the 2006 IPCC GL in developing the GHG inventory for this sector, as shown in Table 2.11.

**Table 2.11 Adjustment of Indonesia's land category by Ministry of Forestry into 2006 IPCC GL categories**

| No. | Forest Category (MoF, 2008) | 2006 IPCC Land Category |
|-----|-----------------------------|-------------------------|
| 1.  | Primary Dry land Forest     | Forest Land             |
| 2.  | Primary Swamp Forest        | Forest Land             |
| 3.  | Primary Mangrove Forest     | Forest Land             |
| 4.  | Secondary Dry land Forest   | Forest Land             |
| 5.  | Secondary Swamp Forest      | Forest Land             |
| 6.  | Secondary Mangrove Forest   | Forest Land             |
| 7.  | Plantation Forest           | Forest Land             |
|     | Other land use (APL)        |                         |
| 8.  | Shrubs                      | Grassland               |
| 9.  | Swamp shrubs                | Wetland                 |
| 10. | Open lands                  | Other Land              |
| 11. | Swamp                       | Wetland                 |
| 12. | Agriculture                 | Cropland                |

| No. | Forest Category (MoF, 2008) | 2006 IPCC Land Category |
|-----|-----------------------------|-------------------------|
| 13. | Mix agriculture shrubs      | Cropland                |
| 14. | Transmigration              | Cropland                |
| 15. | Settlement                  | Settlement              |
| 16. | Grassland                   | Grassland               |
| 17. | Rice field                  | Cropland                |
| 18. | Estate crops                | Cropland                |
| 19. | Dyke                        | Other Land              |
| 20. | Airport                     | Other Land              |
| 21. | Water Body                  | Wetland                 |
| 22. | Clouds                      | Undefined               |

At present, the Directorate of Forestry Planning (Ministry of Forestry) is still in the process of producing time series data of land use and forest cover activity data using satellite imagery. Meanwhile, the GHG inventory presented here used Tier 1 data that compiles information from official reports, such as the Forestry Statistical Report (Ministry of Forestry), Indonesia Statistics (BPS), and other references. For emissions removal/factors, most data (i.e., biomass growth and carbon stock for some land uses) is taken from country-specific studies (Tier 2). Emissions from peat fires were not estimated by the National GHG Inventory Team, thus emissions estimates from peat were taken directly from available studies.

In reporting the GHG inventory, the source categories follow the 1996 IPCC GL as suggested in the Guidelines for National Communications of non-Annex I Parties of the UNFCCC, adopted in decision 17/CP.8. Therefore, the results of emissions estimates were mapped back to the 1996 IPCC GL format.

In 2000, total net emissions from three main GHGs (CO<sub>2</sub>, CH<sub>4</sub> and N<sub>2</sub>O) from this sector reached 821,254 Gg (Table 2.12). In the following years, the rate of emissions fluctuated considerably. In 2001, it decreased by approximately 8%; in 2002, it increased sharply by approximately 139%. This fluctuation was primarily due to high inter-annual variation of deforestation rates and peat fires. Within each year, rates of carbon removal were quite stable.

**Table 2.12 GHG emissions and removal from LUCF (in Gg CO<sub>2</sub>e)**

| Gas                            | 2000           | 2001           | 2002             | 2003           | 2004             |
|--------------------------------|----------------|----------------|------------------|----------------|------------------|
| CO <sub>2</sub> -sequestration | -411,593       | -402,027       | -384,427         | -435,037       | -431,128         |
| CO <sub>2</sub>                | 1,232,766      | 1,156,433      | 2,349,902        | 1,026,507      | 1,488,520        |
| CH <sub>4</sub>                | 56             | 140            | 14               | 13             | 22               |
| N <sub>2</sub> O               | 24             | 0              | 6                | 6              | 9                |
| <b>Net emissions</b>           | <b>821,254</b> | <b>754,546</b> | <b>1,965,495</b> | <b>591,489</b> | <b>1,057,423</b> |

By source category, forest and grassland conversion are the main sources of emissions, followed by peat fires and peat oxidation (i.e., emissions from organic soils) (Figure 2.21). However, from 2000-2004, the contribution of forest and grassland conversion to the total emissions from this sector varied between 48% and 65%, peat fire between 15% and 32% and peat oxidation between 10% and 25%. The 2008 World Bank study, however, gave different estimates. It reported that the average emissions from LUCF were approximately 2,398,000 Gg, in which 53% were from peat fire, 20% were from peat drainage (peat oxidation), 22% were from deforestation and 5% were from palm oil and timber plantation establishment.




Figure 2.21 GHG emissions from LUCF sector from 2000-2004 by source category

The contribution of peat fires to the total national emissions could actually be higher since the estimate of emissions from available studies varies considerably (Table 2.13). The lowest estimates were from Indonesian Forest Climate Alliance (IFCA) study (MoF, 2008) while the highest was from Hooijer *et al.* (2006). Hooijer *et al.* (2006) estimated the emissions based on a Borneo hot-spot count and a carbon calculation method used by Page *et al.* (2002), who estimated emissions from peat fires during the 1997 El Niño. This approach may lead to an overestimate as hot-spot counts in peat lands are not fully proportional to CO<sub>2</sub> emissions, which are governed by additional factors such as the depth and area of burning. Therefore, this relationship may not also be generally applicable for the whole of Indonesia, and so the extrapolation of emission estimates over Indonesia based on limited ground checks in Kalimantan may lead to overestimation. Van der Werf *et al.* (2008) attempted to improve emissions estimates from peat fires using several sources of satellite data with biogeochemical and atmospheric modelling to better understand and constrain peat fire emissions in Indonesia. This resulted in far lower estimates. The NGHGI adopted the study of the van der Werf *et al.* (2007). In 2000, the authors' estimated emissions from peat fires were calculated at about 172,000 Gg CO<sub>2</sub>e, while the ten-year average emissions from peat fires (1997-2007) were approximately 466,000 Gg CO<sub>2</sub>e .

<sup>1</sup> The use of average emissions figures from 1997-2007 could reduce the inter-annual variation

**Table 2.13 Emissions from peat fires according to relevant studies (in Gg)**

| Year    | Heil <i>et al</i> (2007) | Levine (1999) | Hooijer <i>et al</i> , 2002 Lowest | Hooijer <i>et al</i> , 2002 Highest | Duncan, <i>et al</i> . (2007) | van der Werf <i>et al</i> . (2007) | IFCA (MoF, 2008) | Average   |
|---------|--------------------------|---------------|------------------------------------|-------------------------------------|-------------------------------|------------------------------------|------------------|-----------|
| 1997    | 4,026,000                | 898,000       | 2,970,000                          | 9,423,000                           | 2,567,000                     | 1,202,000                          | 16,600           | 3,015,000 |
| 1998    | 1,082,000                | 242,000       | 799,000                            | 2,534,000                           | 689,000                       | 271,000                            | 3,700            | 803,000   |
| 1999    | 623,000                  | 139,000       | 458,000                            | 1,459,000                           | 396,000                       | 190,000                            | 2,600            | 467,000   |
| 2000    | 304,000                  | 66,000        | 224,000                            | 711,000                             | 194,000                       | 172,000                            | 2,400            | 239,000   |
| 2001    | 645,000                  | 143,000       | 477,000                            | 1,511,000                           | 411,000                       | 194,000                            | 2,700            | 483,000   |
| 2002    | 2,204,000                | 491,000       | 1,624,000                          | 5,155,000                           | 1,404,000                     | 678,000                            | 9,400            | 1,652,000 |
| 2003    | 1,188,000                | 264,000       | 876,000                            | 2,783,000                           | 759,000                       | 246,000                            | 3,400            | 874,000   |
| 2004    | 1,907,000                | 425,000       | 1,408,000                          | 4,462,000                           | 1,217,000                     | 440,000                            | 6,100            | 1,409,000 |
| 2005    | 1,694,000                | 378,000       | 1,250,000                          | 3,960,000                           | 1,078,000                     | 451,000                            | 6,200            | 1,260,000 |
| 2006    | 3,560,000                | 796,000       | 2,625,000                          | 8,334,000                           | 2,270,000                     | 1,111,000                          | 15,300           | 2,673,000 |
| 2007    | 524,000                  | 117,000       | 385,000                            | 1,225,000                           | 334,000                       | 175,000                            | 2,400            | 395,000   |
| Average | 1,614,000                | 360,000       | 1,191,000                          | 3,778,000                           | 1,029,000                     | 466,000                            | 6,400            | 1,206,000 |

Note: Figures in italic represent predictions using emissions pattern according to Heil *et al.* (2007), as defined in Aldrian (2008). IFCA (2007) only provided estimates of cumulative emissions from 2000-2005 equal to 30 million tons of CO<sub>2</sub>. For annual emissions, the proportion of emissions followed the proportion used in van der Warf *et al.* (2007)


A different selection of emissions estimates from peat fire would significantly change the emission figures of Indonesia. Table 2.14 shows the change in national emissions figures due to changes in the selection of the peat emissions figures and the relative position of Indonesia as an emitter country.

**Table 2.14 Impact of selecting different emissions estimates from peat fire on national figures and the relative position of Indonesia as an emitter country**

| No. | Option   | Emissions from peatfires (Gg CO <sub>2</sub> ) | Total (+1,244,000 Gg CO <sub>2</sub> ) | Global emitter rangk |
|-----|--|--|--|----------------------|
| 1   | Peat fires in 2000– highest (Page <i>et al</i> , 2002)         | 711,000  | 1,955,000                              | 5th                  |
| 2   | Peat fires in 2000 – lowest (IFCA, 2007)                       | 2,400  | 1,246,000                              | 7th                  |
| 3   | Peat fires in 2000 – latest (van der Warf <i>et al.</i> ,2008) | 172,000  | 1,416,000                              | 7th                  |
| 4   | Average –highest (Hooijer <i>et al.</i> , 2002)                | 3,778,000                                      | 5,022,000                              | 2nd                  |
| 5   | Average –lowest (IFCA, 2007)                                   | 6,400  | 1,250,000                              | 7th                  |
| 6   | Average of all studies, all years                              | 1,206,000                                      | 2,450,000                              | 3rd                  |
| 7   | The latest average (van der Warf <i>et al.</i> ,2008)          | 466,000  | 1,710,000                              | 6th                  |

### 2.5.5 Waste

According to the 2006 IPCC GL, sources of emissions from the waste sector are found in four main categories: solid waste, biological treatment of solid waste, incineration and open burning of waste and wastewater treatment and discharge, as shown in Figure 2.22.



**Figure 2.22 Categories of GHG emissions sources within the waste sector**

- Concerning solid waste disposal: In urban areas, almost 60% of waste is taken to solid waste disposal site (SWDS), while in rural areas or small cities, this figure is only 30% (Indonesian Statistical Data on Environment; BPS, 2007). The major components of solid waste brought to SWDS are organic compounds as the other types of waste (plastics, metal, etc.) are generally recycled for re-utilization. The dominant organic compound in solid waste will affect the degradable organic content (DOC) value and the corresponding correction of the CH<sub>4</sub> emissions factor in inventories. The SWDSs in most big cities in Indonesia are considered to be unmanaged SWDS because they are simply open dumping systems; within the context of GHG emissions, they are categorized as unmanaged deep (>5 m) waste.


Currently, incinerators for municipal solid waste are generally not used in Indonesia. Although several statistical data indicate that incineration is already used for eliminating municipal solid waste, in reality, the so-called ‘incinerator’ is actually an ‘open burning’ system. Therefore, the calculation of CO<sub>2</sub> emissions from municipal solid waste is based on open burning.

- Concerning domestic wastewater: In general, the discharge pathways of domestic wastewater in urban areas in Indonesia are decentralized using individual septic tanks. In rural areas, there is almost no wastewater treatment.
- Concerning industrial wastewater: In general, industrial wastewater is treated by the industry prior to discharge to the environment. The calculation of GHG emissions level varies according to the type of industry and the corresponding treatment technology.

From this analysis, it was found that the 2000 total net emissions from the three main GHGs (CO<sub>2</sub>, CH<sub>4</sub> and N<sub>2</sub>O) of the waste sector reached 157,328 Gg (Table 2.15). Methane contributed to approximately 97% of the total emissions. The rate of emission in the period from 2000-2005 was relatively stable. By source category, the major sources are from wastewater treatment and discharge (Figure 2.23).

**Table 2.15 GHG emissions and removal from waste (in Gg CO<sub>2</sub>e)**

| Gas              | 2000    | 2001    | 2002    | 2003    | 2004    | 2005    |
|------------------|---------|---------|---------|---------|---------|---------|
| CO <sub>2</sub>  | 1,662   | 2,266   | 2,302   | 2,338   | 2,366   | 2,377   |
| CH <sub>4</sub>  | 153,164 | 155,853 | 157,516 | 158,670 | 160,361 | 161,346 |
| N <sub>2</sub> O | 2,501   | 2,699   | 2,982   | 3,066   | 3,072   | 3,108   |
| Total            | 157,328 | 160,818 | 162,800 | 164,074 | 165,799 | 166,831 |



**Figure 2.23 GHG emissions from the waste sector from 2000-2005 by source category**

## 2.6 Emissions Trends

Of the five sectors considered here, there are only three sectors where the rates of emissions have increased. These sectors are energy, agriculture and industry, while emissions from the waste sector were relatively constant and those from the LUCF sector fluctuated considerably

between years (Figure 2.24). The net emissions from the LUCF sector were the lowest in 2003. In 2003, the rate of reforestation was relatively high while rate of deforestation was low.




Figure 2.24 Emissions trends by sector

## 2.7 Key Category Analysis

The key category analysis (Tier 1) without and with LUCF indicated that there are 20 and 5 key categories respectively. Without LUCF, the first three main categories that contributed to more than 50% of the total emissions were (i) energy production (electricity, heat, oil & gas refining), (ii) industrial wastewater treatment and discharge and (iii) manufacturing industries and construction (Table 2.16). With LUCF, the first three main categories are all from the LUCF sector. These include (i) forest and grassland conversion, (ii) peat fire and (iii) CO<sub>2</sub> emissions and removals from soils. This analysis suggests that improvement of activity data and emission factors for these key categories were very important to improve the Indonesian National GHG Inventory. It appears that activity data on land use and forest change area and the forested area are the most important. Local emission factors for these source categories need to be developed.

Table 2.16 Key Category Analysis

| Sector              | Source Categories to be Assessed in Key Source Category Analysis | Gases           | Cumulative Contribution (%) |
|---------------------|--|-----------------|-----------------------------|
| <b>Without LUCF</b> |  | 2,266           | 2,302                       |
| Energy              | Energy production (electricity, heat, oil & gas refining)        | CO <sub>2</sub> | 19.8                        |
| Waste               | Industrial Wastewater Treatment and Discharge                    | CH <sub>4</sub> | 38.5                        |
| Energy              | Manufacturing Industries and Construction                        | CH <sub>4</sub> | 50.9                        |
| Energy              | Transportation   | CO <sub>2</sub> | 61.8                        |



| Sector             | Source Categories to be Assessed in Key Source Category Analysis | Gases            | Cumulative Contribution (%) |
|--------------------|--|------------------|-----------------------------|
| Agriculture        | Rice Cultivation   | CO <sub>2</sub>  | 66.9                        |
| Industrial Process | Cement Production  | CH <sub>4</sub>  | 71.4                        |
| Energy             | Residential  | CO <sub>2</sub>  | 75.6                        |
| Agriculture        | Direct N <sub>2</sub> O Soils                                    | CO <sub>2</sub>  | 78.2                        |
| Energy             | Oil and Natural Gas  | N <sub>2</sub> O | 80.6                        |
| Waste              | Unmanaged Waste Disposal Sites                                   | CH <sub>4</sub>  | 82.6                        |
| Energy             | Non Specified  | CH <sub>4</sub>  | 84.6                        |
| Agriculture        | Enteric Fermentation   | CO <sub>2</sub>  | 86.5                        |
| Waste              | Domestic Wastewater Treatment and Discharge                      | CH <sub>4</sub>  | 88.0                        |
| Waste              | Unmanaged Dumpsite   | CH <sub>4</sub>  | 89.5                        |
| Energy             | Residential  | CH <sub>4</sub>  | 91.0                        |
| Energy             | Commercial/Institutional   | CH <sub>4</sub>  | 92.3                        |
| Industrial Process | Ammonia Production   | CH <sub>4</sub>  | 93.5                        |
| Agriculture        | Indirect N <sub>2</sub> O Soils                                  | CO <sub>2</sub>  | 94.4                        |
| Energy             | Oil and Natural Gas  | CO <sub>2</sub>  | 95.3                        |
| <b>With LUCF</b>   |  |                  |                             |
| LUCF               | Forest and grassland conversion                                  | CO <sub>2</sub>  | 39.6                        |
| LUCF               | Peat Fire  | CO <sub>2</sub>  | 65.2                        |
| LUCF               | CO <sub>2</sub> emissions and removals from soils                | CO <sub>2</sub>  | 79.3                        |
| Energy             | Energy production (electricity, heat, oil & gas refining)        | CO <sub>2</sub>  | 86.9                        |
| Waste              | Industrial Wastewater Treatment and Discharge                    | CH <sub>4</sub>  | 94.1                        |
| Energy             | Manufacturing Industries and Construction                        | CO <sub>2</sub>  | 98.9                        |

## 2.8 Uncertainty Analysis

An uncertainty analysis was conducted (Tier 1) following the Good Practice Guidance (GPG) IPCC 2000. The levels of uncertainty for activity data and emission factors were assessed based on expert judgment and consultation with the Centre for Data and Information in the related sectors (Table 2.17). Overall uncertainty of the Indonesian National GHG inventory for 2000 and 2004 without LUCF was approximately 16.3% and 13.9%, respectively, while the trend uncertainty was 16.9%. With the inclusion of LUCF, the level of uncertainty of the National GHG Inventory increased between 15% and 20% (Table 2.18).

Improving activity data and emissions factors for the LUCF sector (the first three key categories) could increase the certainty of the inventory (cf. Table 2.17 and 2.18). This improvement is possible with the use of satellite data. Emissions/removal factors can also be improved with the use of data from National Forest Inventory (NFI). At the time of this inventory, the NFI data is still being processed and not available for use. With the availability of this data, the overall certainty of the Indonesian National GHG Inventory can be improved significantly.

**Table 2.17 Estimates of uncertainty level for activity data and emissions factors**

| No. | Source/Sink Categories                   | Current Uncertainty |       | Improved |       |
|-----|--|---------------------|-------|----------|-------|
|     |  | AD                  | EF/RF | AD       | EF/RF |
| 1   | Energy and transportation                | 10                  | 5     | Same     | Same  |
| 2   | Industry <sup>1</sup>                    | 10                  | 10    | Same     | Same  |
| 3   | Agriculture                              | 15                  | 30    | Same     | Same  |
| 4a  | Change in forest and other woody biomass | 25                  | 50    | 15       | 25    |
| 4b  | Forest and grassland conversion          | 30                  | 75    | 15       | 25    |
| 4c  | Abandonment of managed land              | 25                  | 50    | Same     | Same  |
| 4d  | Soil emissions                           | 50                  | 75    | Same     | Same  |
| 4e  | Peat burning (van der Werf et al. 2008)  | 25                  | 50    | 15       | 25    |
| 5   | Waste                                    | 50                  | 50    | Same     | Same  |

Note: <sup>1</sup>The level of uncertainty for sub-categories of industrial process varied from 5 to 15 while for other sectors were assumed to be the same. AD = activity data; EF/RF = emissions factor/removal factor

**Table 2.18 Level of current uncertainty of Indonesian National GHG Inventory for 2000 and 2004 and trend and potential reduced uncertainty with the incorporation of necessary improvement for the first three key source categories**

| Sectors      | Year  | Uncertainty |          |
|--------------|-------|-------------|----------|
|              |       | Current     | Improved |
| Without LUCF | 2000  | 16.3        | NA       |
|              | 2004  | 13.9        | NA       |
|              | Trend | 16.9        | NA       |
| With LUCF    | 2000  | 47.0        | 23.0     |
|              | 2004  | 38.5        | 20.0     |
|              | Trend | 32.2        | 20.3     |

## 2.9 Quality Assurance/Quality Control (QA/QC)

At present there is no system in place for QA/QC for the GHG emissions. However, the Bureau of Statistics and several other agencies that are responsible for collecting data from local governments and private companies have set up procedures for data quality checking. In the future, sectors that are responsible for developing the National GHG Inventory will develop QA/QC systems. The Ministry of Environment has proposed SIGN (Sistem Inventarisasi Gas Rumah Kaca Nasional) or a National GHG Inventory System to BAPPENAS (National Agency for Planning and Development) to be one of Indonesia's priority climate change programmes. The overall objective of SIGN is to strengthen the capacity of sectors and local governments

in order to improve the quality of the GHG inventory for the development of a sustainable inventory management system. There are three priority focus areas under SIGN:

1. Improvement of methodologies, activity data and emission factors;
2. Strengthening institutional arrangements, their functions, and operations of archiving, updating and managing of greenhouse gas inventories; and
3. Increasing awareness of local governments on the importance of the National GHG Inventory for developing mitigation strategies. Increasing the capacity of designated personnel of the GHG Inventory within each sector for developing and managing the GHG inventory.

## 2.10 Plans for Improvement

Indonesia plans to improve emissions estimates through the improvement of activity data and emissions/removal factors. Priority will be given to LUCF and the agriculture sector.

For the LUCF sector, priority will be given to the improvement of emissions/removal factors from peat and forestland. The initiatives being undertaken include:

- a. Development of emissions factors from peat lands under different uses. Research is being conducted in Central Kalimantan by National Research Consortium for Climate Variability and Climate Change for Agriculture Sector under the coordination of the Ministry of Agriculture. The research is planned to expand to other provinces with support from Global Environment Facility (GEF) using funding allocated for the 3rd National Communication.
- b. Development of emissions/removal factors from forestlands and emissions factors from fire (both in mineral soils and peat land), including the improvement of activity data using satellite, aerial photography and radar by the Ministry of Forestry and partner agencies. The program is operating under INCAS (Indonesian National Carbon Accounting System), a collaboration between Government of Indonesia and Government of Australia.
- c. Implementation of a pilot study to improve GHG emissions estimates from peat lands in the West Kalimantan and Riau Provinces under the coordination of the State Ministry of Environment.
- d. Estimates of CO<sub>2</sub> emissions from peat land using best available activity data (AD) from the sectors and emissions factor/removal factor (EF/RF) from recent studies under the coordination of BAPPENAS in collaboration with Centre for Climate Risk and Opportunity Management-Bogor Agriculture University (CCROM-IPB), Directorate General of Planology – Ministry of Forestry (BAPLAN-MoF), and Center for Agriculture Land Resources and

Development – Ministry of Agriculture (BBSLDA-MoA). This study attempts to use recent findings from the studies conducted by the Ministry of Agriculture under the National Research Consortium for Climate Variability and Climate Change and Indonesia's National Carbon Accounting System (INCAS) project.

For the agriculture sector, the National Research Consortium is undertaking the improvement of emissions and removal factors for climate variability and climate change. The activities include:

- a. Development of a correction factor for GHG emission from livestock. According to Suryahadi *et al.* (2001), feeds influenced the rate of energy converted to methane. Because feed management systems of livestock production in Indonesia varies, the correction of emissions factors must be based on the feed condition, particularly organic matter (OM) and energy consumption level. The emissions factors from the livestock sub-sector can also be corrected by considering animal population structure.
- b. Development of a correction factor for methane emissions from various rice cultivars and emissions factor for fertilizer application (N<sub>2</sub>O emission factors) and emissions from application of different nitrogen resources and organic fertilizer.
- c. Improvement of activity data on the fraction of biomass burning, to complete information on the proportion of crop residue burned, for animal consumption and industry. Data used to calculate emissions from biomass burning are only based on the proportion of rice being harvested in Indonesian rice fields. It was assumed that the proportion is 1:1; however, this proportion may only represent irrigated rice and not rain-fed and upland rice.
- d. Improvement of activity data on biomass residues from non-rice crops, such as dry land and industrial crops (i.e., corn, peanuts, soybean, palm oil, cacao, sugar cane, etc). Those crops have a high carbon/nitrogen ratio as compared to rice and probably have higher CO<sub>2</sub> emissions compared to rice.
- e. Development of sequestration capacity of different agriculture system particularly perennial crops that increase sharply in last decades.
- f. Establishment of national database on carbon balance (GHG emission and sequestration) on agriculture through regular monitoring.

## CHAPTER III

# General Description of Steps Taken or Envisaged to Implement The Convention

### 3.1 Introduction

Following Indonesia's ratification of the United Nations Framework Convention on Climate Change (UNFCCC) in 2004 and the Kyoto Protocol in 1997, the Government of Indonesia (GOI) stated its strong commitment to support global efforts to combat climate change. As stated in its Article 3.4, the Convention urges member parties of the UNFCCC to incorporate climate change into national development planning. Article 4.1 of UNFCCC also underlines the needs of all countries to develop their specific national development priorities and objectives in accordance to principles of common but differentiated responsibilities. Then, in the 13th Conference of Parties (COP13) in 2007, the UNFCCC delivered the Bali Action Plan, addressing long-term cooperative action on climate change and reiterating references to the need for developing countries to integrate adaptation actions into sectoral and national planning, specific projects and programmes. Moreover, the developing countries also need to enhance mitigation actions in the context of the global sustainable development agenda.


In an effort to maintain the spirit of COP13, the GOI made a number of significant efforts to effectively implement the UNFCCC post-conference. These include (i) the development of a number of policy documents outlining the GOI's efforts in integrating climate change mitigation and adaptation activities into the National Medium-Term Development Plan (Rencana Pembangunan Jangka Menengah Nasional, or RPJMN) and (ii) the preparation and the issuance of regulations that assist sectors and local governments in the implementation of climate change programs.

### 3.2 Process of Integrating Climate Change into the National Development Plan

The GOI has started several initiatives to integrate mitigation and adaptation climate change principles into national development planning agenda. The climate issue has been mentioned clearly in the Long-Term National Development Planning 2005-2035 (RPJMP 2005-2035). The plan highlights the various disasters caused by extreme climatic events in Indonesia, including the recent floods and droughts that have brought about heavy losses to the national economy. The National Mid-Term Development Plan (RPJMN 2004-2009) has also set the stage for integrating climate change into national development planning. Climate change is described in Chapter 19 on Agriculture, Fishery and Forestry Revitalization, Chapter 28 on Improvement of People's Access to Quality Health Services, Chapter 32 on Improvement of Natural Resources and Environmental Management, and Chapter 33 on Acceleration of Infrastructure Development. The main focus of the Plan is to integrate climate change into cross-sectoral development


priorities. In the Annual Government Work Plan (Rencana Kerja Pemerintah, RKP) 2008 (RKP 2008), climate change was addressed through the Disaster Risk Reduction Program (which includes meteorological disasters), one of the RKP's eight priority programs. As a consequence, the budget allocation for the Meteorological Early Warning System (MEWS) program in BMKG (Meteorology, Climatology, and Geophysical Agency) has been augmented to improve the national early warning system for climate variability in Indonesia.

In order to fine-tune the current and the subsequent development plan, particularly in formulating the mid-term national plan (RPJMN 2010-2014) and the annual work plan (RKP 2009 and RKP 2010) to be more responsive and address sectoral and cross sectoral issues sensitive to climate change impacts, BAPPENAS, through collaboration with various government institutions, developed "The National Development Planning: Response to Climate Change" documents also known as "The Yellow Book." The Yellow Book shall also serve as reference for the international community to support climate change-related policies, programs and activities prioritized by the GOI. To develop the Yellow Book, BAPPENAS used recent studies related to climate change in Indonesia such as the Technological Needs Assessment, National Communications, Climate Change National Action Plan, etc. Linkages between climate change related documents, development planning and the budgeting process are shown in Figure 3.1.



**Figure 3.1 Linkages between climate change-related documents, development planning and the budgeting process**


To provide guidance for synergizing programs and actions on climate change adaptations and mitigations within sub-sectors, each sector under the coordination of BAPPENAS developed a sectoral roadmap. The roadmap provides sectoral policy direction, strategies and programs to address climate change, and sectoral long-term commitment to emissions reductions and adaptation measures as well as ongoing, innovative and future climate mitigation and adaptation programs. Thus, the roadmap provides detailed policy guidance for climate change planning, programming and actions. Figure 3.2 presents the process of developing the climate change sectoral roadmap.



**Figure 3.2 Development of the climate change roadmap**

Following commitment of the GOI to actively participate in reducing its emissions through National Appropriate Mitigation Action (NAMA), in 2009, the President of the Republic of Indonesia at the G-20 meeting inPittsburgh and at COP15 in Copenhagen committed to an ambitious, world-leading target of 26% reduction in carbon emissions from Business As Usual (BAU) by 2020. Further emissions reductions of 41% are expected with international support. With this commitment, Indonesia will follow a low carbon development path. As such, BAPPENAS is coordinating the sectors to develop the National Action Plan for Reducing Greenhouse Gas Emissions (RAN-GRK). The GOI lists three principles in the RAN-GRK, stating that the mitigation actions (i) should not hinder economic growth, and should prioritize people’s welfare, especially in with regard to energy resilience and food security, (ii) support protection of poor and vulnerable communities, including environment conservation in the framework of sustainable development and (iii) consist of core activities to reduce emissions and supporting

activities to strengthen the policy framework. Figure 3.3 shows the interconnection between policies, guidelines, roadmaps and investment funds towards low carbon development. One new source of funds to support the implementation of climate change programs in Indonesia is the Indonesian Climate Change Trust Fund (ICCTF), described briefly in the following section.



**Figure 3.3 Process of mainstreaming climate change into national agenda: Policies, guidelines, roadmaps & investment funds (BAPPENAS, 2009b). Note: M is mitigation and A is adaptation.**

### 3.3 Financial Institutions

Climate change issues are one priority focus of the national development planning policy. This focus includes the improvement of environmental management and control, capacity building and institutional set-up for the climate change-related stakeholders and the capacity building related to the disaster management. Current government funding for climate change-related programs is insufficient. Indonesia recognizes that climate change can only be properly addressed by allocating specific funding to the issue. In the past, government activities targeting climate change were embedded into regular development programs; with no specific national budget allocated to climate change, funding was instead main streamed into existing programs.


To avoid economic down turn and overcome funding difficulties that may arise when addressing the impacts of climate change, international support plays a key role in enabling Indonesia to effectively develop and implement climate change programs. In this regard, the GOI prioritizes grant utilization to finance climate change programming. Grants to fund climate change activities can originate directly from bilateral or multilateral donors (under Official Development Assistance [ODA] rules and procedures) or via a trust fund (under related trust fund rules and procedures). However, loan resources can be utilized when grant funding is insufficient and the utilization of loans will be the last alternative (BAPPENAS, 2009b). Nevertheless, the GOI recognizes the importance of establishing a management system to link international financial



resources with national investment strategies since this external cooperation and financing needs to be properly prepared for and managed. The GOI has set up the previously mentioned financing management system for supporting and accelerating the implementation of climate change programs called the Indonesian Climate Change Trust Fund (ICCTF).

The ICCTF aims to be a showcase of alternative financing for climate change mitigation and adaptation programs (BAPPENAS, 2009b). At this stage, the ICCTF has five specific objectives:(i) to facilitate and accelerate investment in renewable energy and efficiency and simultaneously reduce GHG emissions from the energy sector, (ii) to reduce emissions from deforestation and forest degradation and stabilizing carbon stock through sustainable forest and peat land management, (iii) to reduce vulnerability in coastal zones, agriculture and water sectors, (iv) to bridge the financial gaps necessary to address climate change mitigation and adaptation and (v) to increase the effectiveness and impact of external finance for climate change programs.

At the initial phase, the ICCTF will be created as an “Innovative Fund,” which involves grant funding from development partners that will help overcome barriers for early program deployment. At the later stages, the ICCTF may advance by establishing a “Transformation Fund” mechanism, which would involve all available funding (i.e., public-private partnerships, loan and world capital market sources). This Transformation Fund also aims to assist with market penetration. As such, at the initial phase, the ICCTF will be dominated by public funding and at the later stages will draw predominantly on private funds (Figure 3.4). The ICCTF can be accessed by sectoral ministries and other stakeholders to support the implementation of climate change programs. The coordination mechanism of the ICCTF is presented in Figure 3.5.



**Figure 3.4 ICCTF development (BAPPENAS, 2009b)**




Figure 3.5 Coordination mechanism of ICCTF (BAPPENAS, 2009b)

## CHAPTER IV

# Measures to Facilitate Adequate Adaptation to Climate Change

## 4.1 Introduction

In the past four decades, climate-related hazards such as floods, droughts, storms, landslides and wildfires have caused major loss of human lives and livelihoods, the destruction of economic and social infrastructure, as well as environmental damage. In many parts of the world, the frequencies and intensities of these hazards have increased (Sivakumar, 2005; Asian Disaster Reduction Center [ADRC], 2005). Floods and windstorms accounted for 70% of all disasters during this period; droughts, landslides, forest fires, heat waves and others account for the remaining 30%. Between 2003-2005 alone, there were approximately 1,429 disaster incidences in Indonesia. About 53.3% of the disasters during that time period were hydro-meteorological disasters (BAPPENAS and Bakornas PB, 2006). Floods occurred most often (34% of the time), followed by landslides (16%). It is likely that global warming will lead to greater extremes of drying and heavy rainfall which will, in turn, lead to higher risk of climate hazards (Trenberth and Houghton, 1996; Intergovernmental Panel on Climate Change [IPCC], 2007). A 2006 report from UN Office for Coordination of Humanitarian Affairs (UN-OCHA) indicates that Indonesia is one of the countries vulnerable to climate-related hazards.


In the future, a changing climate brought about by global warming is expected to create new patterns of risk, and higher risks in general. Sea level rise due to melting glaciers and polar ice as well as thermal expansion will contribute to the increased risk of coastal flooding. Increasing intensity of tropical cyclones observed in recent decades may be tied to increasing sea surface temperatures (SSTs). By impacting the hydrologic cycle, global warming is expected to alter climatic ranges, shift regional climatic averages and climate zones, and lead to a higher frequency and amplitude of weather events. Climate variability and change occurring against a backdrop of increasing global population and globalization of economic processes may lead to increased competition over resources and new vulnerabilities (International Research Institute for Climate and Society [IRI] and Asian Disaster Preparedness Centre [ADPC], 2008). With the increased risk of climate-related hazards and disasters, many countries, particularly least developed and developing countries, may have difficulties achieving the UN Millennium Development Goals (MDGs) related to poverty, hunger and human health.

## 4.2 Historical Climate Change

The rapid increase in the concentration of greenhouse gases (GHGs) in the atmosphere has been determined to be the main factor causing global warming and climate change. IPCC (2007) reported that the global atmospheric concentration of carbon dioxide (CO<sub>2</sub>) has


increased from a pre-industrial value of about 280 parts per million (ppm) to 379 ppm in 2005. The atmospheric concentration of carbon dioxide in 2005 exceeds the natural range from the last 650,000 years (180 to 300 ppm) as determined from ice cores. Since the beginning of continuous direct atmospheric measurements (1960), it is clear that the annual growth of the carbon dioxide concentration in the atmosphere between 1995 and 2005 (1.9 ppm per year) was larger than that occurring between 1960 and 2005 (1.4 ppm per year), although there is a year-to-year variability in growth rates. The rapid increase of this gas has been the primary factor responsible for the global temperature increase (Figure 4.1).

Based on trend analysis of maximum and minimum temperature data from 1980-2002 for 33 stations in Indonesia, a significant increase in maximum and minimum temperatures was observed in most of the stations (Boer *et al.*, 2007). On average, the rate of changes in minimum and maximum temperatures across the 33 stations was 0.047°C and 0.017°C per year, respectively. In ocean waters surrounding Indonesia, in the period 1993-2008, the average rate of SST increase ranged from 0.020°C/year to 0.023°C/year (Sofian, 2010). The associated increase in global temperature caused an increase in sea level rise due to ice melting and thermal expansion of sea water. Based on analysis of altimeter data from January 1993 to December 2008, it was found that the rate of sea level rise ranged from 0.2 cm/year to 1 cm/year, with an average of approximately 0.6 cm/year (Figure 4.2).



**Figure 4.1 (a) Anomaly of mean global temperature of sea-land and (b) 2001-2005 mean surface temperature relative to 1951–1980 measured at meteorological stations and ship and satellite SST measurements (Hansen *et al.*, 2006)**

Global warming may lead to changes in regional climate, such as changes in precipitation (i.e., the amount of heavy rainfall) and in climate extremes, such as number of hot days and number of long dry spells. The effect of global warming will be superimposed on decadal climate variability, such as that caused by the inter-decadal or Pacific Decadal Oscillation (PDO), and on inter-annual fluctuations caused by the El Niño Southern Oscillation (ENSO) and the North Atlantic Oscillation (NAO) (Salinger, 2005). All this may lead to a century of increasing climate variability and change, expected to be unprecedented in the history of human settlement and agrarian activities.



**Figure 4.2** Sea level rise trends based on altimeter data from January 1993 to December 2008 (Sofian, 2010)

Based on trend analysis of historical rainfall data from 384 stations overtime scales between 20 and 50 years, significant decreases and/or increases in rainfall were detected in many parts of the Indonesian region. A significant decreasing trend in December-January rainfall was observed in small parts of Java, Papua, and Sumatra and over a large portion of the Kalimantan islands, where as a significant increasing trend was observed in most of Java and eastern Indonesia, including Bali, Nusa Tenggara Barat (NTB) and Nusa Tenggara Timur (NTT) (Figure 4.4a). For June-August rainfall I, significantly decreasing trends were observed in most of the Indonesian region with exceptions in Pandeglang (West Java), Makasar (South Sulawesi), Monokwari, Sorong (Papua) and Maluku (Figure 4.4b). The distribution of maximum rainfall during the wet and dry seasons also changed in some of the locations (Figure 4.3).

The monsoon onset has also changed in many parts of Indonesia. Based on analysis of data from 92 stations spread over Indonesia, the monsoon onset has been increasingly delayed in some parts of Indonesia, particularly in Java; similarly, the length of the wet season has tended to shorten, particularly in South Sumatra, Java and Kalimantan (Badan Meteorologi Klimatologi dan Geofisika [BMKG], 2005). Other studies conducted in East Java also suggest that the number of extreme dry months in the Brantas Catchment area has increased in the last five decades, particularly in areas near the coast (Aldrian and Djamil, 2006). In these coastal areas, the number of extreme dry months increased to 4 months during the last ten years and in 2002, the figure reached 8 months, a level considered as the longest dry season in five decades. In mountain areas, the number of dry months over the last ten years is 1-2 months, with a maximum number of 4 months.

### 4.3 Future Climate Change

Projecting climate into the future has great uncertainty, as the magnitude of change depends on many factors such as GHG concentration in the atmosphere and land use/cover changes. Changes in these factors depend on global socio-economic development and other human behaviors. The IPCC has projected different emissions scenarios and GHG concentrations in order to estimate the changes in the planet's climate. For Indonesia, two IPCC scenarios, SRESA2 and SRESB1, were selected as they reflect current understanding and knowledge about underlying uncertainties in the emissions. SRESA2 describes a very heterogeneous world. The underlying theme is self-reliance and preservation of local identities. Fertility patterns across regions converge very slowly, which results in a continuously increasing global population. Economic development is primarily regionally oriented and per capita economic growth and technological change are more fragmented and slow. SRESB1 describes a convergent world with the same global population that peaks in mid-century and declines thereafter, rapid change in economic structures toward a service and information economy, with reduction in material intensity and the introduction of clean and resource-efficient technology (IPCC, 2000). With these characteristics, the SRESA2 will lead to higher future GHG emissions while SRESB1 leads to lower future GHG emissions.


The impact of different levels of GHG emissions was assessed based on simulations from 14 General Circulation Models (GCMs). This analysis was used to evaluate the direction of rainfall change (either increasing or decreasing) suggested by the 14 GCMs. The level of confidence to a particular direction of change is considered to increased if most or all of the models are in agreement. From the analysis, a set of maps showing a change in seasonal rainfall under the two emissions scenarios (SRESA2 and SRESB1) are shown in Figures 4.5-4.8. Areas highlighted with dark red (value equal to 1) mean that all 14 GCMs suggest a decrease in future rainfall, while dark blue areas (values equal to 0) indicate that all the GCMs suggest an increase in future rainfall.

Figure 4.5 suggests that under increasing GHG emissions scenarios (i.e., SRESA2), most of the models are in agreement that in 2025 the wet seasonal rainfall (DJF ) in Java, Bali, NTB, NTT and Papua will increase, while in other parts of Indonesia will decrease. By 2050 and 2080, most of the Indonesian region will experience higher rainfall than under the current condition, with exceptions in the northern parts of Sumatra and Kalimantan. Furthermore, dry season rainfall (JJA ) in most parts of Java might decrease by 2025, increase again by 2050, and then decrease by 2080, particularly in West Java and South Sumatra. Under low emission scenarios (i.e., SRESB1), the pattern of change is similar to that of high emission scenarios, but the magnitude of change is slightly lower.

---

<sup>2</sup> DJF = December, January, February

<sup>3</sup> JJA = June, July, August



**Figure 4.3 Seasonal rainfall trends (mm/year) for Indonesia (Boer *et al.*, 2009)**




Figure 4.4a Distribution of maximum monthly rainfall during the wet and dry seasons in Sibolga-North Sumatra, Maulaboh-West Sumatra and Jatiwangi-West Java (Boer et al., 2009)






Figure 4.4b Distribution of maximum monthly rainfall (mm) during the wet and dry seasons in MuaraTewe-Central Kalimantan and Sorong-Papua (Boer et al., 2009)





Figure 4.5 Level of probability for seasonal rainfall to change in the future (Note: rainfall in areas colored dark red will decrease in the future, and rainfall in areas colored dark blue will increase in the future with relatively high level of confidence (Boer *et al.*, 2009)

The impact of global warming on the monsoon onset in Java and Bali using more GCMs and empirical downscaling models was conducted by Naylor *et al.* (2007). It was found that under the SRESA2 scenario, total AMJ rainfall in Java and Bali is expected to increase relative to the current pattern by approximately 10% on average, but to decrease in JAS by about 10-25%. The decline in JAS rainfall could be up to 50% in West/Central Java and 75% in East Java/Bali at the tail end of the distributions. In East Java/Bali, some models projected that total rainfall would drop close to zero during JAS (Figure 4.6). Under SRESB1, the projection of changes is similar to those of SRESA2 until 2050. The shifted balances between the AMJ and JAS rainfall seasons will lead to the shifted onset of the wet and dry seasons. The onset of the rainy season in Java and Bali will be delayed under a changing climate (Figure 4.6). However, the uncertainty is quite high, as illustrated by the wide range of results among GCMs for a given empirical downscaling model (EDM) (Figure 4.6). Nevertheless, it is clear that a 30-day delay in monsoon onset is very likely to occur more frequently in 2050 than it does today.



**Figure 4.6 (a) Summed precipitation for AMJ and JAS for the present climate (dashed line) and for the future predicted climate, using the A2 scenario, and (b) Likelihood of exceeding the 30-day monsoon threshold in 2050. The thick rectangle shows the middle tercile, and the horizontal lines on either side show the lower and upper terciles. The arrows indicate the mean future probability for all GCMs. The vertical lines show the observed probability for 1983–2004 (Naylor *et al.*, 2007). Note: The analysis was done using 15 different GCMs and 3EDMs**

Many studies indicate that extreme climate events are often associated with the El Niño Southern Oscillation (ENSO). ENSO is one of the natural phenomena that has resulted in devastating consequences on rainfall and caused disasters in Indonesia. El Niño is often related to drought and La Niña to floods. Based on 43 drought events that occurred between 1844-1998, only six drought events were not associated with El Niño (Boer and Subbiah, 2005; Asian Development Bank [ADB] and BAPPENAS, 1999; Quinn *et al.*, 1978). Moreover, ENSO is considered as one of the overriding control factors in major forest/land fires and haze occurrence and frequency.

<sup>4</sup> AMJ = April, May, June


<sup>5</sup> JAS = July, August, September

Climate-related hazards in Indonesia are also caused by the location and movement of the tropical cyclones in the south eastern Indian Ocean (from January to April) and the eastern Pacific Ocean (from May to December). Essential for cyclone genesis are SSTs above 26.5°C, latitude beyond 5 degrees and small vertical wind shear. Since tropical cyclones have sustained surface winds of 32ms<sup>-1</sup> or more, their impacts to the Indonesian region are commonly local strong winds and heavy rainfall on the order of hours to days. Strong winds also often occur during the transition from the northeast to southwest monsoon and vice versa. Many studies have also suggested that increasing SSTs will strengthen the tropical cyclones (Emanuel, 2005).

## 4.4 Climate Change Impacts and Adaptation Assessment


### 4.4.1 Agriculture

Historical data has indicated that the change in the onset of the wet season and in its length will influence rice production levels and cropping patterns in Indonesia. Rice is the major staple food in Indonesia. The delay of the wet season onset due to El Niño phenomena causes decreasing cumulative cropping area during wet season (MH). Naylor *et al.* (2007) found that a one month delay in wet season onset due to El Niño phenomena would decrease wet season rice production by approximately 65% for West Java and Central Java and 11% for East Java and Bali. This decreasing cropping area during the wet season will usually be compensated by increasing the cropping area during the dry season (MK) which generally has a high risk of experiencing drought (Figure 4.7), except when El Niño phenomena turns into La Niña conditions (Figure 4.8). Significant decreases in rainfall during the dry seasons in El Niño years also affects the production of other national food crops. However, the effect is not as significant compared to the effects on rice crops.



**Figure 4.7** Change in cumulative cropping area in the wet season (MH) and dry season (MK) during ENSO years (El Niño and La Niña) compared to normal years (Boer, 2009)

Long dry seasons during El Niño years significantly affect not only annual crops, but also perennial crops. Based on field observations, a long dry season generally destroys young plants. During the 1994 El Niño for example, the percentage of young plants (age of less than 2 years) die back due to the long dry season could go up to 30%. Based on observations in a number of locations, the average young plant dieback for tea crops was approximately 22%, between 4% and 9% for rubber, about 4% for cacao, between 1.5% and 11% for cashew nuts, about 4% for coffee and between 5% and 30% for coconut. For mature plantation crops such as coconut and palm oil, the impact of severe drought appears after 4-9 months (Hasan *et al.*, 1998).



**Figure 4.8 Mean area of dry season rice affected by drought from 1989-2006 (above), during El Niño years (middle) and La Niña years (bottom). Source: Boer (2009)**


The dynamic interactions of crop pests and diseases on crops also appear to be related with ENSO. Rice growing area destroyed by brown plant hopper ('wereng coklat') tended to increase significantly during LaNiña years. For example, in West Java, the total area destroyed by this pest during 1998 LaNiña could increase up to 80 times the size of the area lost during normal years. In addition, there is an indication that types of major crop pests and diseases have shifted recently. Historically, pink rice stem borer (*Sesamia inferens*), for example, was only a minor problem in Java (e.g., Indramayu, Magelang, Semarang, Boyolali, Kulonprogo, and Ciamis) compared to yellow rice stem borer (*Scirpophaga incertulas*) and white rice stem borer (*Scirpophaga innotata*). Today, this disease has become dominant (Nastari Bogor and Klinik Tanaman Institut Pertanian Bogor [IPB], 2007). According to Kalshoven (1981), regions with distinct dry seasons are favorable for pink rice stem borer. Bacterial leaf blight (*Xanthomonas oryzae pv. Oryza*) in the last three years has also become one of the dominant rice crop diseases when historically, this disease was not important. Saddler (2000) stated that the optimal temperature for growth of this disease is around 30°C.

Similar phenomena has also been observed in non-rice crops. Prior to 1997, twisting disease caused by *Fusarium oxysporum* was not an important disease for red onion crops, but now this is a very important disease not only in lowland areas, but also in the highlands. In the last two years, this disease has seriously attacked red onion crops in a number of onion production centers such as Brebes (Wiyono, 2007). From laboratory research, when being exposed to high temperatures, this crop becomes less resistant to this disease (Tondok, 2003). The phenomenal example is the appearance of Gemini disease in chili in the last five years in all main chili and potato production centre of Java (Bogor, Cianjur, Brebes, Wonosobo, Magelang, Klaten, Boyolali, Kulonprogo, Blitar, and Tulungagung; Nastari Bogor dan Klinik Tanaman IPB, 2007). This disease caused by virus which is transmitted to the crops by kutu kebul (*Bemisia tabaci*). It is not clear whether these phenomena are related to global warming. However, many researchers found that the main triggering factor for the crop disease to proliferate is temperature increase. For example, a significant increase in *Bemisia tabaci* populations on tomato occurred when the temperature increased from 17 to 30°C (Bonaro *et al.*, 2007). The explosion of this virus under elevated temperature has been predicted by Boland *et al.*, (2004) in Canada.

Changes in rainfall and increases in temperature will both directly and indirectly affect dairy cattle production and reproduction performance. The direct impacts are the effects on production and their reproduction systems, while the indirect impacts are due to impacts on the quality and quantity of grass or forage needed for livestock feed. Feed shortages limit the dry matter and nutrient intake of the animal resulting in low productivity, reduced reproductive performance, increased mortality rates, increased culling rates and changes in the population of the animal in some regions. High temperatures may have negative impacts on dry matter intake, milk yield, growth rate, resistance of the animal to diseases and reproductive performance of dairy cattle. A decrease in dry matter intake in dairy cattle during high ambient temperatures is main factor reducing productivity of dry cattle.

Based on a study conducted in Java, rainfall and temperature (defined in the form of a “comfort index”) affect length of the pregnancy period (PP) and interval of birth of sheep (Rohman and Boer, 2001). Sheep raised in regions with lower annual rainfall and higher temperatures tended to have a longer pregnancy period. The interval between two births (BI) was also found to be longer (Figure 4.9). The comfort indices for PP and BI are expressed by the following equations:

$CIPP = -0.00044 R + 0.54889 T$  and  $CIBI = -0.00145 R + 1.08133 T$ , where R and T are annual rainfall and mean annual temperature, respectively.



**Figure 4.9** The relationship between comfort index and length of pregnancy period (left) and comfort index and interval between two births (right). Source: Rohman and Boer, 2001

The increase in temperatures due to elevated CO<sub>2</sub> levels also affects crop yields. Based on a simulation study conducted in Java (Boer *et al.*, 2009), it was found that the negative effect of increasing temperatures on rice yield will reduce if increases in CO<sub>2</sub> concentration considered. The increase of CO<sub>2</sub> from 370 to 480 ppm (SRESB1) or 520 ppm (SRESA2) can eliminate the negative effects of increased temperatures. In some districts, rice production in 2025 and 2050 would actually be higher from the current production rate. However, in general, total rice production in Java will experience a decrease from increasing temperatures with or without considering the positive effects of CO<sub>2</sub>. When the effect of CO<sub>2</sub> was not considered without any technology intervention, the decrease in rice production in Java by 2025 and 2050 could reach 1.8 and 3.6 million tons, respectively; when the effect of CO<sub>2</sub> was considered, the decrease in rice production is 33,964 tons and 59,584 tons in 2025 and 2050, respectively, for SRESA2 and 39,3048 tons and 88,8230 tons in 2025 and 2050, respectively, for SRESB1 (Table 4.1)

**Table 4.1 Decrease in rice production in Java compared to current production as increasing temperature and CO<sub>2</sub> concentration in Scenario1 (Boer *et al.*, 2009)**

| Province     | Increasing CO <sub>2</sub> not considered |                   | Increasing CO <sub>2</sub> considered |                |                 |                 |
|--------------|---|-------------------|---------------------------------------|----------------|-----------------|-----------------|
|              |   |                   | SRESA 2                               |                | SRESB1          |                 |
|              | 2025                                      | 2050              | 2025                                  | 2050           | 2025            | 2050            |
| West Java    | -620,389                                  | -1,207,728        | -22,311                               | -33,034        | -142,034        | -309,315        |
| Central Java | -609,644                                  | -1,180,292        | -27,132                               | -36,170        | -143,739        | -305,261        |
| East Java    | -597,734                                  | -1,194,802        | 15,479                                | 9,620          | -107,274        | -273,653        |
| <b>Total</b> | <b>-1,827,767</b>                         | <b>-3,582,822</b> | <b>-33,964</b>                        | <b>-59,584</b> | <b>-393,048</b> | <b>-888,230</b> |

Note: The calculations were based on existing planting area (see Table 4.2) and cropping index (see Table 4.3). Values are expressed in tons.

The decrease in crop production is due not only to climatic factors, but also non-climatic factors. By taking into account the impact of non-climatic factors in the analysis, it will provide better understanding of future crop production under changing climate. Boer *et al.* (2009) introduced three non-climatic factors in assessing impact of climate change in Java – paddy field conversion to non-agriculture land area, cropping index, and technology intervention. Based on these factors, the following six scenarios were evaluated:

1. **Scenario 1:** In the future, there is no rice field conversion in Java, and no change in cropping index. Therefore, in this case, the change in the rice production is only affected by climate change and changes in CO<sub>2</sub> concentration.
2. **Scenario 2:** The rate of rice field conversion was assumed 0.77% per year, while cropping index is constant or unchanged.
3. **Scenario 3:** No rice field conversion occurs in the future, however, the cropping index is increased from the current condition. In this study, the cropping index (CI) of rice in West Java increased to 2.30 and 2.10 in Central Java and East Java by 2025. By 2050, the IP increases significantly to 2.50 in West Java, 2.30 in Central Java and 2.20 in East Java. The current cropping index (CI<sub>0</sub>) was estimated using the data of harvested area during the wet season (LPMH) and dry season (LPMK) because the standard data of cultivated area and rice field area was not available. The equation for estimating IP is as follows:

$$CI_0 = \frac{LP_{MH} + LP_{MK}}{LP_{MH}}$$


4. **Scenario 4:** The rate of rice field conversion is equal to Scenario 2 and the cropping index increases to that of Scenario 3.
5. **Scenario 5:** Like scenario 3, but crop productivity is assumed to increase in intervals of 5%, 10%, 15%, 20%, 25% and 30% with the invention and implementation of agronomy technology.
6. **Scenario 6:** Like scenario 4, but crop productivity is assumed to increase in intervals of 5%, 10%, 15%, 20%, 25% and 30% as existence of intervention of agronomy technology.



The projected decrease in rice production due to agricultural land conversion is much greater than the decrease due to increasing temperatures. By using a rice field conversion rate of 0.77% per year and no change in the cropping index used in Scenario 2, rice production will decrease by 2025 and continue to decrease by 2050 compared to current production. Total decreased rice production in Java will reach 6 million tons by 2025 and will be more than 12 million tons by 2050. When the effects of CO<sub>2</sub> are considered in the model, rice production decreases by more than 5 million tons by 2025 and by more than 10 million tons by 2050 (Figure 4.10). The total rice field area in 2025 and 2050 using a rice field conversion rate of 0.77% is presented in Table 4.2.

**Table 4.2 Total paddy field area (ha) in three provinces in Java for 2025 and 2050 using a rice field conversion rate of 0.77% per year (Scenario 2)**

| Province     | Current   | 2025      | 2050      |
|--------------|-----------|-----------|-----------|
| West Java    | 1,158,945 | 998,315   | 775,218   |
| Central Java | 1,053,889 | 907,820   | 704,947   |
| East Java    | 1,238,335 | 1,066,702 | 828,322   |
| Total        | 3,451,169 | 2,972,837 | 2,308,487 |



**Figure 4.10 Changes in rice production in Java by 2025 and 2050 compared to current rice production due to increasing in temperatures and CO<sub>2</sub> concentrations for SRESB1 and SRESA2 using various cultivated area values and planting indices (Boer *et al.*, 2008)**

Moreover, when rice field conversion does not occur in Java, the negative effects of increased temperature on rice production can be reduced by increasing cropping index. By using the assumption that the cropping index can be increased following the scenarios defined in Table


4.3, the rice production rate in most districts in Java by 2025 and 2050 are remain stable or even increase compared to the current production rate, with exceptions in the Tulungagung, Kediri, Purworedjo, Wonosobo, Magelang, Sleman, Klaten and Sukoharjo districts. However, when the effects of CO<sub>2</sub> are considered in the simulation model, the negative effects of increasing temperature on the decrease in rice production in these districts is not projected to occur in 2050.

**Table 4.3 Cropping index scenario for rice in three provinces in Java for 2025 and 2050 (Boer *et al.*, 2008)**

| Province     | Current | 2025 | 2050 |
|--------------|---------|------|------|
| West Java    | 1.70    | 2.10 | 2.50 |
| Central Java | 1.79    | 2.00 | 2.30 |
| East Java    | 1.62    | 2.00 | 2.20 |

As shown in Figure 4.10, the decrease in rice production of Java by 2025 and 2050 is mainly due to land conversion and by climate change. Efforts to increase CI to maximum levels cannot maintain the current production in 2025 and 2050. This study suggests that loss of rice production due to increasing temperatures and land conversion in Java in 2025 and 2050 cannot be compensated by increasing CI only. On the other hand, the effort to increase CI following the scenario presented in Table 4.3 is hardly possible when the source of irrigated water is not from reservoirs.

The water source for irrigated rice is mostly not from reservoir. If CI for irrigated rice in which the water source comes from reservoir must be increased to 2.75 and not from reservoir also must be raised to 2.0 in West Java, and 1.8 in both Central Java and East Java require the development of new reservoir. Otherwise, CI can only increase to 2.2 for West Java, 2.0 for Central Java and optimal (e.g., 1.8) for East Java. If the scenario in Table 4.4 is reached, the number of reservoirs in Java should increase, so that the percentage of irrigated rice from reservoirs becomes 65% in West Java, 50% in Central Java and 40% in East Java (Figure 4.11).



**Figure 4.11 Percentage of irrigated rice fields from reservoir by increasing CI to 2.5 for West Java, 2.3 for Central Java and 2.2 for East Java (Boer *et al.*, 2008)**

Another effort to maintain the current rice production level of Java in 2025 and 2050 is through increasing crop productivity by improvement of agronomy such as fertilizer application or improvement of variety. When the effects of CO<sub>2</sub> are not considered, land conversion still occurred and CI increased (Scenario 5), hence, the current level of rice production can be maintained in 2025 by increasing the productivity level by 5%. By 2050, the level of productivity should be at least 25% higher from the current productivity level. When CI cannot be increased (Scenario 6), the level of productivity should be increased by 22% to maintain the current production level in 2025. By 2050, the level of productivity should be increased by 70% to maintain current levels of production (Figure 4.12).

Furthermore, when the effects of CO<sub>2</sub> are considered, the increase in productivity is not necessary to maintain current production levels in 2025. Meanwhile, increasing productivity from 10% to 15% is still required to maintain production level equal to the current production level in 2050. In Scenario 6, if CI is unchanged and land conversion still occurred, the levels of productivity in 2025 and 2050 would have to increase approximately 10% and 50%, respectively, to maintain current production levels. The level of rice productivity in 2025 and 2050 for Scenarios 5 and 6 are presented in Tables 4.4 and 4.5, respectively.

**Table 4.4 Productivity levels (t/ha) require to maintain current production levels in 2025 and 2050 under Scenario 5 (CI increased and land conversion occurred; Boer *et al.*, 2008)**


| Season     | Province     | Rice Production           |                         |      |                                     |      |                                     |      |
|------------|--------------|---------------------------|-------------------------|------|-------------------------------------|------|-------------------------------------|------|
|            |              | Current Production (t/ha) | CO <sub>2</sub> ignored |      | CO <sub>2</sub> considered (SRESA2) |      | CO <sub>2</sub> considered (SRESB1) |      |
|            |              |                           | 2025                    | 2050 | 2025                                | 2050 | 2025                                | 2050 |
| Wet Season | West Java    | 4.70                      | 4.94                    | 5.97 | 4.70                                | 5.17 | 4.70                                | 5.31 |
|            | Central Java | 5.14                      | 5.40                    | 6.53 | 5.14                                | 5.66 | 5.14                                | 5.81 |
|            | East Java    | 5.15                      | 5.41                    | 6.54 | 5.15                                | 5.66 | 5.15                                | 5.82 |
| Dry Season | West Java    | 4.86                      | 5.11                    | 6.18 | 4.86                                | 5.35 | 4.86                                | 5.50 |
|            | Central Java | 5.11                      | 5.36                    | 6.48 | 5.11                                | 5.63 | 5.11                                | 5.77 |
|            | East Java    | 4.88                      | 5.13                    | 6.20 | 4.88                                | 5.37 | 4.88                                | 5.52 |
| Average    |              | 4.97                      | 5.22                    | 6.32 | 4.97                                | 5.47 | 4.97                                | 5.62 |

**Table 4.5 Productivity levels needed maintain current production in 2025 and 2050 under Scenario 6 (CI constant and land conversion occurred)**

| Season     | Province     | Rice Production           |                         |      |                                     |      |                                     |      |
|------------|--------------|---------------------------|-------------------------|------|-------------------------------------|------|-------------------------------------|------|
|            |              | Current Production (t/ha) | CO <sub>2</sub> ignored |      | CO <sub>2</sub> considered (SRESA2) |      | CO <sub>2</sub> considered (SRESB1) |      |
|            |              |                           | 2025                    | 2050 | 2025                                | 2050 | 2025                                | 2050 |
| Wet Season | West Java    | 4.70                      | 5.74                    | 7.99 | 5.55                                | 6.96 | 5.55                                | 7.24 |
|            | Central Java | 5.14                      | 6.27                    | 8.74 | 6.07                                | 7.61 | 6.07                                | 7.92 |
|            | East Java    | 5.15                      | 6.28                    | 8.75 | 6.07                                | 7.62 | 6.07                                | 7.93 |
| Dry Season | West Java    | 4.86                      | 5.93                    | 8.27 | 5.74                                | 7.20 | 5.74                                | 7.49 |
|            | Central Java | 5.11                      | 6.23                    | 8.68 | 6.02                                | 7.56 | 6.02                                | 7.86 |
|            | East Java    | 4.88                      | 5.96                    | 8.30 | 5.76                                | 7.23 | 5.76                                | 7.52 |
| Average    |              | 4.97                      | 5.22                    | 6.32 | 4.97                                | 5.47 | 4.97                                | 5.62 |

As previously discussed, increasing CI following the scenarios in Table 4.3 is difficult to attain. Therefore, rice field conversion rate in Java should be minimized if not prevented. In addition, the development of sustainable food crop land, as stated in Act No. 41/2009 about Protecting Sustainable Food Agricultural Land, is necessary to speed up actions for protecting land conversion.

If land conversion in Java is unavoidable and still occurred at a rate of 0.77% per year, and CI is constant (Scenario2), the development of irrigated rice fields should be developed outside Java to compensate for the loss in production in Java. The expanding irrigated area needed is approximately 1.3-1.8 million ha in 2025 and 3.0-3.8 million ha in 2050 (Figure 4.12).



**Figure 4.12 Expansion of irrigated area to compensate the loss of rice production in Java due to climate change and land conversion where increasing CI is not feasible. Source: Boer *et al.* (2008)**

A rapid assessment conducted by Parry *et al.* (1992) in a number of locations in Indonesia suggested that sea level rise due to global warming will also reduce local rice supply in the Krawang and Subang districts by approximately 300,000 tons. Similarly, maize output would likely be reduced by 10,000 tons – approximately half of this is attributed to inundation. Sea level rise would also be likely to affect fish and prawn production. The loss is estimated to be over 7,000 tons and 4,000 tons, respectively (valued at over US\$ 0.5 million). In the lower Citarum Basin, sea level rise could result in the inundation of 26,000 ha of ponds and 10,000 ha of crop land. This could result in the loss of 15,000 tons of fish, shrimp and prawns and approximately 940,000 tons of rice. Parry *et al.* stated that the socio-economic implications of this transition in the Subang District alone could be the loss of employment for about 43,000 farm laborers. In addition, more than 81,000 farmers would have to look for other sources of income due to the inundation of their rice fields or prawn and fish farms due to sea level rise.

Climate change is a long term threat that must be anticipated because of its adverse effects to the sustainability of agriculture. The Ministry of Agriculture has formulated the concept and strategic steps to cope with the threats. General strategy of adaptation for agriculture sector is focussing on food crops sub-sector in order to strengthen and sustain national food security. Some of the policy and strategy include:

- a. Reforming the regulations and agricultural institutions in form of Ministerial Decrees
- b. Development of information system, advocation, and communication
- c. Development of agricultural insurance system related to climate change impacts (drought, flood, pest and diseases)

Some of the important adaptation program on most vulnerable sub-sectors (food crop and horticulture) to the adverse affects of climate change among others are:

- a. Improvement of water management, irrigation scheme, soil and fertilization management including organic fertilizer and development of carbon efficient farming
- b. Development of early, drought, salinity and inundation tolerant crop varieties
- c. Development of farming risks insurance against adverse climate
- d. Cropping pattern adjustment by preparation and dissemination of guidance and tools as dynamic cropping calendar and flood and drought naticipation blue print

Others strategic program to cope with climate change among others are development of climate field schools and system of rice intensification in all food crop production centers

#### 4.4.2 Water Resources


To evaluate the impact of climate change on water resources, analysis based on River Area Unit (Satuan Wilayah Sungai, or SWS) was conducted in order to assess the change of water deficit duration under current and future climate. The water balance model is defined as follows:

$$\textit{Supply} = \textit{Demand} + \textit{Surplus}$$

If the demand is higher than the supply, the surplus becomes negative and vice versa. Water supply is a function of monthly rainfall ( $R_{monthly}$ ) and forest cover. The forest cover will determine the fraction of rainfall that cannot be utilized ( $f$ ). The formula for estimating the supply is as follows:

$$\textit{Supply} = R_{monthly}(1-f)$$

The fraction of rainfall that cannot be utilized ( $f$ ) range from 20% to 60% depending on percent forest cover (Figure 4.13). With the increase of forest cover, the fraction of unutilized rainfall will decrease. The  $f$  value was determined using the Tank model tested in a number of watersheds in Java.



**Figure 4.13 Relationship between fraction of unutilized rainfall (f) and forest cover (Herienschah *et al.*, 2009)**

Water demand comes from three sectors: domestic use (urban and rural), industry and agriculture, expressed as follows:


$$\text{Demand} = KA_{\text{Farm}} + KA_{\text{Population}} + KA_{\text{Industry}}$$

Water demand for domestic use ( $KA_{\text{Population}}$ ) was estimated by multiplying population by water consumption per capita (i.e., 120-150 liters/capita/day). For industrial sectors ( $KA_{\text{Industry}}$ ), water demand is estimated based on the size of each industrial area (i.e., 500m<sup>3</sup>/ha/day). Water demand for industry was estimated based on human development index (HDI), based on assumption that the higher the HDI, the higher the industrial area. The relationship between HDI and industrial areas of a number of districts is shown in Figure 4.14. Water demand for agriculture ( $KA_{\text{Farm}}$ ) was calculated by multiplying agriculture land area by crop coefficient ( $k_c$ ) and potential evapotranspiration. Because all lands will also evaporate soil water, the rate of soil evaporation or evapotranspiration from all land use cover types was also estimated using a similar formula used for agriculture land.

The projection of future population follows the projection from the Bureau of Statistics (BPS), while agricultural and industrial areas were generated using the following assumptions:

1. Forest cover decreases at rate of 1% per year. Deforestation occurs in all districts. The deforested area in each district is assumed to be equal to the proportion of the forest in the district to the total area of the district.

2. The amount of rice field area in Java was assumed to decrease at a rate of 50,000 ha per year, while outside Java it increases at a rate of 150,000 ha per year.
3. The HDIs assumed to increase in the future. After 2025, districts with HDI equal to less than 65 will increase to 65, those with HDI equal to 65 will increase to 75 and those with HDI equal to more than 75 will increase to 85.



**Figure 4.14 Relationship between HDI and industrial areas in Indonesia (Herienschah *et al.*, 2009)**

From the analysis, it was found that the number of months with surplus of water in most of districts in northern parts of Java, Bali, east and west Nusa Tenggara, parts of north Sulawesi, Gorontalo and south Sulawesi, Lampung and south Sumatra are almost zero (Figure 4.15). Some of districts in these provinces are exposed to long periods of water deficit. In some districts, the length of deficits could last up to 12 months, particularly areas in the eastern parts of Indonesia such as east Nusa Tenggara and west Nusa Tenggara. In east Nusa Tenggara, planting even just one crop a year without irrigation is not possible during El Niño years. During El Niño years, the length of the rainy season could be less than 3 months.

Under the future projected climate, the number of districts with no month of water surplus will increase significantly (Figure 4.16). Under the present climate, approximately 14% of the 453 districts across Indonesia have no months with surplus water. Under SRESB1, this is expected to increase to 19% by 2025 and to 31% by 2050, while under SRESA2, it will increase to 21% and 31% by 2025 and 2050, respectively. Table 4.6 shows that most of districts in Bali (more than 70%) do not have any surplus months, followed by NTB, Java (particularly northern parts of Java), Sulawesi (particularly Gorontalo and south Sulawesi) and Sumatra (mostly in southern

and eastern parts of Sumatra). With such conditions, increasing cropping index in these islands is not possible, further restricting options for increasing rice production.

The above analysis implies that under changing a climate, more districts will have water scarcity problems (Herienschah *et al.*, 2009). The development of new initiatives to anticipate the scarcity of water due to climate change and increases on water demand are needed, especially in urban areas where populations are increasing and industrial activities are taking place. Inter-basin transfer of water may be one of the potential options to anticipate the scarcity of water in the future. In Indonesia, many individual basins have surplus water resources even in areas that have reach the ultimate stage of development while others face serious shortages, especially during extreme drought years. Improved storage capacity and inter-basin transfer of water from surplus to deficit regions could therefore be an option for achieving more equitable distribution of water resources and optimal utilization of these resources. The inter-basin technology has been practiced in West Nusa Tenggara.





Figure 4.15 Status of water balance under current and future climate (Herienschah *et al.*, 2009)





**Figure 4.16** Number of districts without surplus water by length under current and future climate scenarios (Herienschah *et al.*, 2009). Note: Both the bar graph (top) and table (below) show the same data.

**Table 4.6** Number of districts that have no months without water surplus (Herienschah *et al.*, 2009)

| Province   | Baseline | B1-2025 | B1-2050 | A2-2025 | A2-2050 | Total Number of District in the island |
|------------|----------|---------|---------|---------|---------|--|
| Bali       | 7        | 8       | 8       | 8       | 9       | 9                                      |
| Java       | 22       | 26      | 23      | 26      | 23      | 120                                    |
| Kalimantan | 7        | 11      | 12      | 11      | 15      | 54                                     |
| Maluku     | 1        | 1       | 2       | 1       | 2       | 16                                     |
| NTB        | 3        | 6       | 8       | 6       | 9       | 10                                     |
| NTT        | 1        | 2       | 4       | 2       | 6       | 16                                     |
| Papua      | 0        | 0       | 3       | 0       | 3       | 29                                     |
| Sulawesi   | 10       | 15      | 30      | 22      | 40      | 63                                     |
| Sumatera   | 14       | 15      | 29      | 20      | 32      | 135                                    |

### 4.4.3 Forestry

The impact of climate change on tropical forest occurs as a result of changes in soil water availability due to the combined effects of changes in temperature and rainfall. A study in India suggested that projected depletion of soil moisture due to climate change would likely affect teak productivity by reducing the productivity of deciduous forest by approximately 16% (Achanta and Kanetkar, 1996). In Indonesia, however, one important potential threat of climate change on is the increased risk of forest fire due to decreasing dry season rainfall and shortening of the length of the wet season in some part of the region.

During El Niño years 1991/92, 1994/95 and 1997/98, the dry season was extended and carbon emissions from fires measured in 97 monitoring stations across Southeast Asian countries increased significantly (Schimel and Baker 2002). For Indonesia in particular, the strongest El Niño (1997) caused land and forest fire on approximately 11.698 million ha (BAPPENAS-ADB, 1999). Areas most affected by the fire were Kalimantan and Sumatra while the impacts in other provinces were not as severe (Table 4.7). The economic loss caused by the fire was tremendous. OFDA/CRED (2007) stated that these fires were one of the top 10 natural hazards occurred during 1997 and 2007 and the value of all damages and economic losses directly or indirectly related to the 1997/98 fires might reach 17,000 million USD (Table 4.8).

Ardiansyah and Boer (2009) have classified the Indonesian region based on its level of risk to fire. The fire risk was defined based on the monthly pattern of the number of hotspots. Although the hotspot data are not for fire alone, hot spots are closely related to forest and land fire occurrence and smoke fog (e.g., Hooijer *et al.*, 2006). The result of this classification suggested that Indonesia can be classified into 6 cluster patterns (Figure 4.17). Kalimantan and Sumatra were found to be islands that are already exposed to high risk of fire. Based on hotspot density patterns, two provinces shown to have very high hot spot density are Riau Province in Sumatra and Central Kalimantan (Figure 4.17). Hotspot densities in these two islands increases rapidly when dry season rainfall decrease or length of dry season extends, particularly during El Niño years. The hotspot density increased rapidly as the monthly rainfall during the dry season decreased more than 50 mm below normal or when anomaly rainfall was -50 mm (Figure 4.18)

**Table 4.7 Area affected by forest fire in 1997/1998 (expressed in ha)**


| Vegetation               | Sumatra          | Java           | Kalimantan       | Sulawesi       | Wesst Papua      | Total by vegetation type |
|--------------------------|------------------|----------------|------------------|----------------|------------------|--------------------------|
| Montane forest           | --               | --             | 213,194          | --             | 100,000          | 313,194                  |
| Lowland forest           | 383,000          | 25,000         | 2,690,880        | 200,000        | 300,000          | 3,598,880                |
| Peat and swamp forest    | 624,000          | --             | 1,100,000        | --             | 400,000          | 2,124,000                |
| Dry scrub and grass      | 263,000          | 25,000         | 375,000          | --             | 100,000          | 763,000                  |
| Timber plantation        | 72,000           | --             | 883,988          | --             | --               | 955,988                  |
| Estate crops             | 60,000           | --             | 382,509          | 1000           | 3000             | 446,509                  |
| Agriculture              | 669,000          | 50,000         | 2,481,808        | 199,000        | 97,000           | 3,496,808                |
| <b>Total by province</b> | <b>2,071,000</b> | <b>100,000</b> | <b>8,127,379</b> | <b>400,000</b> | <b>1,000,000</b> | <b>11,698,379</b>        |

**Table 4.8 Total national economic loss due to fires during the 1997 El Niño year (expressed in million USD)**

| Sector         | MoE and UNDP (1998) | WWF & EEPSEA (1998) |
|----------------|---------------------|---------------------|
| Agriculture    | 88.6                | 130.7               |
| Forestry       | 508.2               | 640.6               |
| Health         | 43.8                | 256.7               |
| Transmigration | 0.2                 | 0.0                 |
| Transportation | 13.6                | 4.9                 |
| Tourism        | 4.9                 | 19.6                |
| Fire control   | 3.2                 | 3.3                 |
| <b>Total</b>   | <b>662.4</b>        | <b>1055.6</b>       |

Under both emission scenarios, the seasonal rainfall during SON may decrease over most of the Indonesian region (refer back to Figure 4.10). Most of the 14 GCMs agree that the seasonal rainfall will decrease, particularly in Sumatra and Kalimantan. The decrease in rainfall during this season may result in the delay of the onset of the rainy season, thereby increasing fire risk. Figure 4.19 shows that the risk of fire will increase in the future in certain months and decrease in other months. In general, for Pattern 5 region see (Figure 4.17), the fire risk in October will increase for both scenarios except for SRESB1 in 2050. Similar pattern also found in Pattern 6, the fire risk January will increase for both scenarios except for SRESB1 in 2050.

The Centre for International Forestry Research (CIFOR) also found that future risk of fires in parts of Kalimantan, Sumatra and west Java would be greater than in the past due to an increase in average drought indices and the number of days with ‘extreme’ danger rating (Figure 4.20).



**Figure 4.17 Classification of regions at risk to fire based on hotspot density patterns – Sumatra (left) and Kalimantan (right) (Ardiansyah and Boer, 2009)**






Figure 4.18 The relationship between hotspot density and rainfall anomaly in Kalimantan and Sumatra for (a) Pattern 1, (b) Pattern 3, (c) Pattern 4, (d) Pattern 5 and (e) Pattern 6



**Figure 4.19 Median value of hot spot density for pattern 5 and pattern 6 regions under current and future climate (Ardiansyah and Boer, 2009)**



**Figure 4.20 Keetch and Byram Drought Index for baseline period of 1960-1990 (top), for A2 projection period of 2070-2100 (middle) and for B2 projection period of 2070-2100 (bottom) forecast Kalimantan, Indonesia. The fire danger index rating classes are: low (0-999); moderate (1000-1449); high (1500-1749); extreme (1750 - 2000). Source: CIFOR (2008)**

Based on the above finding, the risk of repeated fires may increase under a changing climate. Under such conditions, the threat to biodiversity loss will increase. A study in East Kalimantan showed that the number of trees that survived in plots that experienced fire in 1982/83 and again in 1997/98 decrease significantly (Figure 4.21; Ngakan, 2002). Furthermore, even eleven years after fires, the number of species in plots that were lightly, moderately and severely damaged was not significantly different. However, the species composition was still noticeably different. The trees in lightly-damaged plots were dominated by primary (climax) species (Figure 4.22) such as *Aglaiia cinere*, *Madhuca malacensis*, *Mallotus mpritsianus*, *Eusideroxylon zwageri*, etc.




Figure 4.21 The number of surviving and dead trees per hectare after forest fires in 1982/83 and 1997/98 (Ngakan, 2002)




Figure 4.22 Composition of species per hectare in plots which were lightly, moderately and severely damaged by fires in Kutai National Park, East Kalimantan

Other impacts of climate change on forest ecosystems may also include the disappearance or reduction of density of certain species due shifts in ideal temperature and rainfall range, increasing forest disease and pest infestation, the increase or the decrease of forest productivity, etc.

As defined in Chapter 5, the Government of Indonesia has issued a number of policies and regulations to reduce risk of land/forest fire. There are at least 7 legal instruments designed to control fire. These legal instruments are in the form of Acts, Government Regulations, Minister Decrees from the Ministry of Forestry, Ministry of Environment and Ministry of Agriculture, Director General Decrees, etc. A number of programs have been implemented for fire management through the establishment of community-based fire management systems, early warning systems, etc.

#### **4.4.4 Coastal and Marine**

The impacts of climate change on coastal areas can be divided into two categories: biophysical and socio-economic impacts. Biophysical impacts include (IPCC, 2005): (i) increased coastal erosion, (ii) inhibition of primary production processes, (iii) increased flood frequency, (iv) increased intensity of flood events caused by storm-surge, (v) saltwater intrusion into estuaries and aquifers, (vi) changes in the quality of surface water and groundwater characteristics, (vii) changes in the distribution of pathogenic micro organisms and (viii) increasing sea surface temperatures. Economic and social impacts include (i) increasing number of homes lost and coastal habitats, (ii) increased risk of flooding and the number of lives potentially lost, (iii) damage to buildings and other infrastructure, and coastal protection, (iv) increased risk of disease outbreaks, (v) loss of resources that can be replaced (vi) loss of tourism, recreation and transportation, (vii) loss of cultural assets and historical values and (viii) impacts on inland fisheries and agriculture due to declining quality of soil and water as a result of saltwater intrusion.

In Indonesia, the potential impact of climate change on coastal areas and fisheries is significant. From historical data, it is clear that the increase in sea temperatures during El Niño years has caused serious coral bleaching. Coral bleaching occurred in Indonesian waters in 1983, including the Sunda Strait, and the Thousand Islands, and Karimun Java Sea. This resulted in high numbers of dead coral. In 1998, widespread coral bleaching occurred again, but indifferent locations, this time in East Sumatra, Java, Bali and Lombok. In the Thousand Islands, 90-95% of coral at depths up to 25 m suffered bleaching and death (World Resources Institute [WRI], 2002). Meanwhile, Bali experienced 75-100% coral bleaching. According to the World Wildlife Fund (WWF), the 1998 coral bleaching event was triggered by El Niño. The 1982/83 and 1997/1998 El Niño events have been defined as the two strongest El Niño events during the last century. The Indonesian Coral Reef Foundation (TERANGI) (2007) also reported coral bleaching in the Thousand Islands in February 2006 and February 2007, even though the El Niño during this period was not as strong as those that occurred in 1982/83 and 1997/98.

Coral bleaching has significant impacts on fish populations as it is the habitat of thousands of species of reef fish. In addition, the reef also functions as a high energy and wave absorber that reduces the risk of coastal erosion. Therefore, damage to reefs increases the risk of coastal erosion. The increasing frequency of storms resulting in high waves can also cause clusters of coral damage. Due to its world renowned biodiversity and aesthetic beauty, beaches in Indonesia have become a popular tourist destination and is a significant source of revenue for Indonesia. In general, the number of tourists who visit Indonesia tends to increase and provide major revenue for the government (Ministry of Culture and Tourism, 2010). However, if not managed appropriately, tourism activities can also cause damage to coastal areas, including coral reefs, and beaches.

Rising ocean temperatures will also inhibit the growth of phytoplankton, which many important species of fish depend on. Fish area resource that can move from one place to another, across borders between countries. For example, the migration of Pacific salmon from USA to Canada is suspected to be attributed to a rise in Pacific Ocean temperatures. This situation pushed the two countries to establish a single agreement on salmon fishing (Miller, 2000). The increase in ocean temperatures will also change the pattern of currents, wind and rain patterns. Warmer ocean water can prevent the proliferation of plankton thereby reducing the availability of fish food. Some fish species such as pelagic will likely migrate to other regions that offer suitable temperature conditions and food availability. This will adversely affect the attainment of development objectives in the fisheries sector.

Coastal erosion has also been observed in many parts of Indonesian coastal areas due primarily to sea level rise (SLR) and strong wave action. Based on data released by the Ministry of Public Works, about 40% of the 81,000 km coastline has been eroded. Along the northern coast of Java, the coastal erosion impacts on 5,500 hectares of land are spread over 10 districts. For example, in Tegal Regency, erosion of 40 km of shoreline has resulted in moving the coast line inland as far as 50 to 100 meters. Approximately 3 km out of the 10.5 km of shoreline erosion has also resulted in a loss of 25 ha of ponds.

The increase in wind speed, sea level, and wave and tidal action due to climate change will have significant impacts on coastal and marine resources. As an island country, Indonesia will suffer significantly from climate change-induced sea level rise. The following sections describe briefly the impact of sea level rise on four coastal cities of Indonesia, Jakarta, Medan, Semarang and Surabaya using three scenarios. The first scenario assesses the impact of sea level rise that may cause the permanent inundation of some of the coastal areas. The second scenario assesses the impact of sea level rise combined with high tides. The third scenario assesses the impact of sea level rise combined with land subsidence. The economic losses due sea level rise were assessed based on the area and type of land use affected by the sea level rise. The four main land use types affected by sea level rise in the coastal cities include settlement, rice fields, ponds and harbors/airports. The value of economic losses due to the inundation of the respective land use types were collected based on literature studies (Table 4.9).




**Table 4.9 Economic loss per ha of affected land use type**

| Land Use/Land Cover | Damages in rupiahs (IDR) |                |           | Source |
|---------------------|--------------------------|----------------|-----------|--------|
|                     | Low                      | Medium         | High      |        |
| Harbors/airports    | -                        | 37,500,000,000 | -         | 1      |
| Ponds (prawn/fish)  | 2,150,000                | 2,200,000      | 2,250,000 | 3      |
| Settlements         | 1,640,000                | 2,695,000      | 3,750,000 | 2      |
| Wet rice fields     | 1,750,000                | 3,000,000      | 5,500,000 | 4      |


Source: 1Batam Pos, 2BPSDA (2003), 3 Kartikasari and Boer et al (2007) and 4 Sasmita (2006)

## MEDAN CITY

This case study covers the area of Medan and part of the Deli Serdang Regency. Medan topography is undulating with an altitude between 2.5 and 37.5 meters above sea level (a.s.l.); Deli Serdang area topography is similar to the Medan municipality. Due to the topography, this region is not vulnerable to flooding due to sea level rise equal to one meter or less. The probability of coastal areas affected by flooding due to sea level rise alone (Scenario 1) is low even with sea level rise up to 1 m. However, when combined with high tide (2.69 m), a significant area will be inundated. The time scale of this flood is not permanent and only temporary. Based on this scenario, the most extensive areas affected are Harapan Perak in Deli Serdang District and Medan Labuhan in Medan Municipality (Figure 4.23). The extent of water flooding inland due to sea level rise and high tide is varied depending on the topographic of the area. The maximum height of flooding and the total area temporarily inundated is shown in Figure 4.24. Because of the nature of the floods in this region are only temporary, the damage to infrastructure is small, but if the flood affected the functions of wetlands and ponds, damage will be high and lead to large economic losses.



**Figure 4.23 Distribution of floods based on the scenario of sea level rise + high tide: 0.25+2.69 m (A) and 1.0+2.69 m (B). Source: Hariati *et al.*, 2009**



**Figure 4.24** Distribution breadth and depth of inundation due to sea level rise and high tide of Medan and surrounding areas (Hariati *et al.*, 2009)

The main land use types situated below an elevation of 100 m a.s.l. are ponds, dry land farming and settlements. Distribution of settlement is more concentrated in Medan than in Deli Serdang. The size of settlements likely to be affected by flooding are 320-770 ha, and this will threaten the residents who live in the area. With sea level rise of 2.94 meters (0.25+2.69 m), the total population affected by the temporary flood is approximately 370,000 people; when sea level rise increases by 3.69 meters, the population affected increases to 1 million people. The damage on settlements may not be significant, so the loss is assumed to be zero. However, the impact on rice fields and ponds will be high as it will damage the crop due to saltwater intrusion and flush out the fish in the ponds. With sea level rise of 2.94 meters, the total area of wetland ponds affected by flooding is almost 18,000ha, equivalent to an economic loss of approximately 47 billion IDR. With sea level rise of 3.69 meters, the amount of economic loss increases to approximately 69 billion IDR (Table 4.10).

**Table 4.10** Impact of sea level rise and high tide on population and land use in Medan (Hariati *et al.*, 2009)

| Scenario                | Sea Level Rise (m) | Population (x 1000) | Economic Loss (billion IDR) |       |       |                |       |
|-------------------------|--------------------|---------------------|-----------------------------|-------|-------|----------------|-------|
|                         |                    |                     | Settlement                  | Rice  | Ponds | Harbor/Airport | Total |
| SLR + Tides + Waves (m) | 2.94               | 368                 | 0.00                        | 29.24 | 17.50 | 0.00           | 46.74 |
|                         | 3.19               | 588                 | 0.00                        | 40.41 | 21.06 | 0.00           | 61.48 |
|                         | 3.44               | 761                 | 0.00                        | 43.23 | 21.95 | 0.00           | 65.19 |
|                         | 3.69               | 998                 | 0.00                        | 46.02 | 22.73 | 0.00           | 68.75 |


## JAKARTA CITY

Some areas of Jakarta are located at low elevation or even below sea level (Satkorlak PBP Jakarta 2006). Based on SRTM data analysis, the low elevation areas are distributed across four municipalities (Table 4.11). The size of the areas with an elevation below sea level rise ranges from 90 to 2020 ha. The largest area is situated in North Jakarta and the smallest area is in Central Jakarta. In North Jakarta, low elevation areas are distributed among the six sub-districts of Cilincing, Kelapa Gading, Koja, Pademangan, Penjaringan and Tanjung Priok, while the specific villages with the lowest elevation are Sunter Agung, Papango, and Tanjung Priok (Figure 4.25).

**Table 4.11 Percent distribution area of low elevation ( $\leq 0$  meters below sea level in DKI Jakarta**


| No.   | Municipality    | Area   |       |
|-------|-----------------|--------|-------|
|       |                 | (ha)   | (%)   |
| 1.    | West Jakarta    | 244.5  | 9.9   |
| 2.    | Central Jakarta | 91.4   | 3.7   |
| 3.    | East Jakarta    | 105.1  | 4.3   |
| 4.    | North Jakarta   | 2020.7 | 82.1  |
| Total |                 | 2461.7 | 100.0 |

Source: Analysis of SRTM data




**Figure 4.25 Distribution of low elevation ( $\leq 0$  meterdpal) areas in DKI Jakarta**

Under Scenario 1, some of the Jakarta area will be permanently inundated due to a sea level rise of 0.25 m. The most affected areas are located in the Sunter Agung and Papango of Tanjung Priok sub-districts. The flood depth may reach 11 cm and in some locations can increase up to 20 cm. The coastline will move inland by approximately 1.5 km from its current location. If sea level rises up to 1 m, the area that will be permanently inundated will increase significantly and more sub-districts will be affected, including Tanjung Priok and Koja (North Jakarta) and Penjaringan and Kalideres (West Jakarta) with a mean flood depth of approximately 30 cm. The coastline will move inland by approximately 2.0 km from its current location (Figure 4.26). Combined with high tide (Scenario 2), the total area that will be inundated will be much larger. The depth of flood and total flooded area under Scenarios 1 and 2 are shown in Figure 4.27.




**Figure 4.26** Distribution of floods based on sea level rise of 0.25 m (A), 1.0 m (B), sea level rise + high tide of 2.28 m (C) and sea level rise + high tide of 3.03 m (D). Source: Hariati *et al.*, 2009

Based on the overlay of population density maps and flood area, the size of the population affected by a sea level rise of 0.25 and 1.0 m will be approximately 74,000 and 130,000, respectively (Figure 4.28). If the level of sea level rise is combined with high tide, the total number of people affected will increase to 236,000 and 381,000 people, for 2.28 m and 3.03 m, respectively (Figure 4.28).



**Figure 4.27** Area of flooding and depth of flood due to sea level rise and high tide in the Jakarta region. Source: Hariati *et al.*, 2009



**Figure 4.28** Number of residents affected by flooding due to sea level rise and high tide in Jakarta. Source: Hariati *et al.*, 2009

The impact of sea level rise on the city of Jakarta was assessed based on four land use types: residential, wetland, ponds and ports. With sea level rise of less than 0.25 m, the economic loss will reach approximately 6,500 billion IDR; if sea level rise increased to 1.0 m, the loss will increase to more than 8,600 billion IDR (Scenario 1). If the impact of sea level rise is combined with high tide (Scenario 2), the loss will not be much different from that of Scenario 1 (Table 4.12). This is due to the nature of floods in the second scenario, which are only temporary. It is assumed that the damage caused by the temporary flood in settlements and airport is not

significant. The impact of temporary flooding is only significant on rice field and ponds. In this analysis, the economic impact of temporary floods through their effect on flight landing and take-off at airports is not taken into account.

**Table 4.12 Impact of sea level rise and high tide on the population and land use in Jakarta.**


Source: Hariati *et al.*, 2009

| Scenario                        |      | Number of people affected (in thousands) | Economic Loss (billion IDR) |       |       |                |         |
|---------------------------------|------|--|-----------------------------|-------|-------|----------------|---------|
|                                 |      |  | Settlements                 | Rice  | Ponds | Harbor/Airport | Total   |
| SLR (m)                         | 0.25 | 74                                       | 1.00                        | 0.06  | 0.12  | 6,489.56       | 6490.74 |
|                                 | 0.50 | 93                                       | 1.27                        | 0.07  | 0.23  | 7,202.44       | 7204.01 |
|                                 | 0.75 | 109                                      | 1.63                        | 0.08  | 0.62  | 7,881.83       | 7884.16 |
|                                 | 1.00 | 130                                      | 2.15                        | 0.28  | 0.71  | 8,622.76       | 8625.90 |
| SLR + tides and wave action (m) | 2.28 | 236                                      | 1.00                        | 8.49  | 1.48  | 6,489.56       | 6500.53 |
|                                 | 2.53 | 280                                      | 1.27                        | 9.50  | 1.55  | 7,202.44       | 7214.76 |
|                                 | 2.78 | 325                                      | 1.63                        | 10.15 | 1.63  | 7,881.83       | 7895.24 |
|                                 | 3.03 | 381                                      | 2.15                        | 11.05 | 1.71  | 8,622.76       | 8637.67 |

It should be noted that land subsidence problems already exist in Jakarta. However, due to unavailability of spatial data on land subsidence, analysis for Scenario 3 was not conducted. Based on a number of reports (Abidin *et al.*, 2009; Djaya *et al.*, 2004), subsidence in several places in the metropolitan area of Jakarta ranging from 20 to 200 cm occurred over the period from 1982-1997. In general, land subsidence exhibits spatial and temporal variations, with the rates of about 1-15 cm/year, but in a few locations, the subsidence rates reached 20-25 cm/year (Abidin *et al.*, 2009). The four main factors causing land subsidence in Jakarta are ground water extraction, load of constructions (i.e., settlement on high compressibility soil), natural consolidation of alluvial soil and tectonic subsidence (Abidin *et al.*, 2009). Overall, land subsidence may have more serious impacts than that of the global warming on land inundation on the City of Jakarta.

## SEMARANG CITY


The impact of sea level rise on Semarang City will be severe, as the city already experiences problems with land subsidence (Georisk Project, 2008). In Semarang, the primary factor that triggers land subsidence is the increased consumption of ground water. As a result, hydraulic pressure in the soil layer declines (referred to as cone depression of groundwater), followed by the decrease of the soil surface. This condition will be more severe when the area above ground incurs additional burden, such as settlement building. Based on the Georisk Project report (2008), the cone depression in the area of Semarang widened east wards from 1984 to 2000. Based on these observations, it is assumed that the direction of growing land subsidence is toward the east (Figure 4.29A). The rate of land subsidence ranged from 0-1 cm/year up to 8-9 cm/year (Figure 4.29B). The highest rate of land subsidence (8-9 cm/year) occurred in most of North Semarang and Genuk; the lowest rate of land subsidence (0-1 cm/tahun) occurred in Tugu, Central of Western Semarang, and southern part of Central Semarang, East Semarang, Gayamsari and Padurungan. This data was used as the basis for projecting land subsidence up to 2100.



**Figure 4.29 (A) Growth of cone impairment (cone depression) in the groundwater aquifer from 1984 until 2000, and (B) land subsidence rates in Semarang City. Source: Georisk Project, 2008**

Under Scenario 1, it was found that total area in Semarang City that will be permanently inundated by sea level rise of 0.25 m is less than 600 ha with a flood depth of approximately less than 20 cm. The sub-district with the greatest area affected by flood is Genuk, while the sub-district with the smallest area affected is Tugu. With sea level rise up to 0.50 and 1.0 m, the coastline is projected to move inland between approximately 700-1000 m, and 1000-1300 m, respectively. In the three most affected sub-districts of Semarang (Tugu, Semarang Utara and Genuk), the coastline will move inland by approximately 680, 860 and 1000 m respectively, with a sea level rise of 0.5 m, (Figure 4.30A); with sea level rise of 1.0 m, the coastline will move inland by 1300, 1000 and 300 m, respectively (Figure 4.30B). Under Scenario 2, combining sea level


rise with a tidal height of 2.03 m, more land will be inundated, but it will be only temporarily (Figures 4.30C and 4.30D). The total area flooded and the depth of the flood under Scenarios 1 and 2 are presented in Figure 4.31.




**Figure 4.30** Distribution of floods based a sea level rise of 0.25 m (A), 1.0 m (B), and sea level rise + high tide equal to 2.28 m (C) and 3.03 m (D) in Semarang. Source: Hariati *et al.*, 2009

Under Scenario 3 (sea level rise combined with land subsidence), the areas permanently inundated are similar to those of Scenario 2 (Figure 4.32). With sea level rise of 1.0 m combined with land subsidence, the total area inundated may reach approximately 4,000 ha with a flood depth of about 1.6 m (Figure 4.33). Based on this condition, the plan to re-locate people who live in the coastal areas of Semarang, particularly in Tugu, Semarang Utara and Genuk sub-districts should be designed as soon as possible. At present, in order to adapt to sea level rise and land subsidence, the community has to raise their floors every 5 years by about 50 cm to avoid flooding in their houses during high tide. Table 4.13 shows the projected shoreline retreat in the three sub-districts due to sea level rise and land subsidence.






**Figure 4.31** Flood area and flood depth due to sea level rise and high tide in Semarang.  
 Source: Hariati *et al.*, 2009



**Figure 4.32** Distribution of floods based on land subsidence and sea level rise of 0.25 m (A), 0.50 m (B), 0.75 m (C) and 1.0 m (D). Source: Hariarti *et al.*, 2009




**Figure 4.33 Flood area and flood depth due to sea level rise and land subsidence in Semarang area.**  
Source: Hariarti *et al.*, 2009

**Table 4.13 Shoreline retreat due to sea level rise and land subsidence (Hariarti *et al.*, 2009)**

| No. | Sub-District   | Coast line movement in land due to sea level rise (m) |       | Coast line movement in land due land subsidence (m) |                         |
|-----|----------------|---|-------|---|-------------------------|
|     |                | 0.5 m   | 1.0 m | 0.5 m + land subsidence                             | 1.0 m + land subsidence |
| 1   | Tugu           | 700   | 1300  | 930   | 1800                    |
| 2   | North Semarang | 800   | 1030  | 2300  | 2800                    |
| 3   | Genuk          | 950   | 1200  | 2970  | 4300                    |

Sea level rise of 0.25 and 1.0 will threaten approximately 31,000 and 114,000 people, respectively. If sea level rise occurs simultaneously with high tide, the number of people threatened by floods increases. With a total increase in sea level of 2.28 and 3.03 m as a result of high tide and sea level rise, the number of people threatened increases to approximately between 177,000 and 443,000, respectively (Figure 4.34).

The economic loss due to sea level rise ranges considerably depending on whether the area is inundated temporarily or permanently (Table 4.14). Under Scenario 1, sea level rise of 0.25 m will result in an economic loss of approximately 1.5 billion USD, while if the sea level rises 1 m and is combined with land subsidence, the total economic loss may reach 378 billion USD (Table 4.14).



**Figure 4.34** The number of residents affected by flooding due to sea level rise and high tide.  
Source: Hariarti *et al.*, 2009


**Table 4.14** Impact of sea level rise, high tide and land subsidence on the population and land use in Semarang. Source: Hariarti *et al.*, 2009

| Scenario                  | Number of people affected (x 1000) | Economic Loss (billion IDR) |      |       |                |        |        |
|---------------------------|------------------------------------|-----------------------------|------|-------|----------------|--------|--------|
|                           |                                    | Settlement                  | Rice | Ponds | Harbor/Airport | Total  |        |
| SLR (m)                   | 0.25                               | 31                          | 1.12 | 0.00  | 0.37           | 0.00   | 1.49   |
|                           | 0.50                               | 51                          | 1.49 | 0.00  | 0.65           | 0.00   | 2.14   |
|                           | 0.75                               | 77                          | 1.76 | 0.00  | 0.94           | 0.00   | 2.69   |
|                           | 1.00                               | 114                         | 2.08 | 0.00  | 1.24           | 8.76   | 12.08  |
| SLR + high tide (m)       | 2.28                               | 177                         | 1.12 | 0.00  | 2.12           | 0.00   | 3.25   |
|                           | 2.53                               | 244                         | 1.49 | 0.01  | 2.13           | 0.00   | 3.63   |
|                           | 2.78                               | 330                         | 1.76 | 0.01  | 2.14           | 0.00   | 3.91   |
|                           | 3.03                               | 443                         | 2.08 | 0.01  | 2.15           | 8.76   | 13.00  |
| SLR + land subsidence (m) | 0.25                               | 69                          | 2.54 | 0.00  | 0.83           | 0.00   | 3.37   |
|                           | 0.50                               | 235                         | 6.15 | 0.00  | 1.30           | 0.00   | 7.45   |
|                           | 0.75                               | 360                         | 7.61 | 0.01  | 1.67           | 59.91  | 69.19  |
|                           | 1.00                               | 495                         | 8.77 | 0.01  | 1.92           | 367.53 | 378.22 |

## SURABAYA CITY

Based on sea level rise of 0.25 m to 1.00 m (Scenario 1), the area of flooding in Surabaya will be less than 2,000 ha, but if it is combined with high tide, a 2.91 m rise (Scenario 2), the area of flooding will reach 10,000 ha (Figures 4.35 and 4.36). The sub-districts which will suffer the most from flooding will be Sukolilo, Gunung Anyar and Rungkut, in that order. Visual observations using satellite imagery and Google Earth found that the areas most under threat from sea level rise are settlements. In addition to settlements, the other land uses potentially affected by flooding are paddy fields, ponds and other infrastructure. The number of people threatened by flooding under Scenarios 1 and 2 will be less than 150,000 and more than 700,000 people, respectively (Figure 4.37).

The expected economic loss due to flooding is presented Table 4.15. For Scenarios 1 and 2, the total economic loss due to sea level rise of 0.25 m will be about 133 billion IDR and 145 billion IDR, respectively. There is only a slight increase from Scenario 1 to Scenario 2. As described previously, this is due to the differences in the nature of flood inundation in Scenario 1 and 2 whereas the floods are permanent in Scenario 1 and only temporary in Scenario 2.



**Figure 4.35** Distribution of floods based sea level rise of 0.25 m (A), 1.0 m (B), and sea level rise + high tide equal to 3.16 m (C) and 3.91 m (D) in Surabaya. Source: Hariati *et al.*, 2009




Figure 4.36 Flood area and flood depth due to sea level rise and high tide in Surabaya.  
Source: Hariati et al., 2009




Figure 4.37 The number of residents affected by flooding due to sea level rise and high tide in Surabaya.  
Source: Hariati et al., 2009

Table 4.15 Impact of sea level rise and high tide on population and land uses in Surabaya.  
Source: Hariati et al., 2009

| Scenario                    | Number of people affected (x 1000) | Economic Loss (billion IDR) |      |       |                |        |        |
|-----------------------------|------------------------------------|-----------------------------|------|-------|----------------|--------|--------|
|                             |                                    | Settlement                  | Rice | Ponds | Harbor/Airport | Total  |        |
| SLR                         | 0.25                               | 61                          | 0.34 | 0.00  | 4.01           | 128.52 | 132.87 |
|                             | 0.50                               | 73                          | 0.40 | 0.00  | 4.27           | 141.36 | 146.03 |
|                             | 0.75                               | 96                          | 0.56 | 0.19  | 4.53           | 191.06 | 196.34 |
|                             | 1.00                               | 122                         | 0.68 | 0.22  | 4.95           | 316.61 | 322.46 |
| SLR + tides and wave action | 3.16                               | 310                         | 0.34 | 4.22  | 11.65          | 128.52 | 144.73 |
|                             | 3.41                               | 401                         | 0.40 | 4.58  | 12.51          | 141.36 | 158.84 |
|                             | 3.66                               | 546                         | 0.56 | 4.83  | 13.03          | 191.06 | 209.49 |
|                             | 3.91                               | 751                         | 0.68 | 5.18  | 13.56          | 316.61 | 336.03 |

The increase in sea level by approximately 25 to 50 cm in 2050 and 2100, respectively, as projected by many models, will inundate many parts of the coastal cities of Indonesia. Land subsidence will exacerbate this, increasing the total area that will be permanently inundated. Between 25% and 50% of the total area in coastal cities such as Medan, Jakarta, Semarang, and Surabaya will be permanently underwater. The increase of sea level rise may also inundate the outer islands of the country, thereby reducing the size of Indonesian territory. The analysis suggests that an increase in sea level of 50 cm will not inundate the outer islands of Indonesia permanently, however, combined with tidal patterns in the region, five outer islands will be temporarily inundated. These islands include Alor (next to Timor Leste), Pelampong (next to Singapura), Senua (next to Malaysia), Simuk and Sinyaunyu (next to India).

Some of adaptation measures done in the cities (among other places) to cope with flooding from sea level rise, tides and extreme waves include:

1. Development of dike equipped with polder system to protect area behind the dike (see Figure 4.38);
2. Mangrove rehabilitation in an effort to increase soil surface and reduce wave energy destruction so that the rate of erosion can be reduced; and
3. Practicing fish culture using sylvofishery.



**Figure 4.38 Construction of a dike to reduce flooding from high tides and extreme waves in East Harbour of Muara Bary, Jakarta. Source: Media Indonesia, 2007**

#### 4.4.5 Health

The impact of climate change can be learned from the influence of climate variability, especially related to extreme climatic conditions. The effects of climate variability and change on human life can be grouped into two (Tromp, 1980): (a) direct climate effects (effects on body physiology and psychology) and (b) indirect climate effects (effects on almost all aspects of life and environment that ultimately affect humans). According to Patz *et al.* (2003) from the World Health Organization (WHO), climate change and variability can affect:


1. mortality, physical injury, physical and mental health (directly); and
2. changes in biological processes and ecological impacts on food security such as food sufficiency status and resilience of human body, the quality of water resources, and transmission of infectious diseases and water-borne diseases(indirectly).

Climate change, combined with environmental and behavioral changes in society, causes pressures on public health conditions in Indonesia that have become increasingly difficult to overcome. Indonesia suffers from some vector borne diseases, which may become more severe, especially during extreme climatic periods, as experienced by other developing countries. Transmission periods and distribution areas change and the types of diseases become more varied. Increasing cases and changing patterns of disease distribution during extreme climate periods are an indication of the influence of climate change on public health. Among the ten highest ranked health cases illustrated by the number of patients who were treated at the State Hospitals in Indonesia in 2006 (Table 4.16), outbreaks of dengue fever, malaria, leptospirosis and diarrhea were found to be significantly correlated with the occurrence of extreme climate events associated with ENSO.

**Table 4.16 The ten main of health problems recorded by hospitals in Indonesia in 2006 (Source: Ditjen Pelayanan Medik - Ministry of Health, 2006)**

| No. | Health Problem                       | No. of Cases | % of total Health cases | Magnitude Impact of climate change |
|-----|--------------------------------------|--------------|-------------------------|------------------------------------|
| 1.  | Diarrhea, Gastroenterities infection | 177,517      | 9.95                    | > 50 %                             |
| 2.  | Dengue fever                         | 81,392       | 3.64                    | > 80 %                             |
| 3.  | Tiphoid and paratifoid               | 72,804       | 3.26                    |                                    |
| 4.  | Obstetrics and gynecology problems   | 63,580       | 2,85                    |                                    |
| 5.  | Intracranial injury                  | 48,645       | 2.18                    |                                    |
| 6.  | Fever by unknown cause               | 46,175       | 2.07                    |                                    |
| 7.  | Multiple injury                      | 46,081       | 2.06                    |                                    |
| 8.  | Pneumonia                            | 37,634       | 1.69                    | > 80 %                             |
| 9.  | Malaria                              | 36,865       | 1.65                    | > 90 %                             |
| 10. | Dyspepsia                            | 34,029       | 1.52                    |                                    |

Cases of dengue fever were found to increase significantly during La Niña years (Figure 4.39) when seasonal rainfall levels were above normal. A significant upward trend in the number of dengue cases was also observed in Java, especially in large cities (Figure 4.40). In other cities, such as Dhaka (Bangladesh), cholera cases correspond significantly to local maxima in ENSO, and this climate phenomenon accounts for over 70% of disease variance (Rodo *et al.*, 2002). In Africa, malaria disease outbreaks were triggered by the occurrence of above normal rainfall (Moji *et al.*, 2002).




**Figure 4.39 Incidence rate of dengue fever and the number of affected cities and districts in Note: 1973, 1988 and 1998 were La Niña years. Source: Ministry of Health.**

To predict how climate change will affect vector borne-diseases such as malaria and dengue fever, the following formula from Reisen (1989) was used:

$$c = \frac{ma^2Vp^n}{-\ln p}$$

- c : vectorial capacity, the number of infective bites received daily by a single host
- m: density of vectors in relation to density of host
- a : proportion of vectors feeding on a host divided by the length of the gonotrophic cycle in days
- V : vector competence
- p : daily survival of vectors
- n : the length of the sporogonic cycle(extrinsic incubation period)





**Figure 4.40 Annual trend of dengue incidence rates in districts in Java in period between 1992-2005 (cases per 100,000 people per year). Source: MoE, 2007b.**

The transmission risk or transmission potential was predicted by multiplying the vectorial capacity with a correction factor. The following correction factor for malaria (CF<sub>m</sub>) was developed based on temperature and rainfall using the following formula (Hidayati *et al.*, 2009):

$$CF_m = (T_g * T_s) * (R_s * R_{s-})$$

Where T<sub>g</sub> will be equal to 1 if air temperature is higher than the base temperature of the gonotrophic cycle (18°C), otherwise, T<sub>g</sub> equal to 0. T<sub>s</sub> is equal to 1 if air temperature is higher than the base temperature of the sporogonic cycle (10°C), otherwise, T<sub>s</sub> equal to 0. R<sub>s</sub> is equal to 1 if rainfall is higher than or equal to 25 mm, and equal to R/25 if rainfall is higher than 25 mm. This correction factor was adopted considering the water required for mosquito to lay eggs. For dengue fever, the correction factor was developed based on population density (CF<sub>d</sub>). The CF<sub>d</sub> in a particular region is calculated using the following formula (Hidayati *et al.*, 2009):

$$CF_d = W_d / W_{d\_Mataram}$$

Where W<sub>d</sub> is the weight of population density for that particular region, W<sub>d</sub> is equal to 1 for a region with population density (P) of more than 5000 people/km<sup>2</sup>, 0.75 for a region where 3000 < P < 5000, 0.5 for a region where 1000 < P < 3000 and 1.4 for a region where P < 1000 people/km<sup>2</sup>. W<sub>d</sub>–\_Mataram is the weight of population density of Mataram, a district where a detailed survey on dengue vectors was conducted.

The results of the analysis showed the increasing potential of malaria and dengue fever transmission in 2025 and 2055. The increase in 2055 is not significantly different compared with changes in 2025. Changes in the potential of malaria transmission are predicted to rise much higher than the dengue fever transmission. Under the B1 scenario, changes in the value of transmission potential are predicted to be greater than under the conditions based on the A2 scenario (Table 4.17).


**Table 4.17 Percent change in transmission potential for malaria and dengue fever (Hidayati *et al.*, 2009)**

| Scenario | Malaria |      | Dengue Fever |      |
|----------|---------|------|--------------|------|
|          | 2025    | 2055 | 2025         | 2055 |
| SRESA2   | 19      | 20   | 4            | 4    |
| SRESB1   | 74      | 85   | 13           | 15   |
| Average  | 46.5    | 52.5 | 8.5          | 9.5  |


The average potential transmissions (TP) of dengue fever and malaria at present, from 2020-2030 and from 2050-2060 are shown in Figures 4.41 and 4.42. The spatial differences in TP value is much greater than the difference over time, so the difference depicted in the map over time cannot be easily observed.

To minimize the potential impacts of climate change on public health, adaptation measures need to be developed. In general, adaptation efforts in health can be done via six options. Each option has its own operational level, benefits, feasibility and cost barriers (Mc Michael *et al.* 2003; Table 4.18). Some of the adaptations have already been developed. Haryanto (2008) suggest the following adaptation strategies:

1. Improvement of the disease ecology surveillance system and development of early warning system for outbreaks;
2. Enhancement of capacity building for the government, private sectors, civil society institutions related to the prevention and mitigation of public health aspects affected by climate change;
3. Increasing political awareness of the effects of climate change on public health;
4. Empowering community health service systems for the prevention and control of diseases;
5. Conducting research and developing methods for epidemiology and medicine to break the chains of disease transmissions; and
6. Prevention and eradication of contagious and vector-borne diseases affected by climate change.



**Figure 4.41** Transmission potential of dengue fever under current and future climate (Hidayati *et al.*, 2009). Note: Red represents the highest risk and blue represents the lowest risk



**Figure 4.42 Transmission potential of malaria under current and future climate (Hidayati *et al.*, 2009).  
Note: Red represents the highest risk and blue represents the lowest risk**

The Ministry of Health has developed a short-term climate change adaptation program for 2010-2014 (Health Dept., 2009). This includes (i) building a response system for the impacts of climate change on the health sector, (ii) increasing community access to health facilities and services, especially for the poor, (iii) carrying out training programs on health services for community, (iv) building an emergency response system on disasters and extraordinary events, and (v) carrying out disease prevention and control programs.

As climate change will generally increase the intensity and extent of area under risk and the possibility of the emergence of new diseases related to climate, current resources may not be enough. Some of the important aspects that have to be improved to increase the resilience of the health sector to the current and future climate risks are (i) implementation of existing laws/regulations, (ii) accurate early warning system related to climate-induced health problems, (iii) adequate mass health centers in each susceptible area, (iv) map of spatial susceptibility, (v) sufficient medical officers and paramedics in susceptible areas, (vi) medicines and facilities for diagnoses and treatment of victims, (vii) cooperation between various sectors and institutions both within the country and abroad in order to prevent and overcome health problems, (viii) adequate research and results, (ix) fund for carrying out programs (today, the health budget is

**Table 4.18 General adaptation options to reduce the potential health impacts of climate change**

| Adaptation Options                             | Level | No. of people that benefit | Feasibility | Barriers | Cost |
|--|-------|----------------------------|-------------|----------|------|
| Interagency co-operation                       | G R N | +++                        | ++          | ++       | +    |
| Improvements to public health infrastructure   | NL    | +++                        | +           | +        | ++   |
| Early warning and epidemic forecasting         | NL    | ++                         | ++          | +        | +    |
| Support for infectious disease control         | NL    | ++                         | +++         | +        | +    |
| Monitoring and surveillance                    | L     | ++                         | +++         | +        | +    |
| Integrated environmental management            | L     | +                          | ++          | +        | ++   |
| Urban design (including transport systems)     | L     | +                          | +           | ++       | ++   |
| Housing, sanitation, water quality             | L     | +                          | +           | +        | +    |
| Specific technologies (e.g., air conditioning) | L     | +                          | +++         | +        | +    |
| Public education                               | L     | +++                        | +++         | +        | +    |

Notes: G = Global, R = Regional, N = National, L = Local, + + + = large effect, + + = medium effect, + = small effect (Source: McMichael et al. 2003)

less than 5% of the gross domestic brutto, (x) surveillance systems, (xi) knowledge, awareness and active and full participation of the community and (xi) the capacity to successfully use climate information in managing current and future risks to public health.



## CHAPTER V

# Measures to Mitigate Climate Change

To meet the objectives of the Convention, Indonesia has set up various programs to reduce its emissions. The rate of future greenhouse gas (GHG) emissions under the absence of climate change policy – normally referred to as the base case scenario or business as usual (BAU) – has to be developed. This BAU emissions level will be used to determine how much emissions reduction has been achieved by the country through the implementation of climate change mitigation programs. Baseline emissions should be a key consideration in setting GHG emissions targets as a mitigation strategy to respond to global climate change. At a minimum, the emissions target must be less than the baseline level to induce behavior change and new investment. However, there is substantial uncertainty regarding baseline GHG emissions projection. The use of different assumptions and methodologies may result in different projections. In addition, projecting emissions by sector could not be done independently to some extent. Assumptions used in one sector for developing emissions projections should consider the assumptions used by other sectors. For example, the projected increasing demand for agriculture land (i.e., for rice cultivation) in the future should be captured consistently by the forestry sector in projecting its future deforestation rate. However, in the Second National Communication (SNC), the sectors have not been able to develop integrated emissions projections. The following sections describe briefly the sectoral policy on climate change mitigations and sectoral emissions projections.

## 5.1 Energy Sector

### 5.1.1 Policy Framework

Renewable energy development in Indonesia is supported through National Energy Policy and Energy Law as well as through Green Energy Policy (2004). The Green Energy Policy is basically an effort to develop a “green” energy system characterized by maximum use of renewable energy, efficient use of energy and clean energy technology such as clean coal technology, fuel cell and nuclear energy.

The Government of Indonesia (GOI) has issued a number of policies and regulations that will directly affect the GHG emissions level of the relevant sector. In Energy Law No. 30/2007, there are several provisions, the implementation of which will affect GHG emissions mitigation – for example, provisions that support energy conservation and the development of new and renewable energy through incentive mechanisms. Implementation of the provisions concerning energy conservation and renewable energy development are to be regulated by a government regulation. The operational provision of this government regulation is under preparation (status June 2009).

- a. In 2006, the GOI has launched National Energy Policy under Presidential Decree No. 5/2006 (called PERPRES). One major issue of the policy is the energy supply mix target that was to be achieved in 2006. The energy supply mix target developed was based on domestic resource availability and national energy security objectives. The national energy security objectives are to move away from oil due to declining domestic reserves and maximize the utilization of abundantly available domestic energy resources. More specifically, the energy supply mix target for year 2025 is as follows: To reduce the share of oil from 54.78% (in 2005) to less than 20%
- b. To increase the share of natural gas from 22.2% (in 2005) to 30%
- c. To increase the share of coal from 16.77% (in 2005) to more than 33%
- d. To increase the share of geothermal energy from 2.48% (in 2005) to more than 5%
- e. To increase the share of other new and renewable energy, particularly biomass, nuclear, hydropower, solar and wind to more than 5%
- f. To develop biofuel to achieve a share of at least 5%
- g. To develop liquefied coal to achieve at least 2% (which will start 50,000 barrels per day (bpd) production or 18 million barrels of oil equivalent [MMBOE] in 2020).

It should be noted that although the Government is to increase the share of renewable energy in the national energy supply mix, the share of coal is also to be increased to fulfill the high domestic demand for energy. Therefore, overall, it cannot be guaranteed that total GHG emissions from the supply side will decrease in the future.

In addition to support through the National Energy Policy and Energy Law, renewable energy development in Indonesia is also promoted through the aforementioned Green Energy Policy (2004). The objective of this policy is to develop a “green” energy system that maximizes the use of renewable energy as well as the efficient use of energy and clean energy technology such as clean coal technology, fuel cell and nuclear energy.

In recent developments regarding coal utilization, the Government launched the first crash program for large additional power generation (10,000 MW coal PP), beginning in 2003. The GOI also launched the second crash program for another large additional power generation with 10,000 megawatts (MW) (68% is coal polypropylene [PP], 22% is renewable, and the remaining is 10% is natural gas), beginning in 2007. It can be concluded that in power generation, the contribution from renewable sources will remain low (11% out of the additional 20,000 MW).

The latest policy that supports the development of biofuel in Indonesia is the Biofuel Price Subsidy (2010), where GOI has launched subsidy for biofuel and the obligation of Pertamina (the national oil company) to buy domestic biofuel production. Before the subsidy policy was launched, the Ministry of Energy and Mineral Resources (MEMR) released MEMR Decree No. 32/2008 to regulate (i) supply, utilization and trade of biofuel and (ii) obligation of biofuel utilization in the transport, industry and power sectors. The obligation will be implemented in stages. For example, the biofuel obligation in the transportation sector was 1% in 2009, increasing up to 15% in 2025. This target will be increased every year so that in 2025, the total



contribution of biofuel in the energy supply mix will reach 5% of the total energy supply in Indonesia (i.e., ± 22 billion liters), as written in the National Energy Policy.

The subsequent regulations and policies supporting biofuel development include:

- President Instruction No. 1/2006 concerning biofuel supply and utilization and President Instruction No. 10/2006 concerning the establishment of a national team for biofuels development for poverty alleviation and unemployment reduction; the main objectives of these policies are to reduce dependency on fossil fuels and to strengthen national energy security;
- National Standard (SNI) No. 04-7182-2006 for biodiesel, SNI No. DT27-0001-2006 for bioethanol and Director General for Oil and Gas Decree no. 13483K/24/DJM/2006 concerning Biodiesel Specification for Domestic Market;
- Income tax facility and accelerated amortization for investment in specified business sectors or other areas (Government Regulation No. 1/2007) where renewable energy is included in this category;
- Credit facility for bio-energy development and revitalization of plantations (Minister of Finance Decree No.117/PMK.06/2006) for investment and working capital for activities that support efforts to achieve food and energy security, including biofuel development (Minister of Finance Decree No.79/PMK.05/2007).

To boost the utilization of renewable energy for power generation, the GOI has launched several regulations with provisions related to renewable energy. Specific regulations that govern the sale of electricity generated using renewable energy and/or excess power by an entity to PT PLN (the national electricity enterprise) are:

- MEMR Decree No. 1122/2002 concerning small-scale distributed renewable power generation that regulates small-scale power generation (< 1 MW) using renewable energy, in which, if an entity (i.e., small enterprises) wants to sell electricity generated using renewable energy to PLN (the country's public electric utility), then PLN is obliged to purchase the electricity.
- MEMR Decree No. 02/2006 concerning medium-scale renewable power that regulates medium-scale power generation (1-10 MW) using renewable energy or excess power from captive power plants. If an entity (enterprise) wants to sell electricity generated using renewable energy to PLN (the country's public electric utility), then PLN is obliged to purchase the electricity.
- Government Regulation No. 26/2006 concerning supply and utilization of electricity (revision of Government Regulation No. 10/1989) with new provisions, such as open access

for transmission, direct purchase/without bidding for renewable and excess power or under emergency situations. Procedures for power purchase and transmission rental are covered in the Ministerial Decree No. 001/2006. One of the articles in the decree regulates procedures for the purchase of electricity from renewable sources. PLN is allowed to do direct appointment (without bidding procedures) for the purchase of electricity that is generated using renewable energy. For non-renewable electricity, PLN's power purchase is based on competition (bidding process).

There are also a number of policies/laws related to energy efficiency (EE). However, the current policy (National Policy 2006) does not explicitly state or mention the target of EE achievement, affected market players of EE or main features of EE. There are no incentives or penalties mentioned in the current policies/laws that are related to EE. The latest national energy policy (Presidential Regulation No. 5/2006) mentions the need to pursue energy conservation in all sectors. The policy aims to reach energy elasticity equal to less than 1 by 2025.

The Government is also implementing an energy pricing policy whereby energy prices are to be adjusted gradually to reflect the economic value of energy. Staging and adjustment of energy prices are to be designed in such a way that it would promote and optimize energy diversification. With regard to incentives, the Government is to provide incentives for efforts leading to the implementation of energy conservation and for the development of alternative energy.

### 5.1.2 Emissions Projections and Mitigation Measures

In the SNC, energy supply demand projections are used as sources of activity data in the estimation of GHG emissions. The energy supply demand projection model used in the SNC is the same model used in the Indonesian Energy Outlook 2010-2030 (e.g., Dynamic Model developed by Institut Teknologi Bandung, 2009). The model is used to develop energy supply-demand projections under the BAU scenario and mitigation (CLIMATE) scenario. The model is principally similar to other energy projection models (i.e., the Econometric Model, MARKAL-MAED, LEAP, etc.), where the drivers of energy development are gross domestic product (GDP) and population. The GDP and population growth scenario will determine energy demand growth, which in turn will determine energy supply and GHG emissions level. The relationship between energy supply and demand is determined by energy type and conversion technology.

GDP growth is one of the socioeconomic indicators projected by the Government of Indonesia through BAPPENAS (National Summit, 2009). BAPPENAS has released the target of future Indonesian Economic Development for 2015. The GDP of Indonesia is assumed to grow at an annual rate of 5.5% (2005–2010), 6.6% (2010-2014) and reaching 7.2% in 2015. After 2015, it is rather difficult to determine the GDP growth. Several studies assume the average growth in the GDP will be approximately 7.2% between 2015-2030. This figure is slightly higher than those commonly used by other studies (i.e. about 7%). The estimate of the population growth in the SNC used the estimate from the National Bureau of Statistic (BPS, 2009). The population of Indonesia is estimated to reach 273 million in 2025 (growing at 1.05 %/year, on average, by

2025). Assuming that in the future the population will keep growing at the same rate at 0.94%/year, on average, from 2020-2025, it is projected that the population will reach 286 million by 2030.

Based on the above assumptions, the MEMR developed energy supply/demand projections for Indonesia in the Indonesian Energy Outlook 2010-2030 (Pusdatin-MEMR, 2009b). Three energy projection scenarios from the Indonesian Energy Outlook 2010-2030 were used in the SNC for estimating GHG projections for the energy sector – BAU, mitigation-1 (CLIMATE 1), and mitigation-2 (CLIMATE 2).

The assumptions used in the development of the BAU scenario from the Indonesian Energy Outlook 2010-2030 were developed using existing policies and plans that have been committed by the GOI, namely Crash Programs I and II for Energy Development that have been covered under RUPTL (Business Plan of Electricity Supply) PT PLN 2009-2018 (PT. PLN, 2009), RUKN (General Plan of National Electricity) 2008-2027 (MEMR, 2008), and road map of biofuel development and utilization (although the biofuel development and utilization is assumed still not maximal). Under this BAU scenario, the projected energy supply is shown in Figure 5.1, while the capacity of power plants that need to be developed is presented in Figure 5.2.





Figure 5.1 Energy projections by type of energy under the BAU scenario, 2010-2030



**Figure 5.2 Power plant projections by fuel type under the BAU scenario, 2010-2030**

In the GHG emissions projections under BAU, it was assumed that development will continue according to the historical trend. The BAU scenario shows how GHG emissions will grow over time in the absence of mitigation policy interventions. Natural energy efficiency and existing national (government and private) plans are considered as components of BAU scenario. The resulting emissions from this scenario in 2025 would reach approximately 1.5 carbon dioxide equivalents (CO<sub>2</sub>e), respectively. The rate of annual projected emissions by fuel types and sectors are presented in Figures 5.3 and 5.4, respectively.

The assumptions used for the GHG emissions projections under mitigation scenarios (CLIMATE 1 and 2) are similar to the BAU case with additional assumptions that there will be additional efforts related to energy conservation and new and renewable energy development (i.e., enhance geothermal program, micro-hydro, biofuel, biomass waste to energy, solar PV, wind energy and coal bed methane [CBM]).

Another mitigation scenario used is Presidential Regulation No. 5/2006 - Blue Print PEN 2010, or the PERPRES scenario. The energy supply projection under the PERPRES scenario was developed with an objective to meet energy security. The target of renewable energy in the supply mix is more than 17.93% of the total energy supply. The shift of new and renewable energy from 4.3% in 2005 to 17.93% in 2025 is positive for climate change mitigation, however, the increase in coal share from 19.4% in 2005 to 34% coal and 3.56% liquefied coal by 2025 will negatively




Figure 5.3 Projection of GHG emissions generation by fuel type under the BAU scenario, 2010-2030




Figure 5.4 Projection of GHG emissions generation by sector under the BAU scenario, 2010-2030

affect climate change mitigations so that there is little net decrease in GHG emissions. The energy supply mix under the PERPRES scenario is formulated based on least cost and resource availability.

Both the CLIMATE 1 and CLIMATE 2 mitigation scenarios and PERPRES scenario have excluded advanced clean coal technologies (i.e., integrated gasification combined cycle [IGCC]) and carbon capture and sequestration [CCS] because these technologies are considered to be very expensive. The clean coal technologies considered under both scenarios are fluidized bed, sub-critical and super-critical coal power plants.

The primary energy supply mix, sectoral energy demand, and power plant development projections along with the carbon dioxide (CO<sub>2</sub>) emissions under the BAU, the two mitigation scenarios (CLIMATE 1 and CLIMATE 2), and PERPRES scenario (for comparison) are presented in Table 5.1, Table 5.2, Table 5.3, Figure 5.5, and Figure 5.6. Table 5.1 presents the projection of primary energy supply mix (by fuel type), Table 5.2 presents the projection of final energy demand by sector and Table 5.3 presents demand projection of installed capacity of power generation. These projections are developed using dynamic models with several assumptions, as was discussed previously.

Figures 5.5 and 5.6 show that there are potential reduction in the CO<sub>2</sub> emissions projections under the CLIMATE 1, CLIMATE 2 and PERPRES scenarios compared to the BAU scenario. The potential of CO<sub>2</sub> emissions reductions for each scenario is presented in Figure 5.7. The potential CO<sub>2</sub> emissions reduction by 2020 under the CLIMATE 1 scenario is approximately 0.18 Gton CO<sub>2</sub>e, or approximately 18% of the BAU emissions; under the CLIMATE 2 scenario, the potential emissions reduction is approximately 0.28 Gton CO<sub>2</sub>e, or approximately 28% of the BAU emissions. Under the PERPRES scenario, the CO<sub>2</sub> emissions reductions by 2020 will reach 0.1 Gton CO<sub>2</sub>e, or 10% of BAU emissions. By 2025, the CLIMATE 1, CLIMATE 2, and PERPRES scenarios will result in a 20% reduction (0.3 Gton CO<sub>2</sub>e), 31% reduction (0.47 Gton CO<sub>2</sub>e) and 21% (0.31 Gton CO<sub>2</sub>e), respectively.

**Table 5.1 Projection of energy supply mix under BAU, CLIMATE and PERPRES scenarios (million BOE)**

| Type Energy       | 2010        |                   |                   |              | 2015        |                   |                   |              | 2020        |                   |                   |              | 2025        |                   |                   |              | 2030        |                   |                   |              |   |
|-------------------|-------------|-------------------|-------------------|--------------|-------------|-------------------|-------------------|--------------|-------------|-------------------|-------------------|--------------|-------------|-------------------|-------------------|--------------|-------------|-------------------|-------------------|--------------|---|
|                   | BAU Outlook | CLIMATE 1 Outlook | CLIMATE 2 Outlook | PERPRES 2009 | BAU Outlook | CLIMATE 1 Outlook | CLIMATE 2 Outlook | PERPRES 2009 | BAU Outlook | CLIMATE 1 Outlook | CLIMATE 2 Outlook | PERPRES 2009 | BAU Outlook | CLIMATE 1 Outlook | CLIMATE 2 Outlook | PERPRES 2009 | BAU Outlook | CLIMATE 1 Outlook | CLIMATE 2 Outlook | PERPRES 2009 |   |
| Coal              | 353         | 353               | 353               | 330          | 600         | 462               | 375               | 482          | 858         | 669               | 495               | 725          | 1,412       | 1,078             | 751               | 1,007        | 2,201       | 1,677             | 1,079             | -            |   |
| Coal Liquefaction | -           | -                 | -                 | -            | -           | -                 | -                 | -            | 36          | 36                | -                 | 30           | 110         | 110               | -                 | 91           | 218         | 218               | -                 | -            |   |
| Oil               | 422         | 422               | 422               | 495          | 538         | 515               | 514               | 555          | 699         | 635               | 651               | 627          | 840         | 743               | 786               | 589          | 997         | 841               | 909               | -            |   |
| Natural Gas       | 207         | 207               | 207               | 269          | 302         | 275               | 314               | 349          | 494         | 415               | 460               | 417          | 687         | 527               | 678               | 616          | 841         | 664               | 971               | -            |   |
| CBM               | -           | -                 | -                 | -            | 7           | 7                 | 7                 | 11           | 18          | 18                | 18                | 43           | 64          | 79                | 79                | 96           | 98          | 105               | 105               | -            |   |
| Hydro             | 45          | 45                | 45                | 15           | 65          | 68                | 78                | 23           | 99          | 100               | 117               | 32           | 102         | 110               | 129               | 39           | 110         | 121               | 140               | -            |   |
| Geothermal        | 10          | 10                | 10                | 36           | 36          | 41                | 45                | 90           | 57          | 66                | 69                | 116          | 61          | 74                | 79                | 184          | 66          | 82                | 90                | -            |   |
| Biomassa          | 41          | 41                | 41                | -            | 40          | 39                | 47                | -            | 38          | 35                | 53                | -            | 37          | 41                | 58                | -            | 34          | 52                | 67                | -            |   |
| MSW               | -           | 2                 | -                 | -            | -           | 22                | -                 | -            | -           | 65                | -                 | 1            | -           | 111               | -                 | 1            | -           | 167               | -                 | -            |   |
| Biofuel           | 2           | -                 | 2                 | 34           | 6           | 0                 | 23                | 75           | 15          | 0                 | 66                | 170          | 47          | 0                 | 113               | 298          | 83          | 0                 | 169               | -            |   |
| Solar             | -           | -                 | -                 | -            | -           | -                 | -                 | -            | -           | -                 | 1                 | 1            | -           | 0                 | 1                 | 1            | -           | 0                 | 2                 | -            |   |
| Wind              | -           | -                 | -                 | -            | -           | -                 | -                 | -            | -           | -                 | -                 | -            | -           | -                 | -                 | 1            | -           | -                 | -                 | -            |   |
| Nuclear           | -           | -                 | -                 | -            | -           | -                 | -                 | -            | -           | -                 | -                 | -            | -           | -                 | -                 | -            | -           | -                 | 12                | -            |   |
| Total             | 1,080       | 1,080             | 1,080             | 1,179        | 1,591       | 1,429             | 1,402             | 1,585        | 2,314       | 2,038             | 1,931             | 2,162        | 3,338       | 2,871             | 2,674             | 2,923        | 4,649       | 3,927             | 3,543             | -            |   |
| Efficiency        | -           | -                 | -                 | -            | -           | 10%               | 12%               | -            | -           | 12%               | 17%               | -            | -           | -                 | 14%               | 20%          | -           | -                 | 16%               | 24%          | - |

Notes: Primary energy supply mix under Perpress scenario 2010 – 2025

Table 5.2 Projection of sectoral energy demand under BAU, CLIMATE 1 and CLIMATE 2 scenarios (million BOE)

| Sector Activity     | 2010 |           |           | 2015  |           |           | 2020  |           |           | 2025  |           |           | 2030  |           |           |
|---------------------|------|-----------|-----------|-------|-----------|-----------|-------|-----------|-----------|-------|-----------|-----------|-------|-----------|-----------|
|                     | BAU  | CLIMATE 1 | CLIMATE 2 | BAU   | CLIMATE 1 | CLIMATE 2 | BAU   | CLIMATE 1 | CLIMATE 2 | BAU   | CLIMATE 1 | CLIMATE 2 | BAU   | CLIMATE 1 | CLIMATE 2 |
| Transportation      | 225  | 225       | 225       | 317   | 316       | 316       | 472   | 468       | 467       | 645   | 653       | 624       | 840   | 814       | 780       |
| Industry            | 386  | 386       | 386       | 564   | 512       | 505       | 828   | 691       | 649       | 1,124 | 915       | 863       | 1,423 | 1,138     | 1,076     |
| Households          | 273  | 273       | 273       | 313   | 309       | 309       | 370   | 362       | 362       | 417   | 402       | 402       | 440   | 417       | 417       |
| Commercial          | 47   | 47        | 47        | 54    | 51        | 50        | 68    | 59        | 57        | 87    | 76        | 73        | 127   | 108       | 104       |
| ACM                 | 39   | 39        | 39        | 46    | 46        | 46        | 54    | 54        | 54        | 65    | 65        | 66        | 86    | 86        | 86        |
| Total               | 970  | 970       | 970       | 1,292 | 1,233     | 1,226     | 1,790 | 1,633     | 1,589     | 2,339 | 2,093     | 2,028     | 2,915 | 2,562     | 2,463     |
| Efficient Potential |      |           |           |       |           |           |       |           |           |       |           |           |       |           |           |
| - Transportation    |      |           |           |       | 0%        | 0%        |       | 1%        | 1%        |       | 2%        | 3%        |       | 3%        | 7%        |
| - Industry          |      |           |           |       | 9%        | 10%       |       | 17%       | 22%       |       | 19%       | 23%       |       | 20%       | 24%       |
| - Households        |      |           |           |       | 1%        | 1%        |       | 2%        | 2%        |       | 3%        | 3%        |       | 5%        | 5%        |
| - Commercial        |      |           |           |       | 6%        | 7%        |       | 11%       | 15%       |       | 13%       | 16%       |       | 15%       | 18%       |
| - ACM               |      |           |           |       | 0%        | 0%        |       | 0%        | 0%        |       | 0%        | 0%        |       | 0%        | 0%        |
| TOTAL               |      |           |           |       | 5%        | 5%        |       | 9%        | 11%       |       | 10%       | 13%       |       | 12%       | 16%       |

Table 5.3 Projection of installed capacity demand of power generation under BAU, CLIMATE 1 and CLIMATE 2 scenarios (in MW)

| Sector Activity | 2010   |           |           | 2015   |           |           | 2020   |           |           | 2025    |           |           | 2030    |           |           |
|-----------------|--------|-----------|-----------|--------|-----------|-----------|--------|-----------|-----------|---------|-----------|-----------|---------|-----------|-----------|
|                 | BAU    | CLIMATE 1 | CLIMATE 2 | BAU    | CLIMATE 1 | CLIMATE 2 | BAU    | CLIMATE 1 | CLIMATE 2 | BAU     | CLIMATE 1 | CLIMATE 2 | BAU     | CLIMATE 1 | CLIMATE 2 |
| Steam Coal      | 19,890 | 19,890    | 19,890    | 37,032 | 26,399    | 18,997    | 54,835 | 42,404    | 28,363    | 104,418 | 82,458    | 53,180    | 178,147 | 142,260   | 88,627    |
| Natural Gas     | 12,875 | 12,875    | 12,875    | 15,756 | 14,474    | 19,974    | 25,414 | 21,832    | 30,458    | 31,604  | 26,874    | 49,011    | 37,626  | 31,657    | 76,020    |
| Diesel          | 3,221  | 3,221     | 3,221     | 3,094  | 2,985     | 2,998     | 2,943  | 2,621     | 2,798     | 2,806   | 2,278     | 2,295     | 2,677   | 2,093     | 1,894     |
| Biofuel         | 33     | 33        | 33        | 57     | 184       | 296       | 63     | 463       | 614       | 87      | 569       | 765       | 97      | 629       | 812       |
| Hydro           | 3,746  | 3,746     | 3,746     | 5,382  | 5,702     | 6,513     | 8,211  | 8,313     | 9,750     | 8,479   | 9,177     | 10,737    | 9,171   | 10,091    | 11,651    |
| Geothermal      | 1,093  | 1,093     | 1,093     | 3,875  | 4,359     | 4,776     | 6,060  | 7,053     | 7,333     | 6,475   | 7,866     | 8,463     | 7,055   | 8,706     | 9,550     |
| Agro-Waste      | 100    | 100       | 100       | 95     | 708       | 701       | 172    | 1,237     | 1,186     | 212     | 1,785     | 1,718     | 271     | 2,304     | 2,222     |
| MSW             | -      | -         | -         | 5      | 5         | 5         | 20     | 20        | 20        | 40      | 40        | 40        | 50      | 50        | 50        |
| Landfill Gas    | -      | -         | -         | -      | -         | -         | 3      | 3         | 3         | 10      | 10        | 10        | 20      | 20        | 20        |
| Solar           | 12     | 12        | 12        | 19     | 19        | 31        | 40     | 40        | 96        | 62      | 62        | 178       | 76      | 76        | 288       |
| Wind            | 1      | 1         | 1         | 3      | 3         | 3         | 7      | 7         | 8         | 12      | 12        | 17        | 20      | 20        | 30        |
| Nuclear         |        |           |           |        |           |           |        |           |           |         |           |           |         |           | 4,000     |
| TOTAL           | 40,969 | 40,969    | 40,969    | 65,318 | 54,838    | 54,295    | 97,768 | 83,993    | 80,630    | 154,204 | 131,130   | 126,415   | 235,210 | 197,906   | 195,164   |

Note: There is no power generation projection plan under the PERPRES scenario for 2010-2030






Figure 5.5 GHG emissions projections by fuel type under BAU, CLIMATE and PERPRES scenarios





Figure 5.6 Total GHG emissions projections under BAU, CLIMATE and PERPRES scenarios



**Figure 5.7 GHG emissions reduction potential under BAU, CLIMATE 1, CLIMATE 2, and PERPRES scenarios**

In selecting mitigation measures, factors considered were as follows: (i) sectors that consumed a significant amount of fossil energy, especially with high GHG emissions, such as coal and other fossil fuels, (ii) magnitude of GHG emissions reduction potential of a technology for a given demand and (iii) maturity of the technology to be used for mitigation. Based on these considerations, several mitigation options have been selected, as shown in Table 5.4. Not all of these options are included in the CLIMATE scenarios. The CLIMATE scenarios are developed by considering the domestic resources potential and technological and financial limitations in acquiring more advanced mitigation approaches, such as CCS and advanced clean coal technology.

In addition to that, the selected options also consider that Indonesia is not an Annex I country, and therefore has no international obligation to reduce GHG emissions. However, the GOI released a statement that Indonesia will reduce up to 26% by 2020 with domestic capacity and will further reduce up to 41% by 2020 with international support; this statement is a non-binding GHG emissions reductions statement. It should be noted that the selected mitigation options could be sufficient to achieve the above mentioned emissions reduction statement (26% by 2020). Emissions reductions, which can be achieved from the selected mitigation options under the CLIMATE scenario, are the differences between the emissions projections under the BAU scenario and the emissions projections under the CLIMATE scenarios. The GHG emissions difference level between the BAU scenario and CLIMATE scenarios is more appropriately referred to as “emissions avoidance.”

**Table 5.4 Selected mitigation options**

| Sector                 | Sub-sector                            | Mitigation Options  |
|------------------------|---------------------------------------|---|
| <b>Energy Supply</b>   | Power generation                      | <ul style="list-style-type: none"> <li>a. Include the energy system with fewer GHG emissions that has not been selected in BAU, such as biomass-based power (electricity generation from municipal solid waste [MSW] incineration, , landfill gas (LFG), liquid and solid biomass waste, etc.), solar, wind, etc., although the share of these types of energy is still relatively very small;</li> <li>b. Increase the share of new and renewable energy that has been included in BAU, but is still below the level of PERPRES target, such as geothermal and nuclear energy;</li> <li>c. Increase the use of efficient technology in power plants, such as circulated coal fluidized bed combustion (CFBC), sub- or super-critical power plants, coal IGCC, gas combined cycle, combined heat and power;</li> <li>d. Develop technology that is appropriate for utilizing fuels with low GHG emissions (i.e., natural gas); and</li> <li>e. CCS technology application for large coal power plants.</li> </ul> |
|                        | Oil and gas production and refineries | Gas flaring and venting reduction   |
| <b>Energy End User</b> | Transportation                        | <ul style="list-style-type: none"> <li>a. Increase biofuel development efforts for personal vehicles</li> <li>b. Increase the use of compressed natural gas and liquified petroleum gas (LPG) in transportation</li> <li>c. Introduce fuel cell vehicles and electric cars/motorcycles</li> <li>d. Promote mass rapid transport (MRT)</li> <li>e. Traffic management, etc. (Appendix 3 presented a policy and air control program issued by the Directorate General of Land Transportation, Ministry of Transportation to reduce air pollution due to exhaust gas from vehicles)</li> </ul>   |
|                        | Industry                              | <ul style="list-style-type: none"> <li>a. Increase the use of more EE equipment or technology (i.e., more efficient conversion process, more efficient combustion system, variable speed electric motors, etc.)</li> <li>b. Increase the use of more efficient material conversion process</li> <li>c. Increase the use of recycled materials, etc.</li> <li>d. Introduce co-processing or co-firing technology</li> </ul>  |
|                        | Residential and commercial end uses   | BAU, CLIMATE 1 and CLIMATE 2 scenarios and is therefore also considered as a mitigation option  |


Under the CLIMATE mitigation scenarios, there are three important options to reduce GHG emissions levels: (i) efficiency energy (demand side as well as power supply side), (ii) promoting biofuel to replace the utilization of fossil fuels (refinery products) and (iii) the utilization of renewable energy in power generation (geothermal, biomass based fuels, hydro, etc.). Detailed mitigation scenarios and investment costs will be discussed in the following sub-sections.

### 5.1.3 Mitigation Costs

The estimated investment costs needed for implementation of the mitigation options includes (i) development of biofuel production to supply energy demand in the Indonesian transportation sector, (ii) development of advanced coal power plants (more efficient) and less GHG emitting power plants (i.e., natural gas power generation), (iii) develop other renewable energy (e.g., geothermal, hydropower, biomass-based power plants, municipal solid waste [MSW]-based power plants, etc.) and (iv) increasing efficiency on the energy demand side.

#### Biofuel Production for the Transportation and Industrial Sectors

Recently, biofuels have become one of the most important forms of energy in Indonesia, particularly in the transportation and industry sectors. According to the GOI's plan, the future demand of biofuel in the transportation and industry sectors will be supplied from domestic production. Figure 5.8 presents the demand of biofuel in Indonesia and total investment for the production systems. The total investment cost is estimated based on the investment cost of 30 million USD per 300,000 tons per year. Table 5.5 presents additional investment costs and the potential reduction of GHG emissions.



**Figure 5.8 Total projected demand of biofuel in the transportation and industrial sectors and estimated investment cost of the biofuel production system**

**Table 5.5 Additional demand of biofuel and GHG emissions reduction potential and cost**

| Year | Biofuel Add. Demand (from 2010), BE0 |             |             | Reduction Potential, Ton CO <sub>2</sub> e |            | Additional Cost, Million USD |           | Av. Reduction Cost, USD/ton |           |
|------|--------------------------------------|-------------|-------------|--|------------|------------------------------|-----------|-----------------------------|-----------|
|      | BAU                                  | CLIMATE 1   | CLIMATE 2   | CLIMATE 1                                  | CLIMATE 2  | CLIMATE 1                    | CLIMATE 2 | CLIMATE 1                   | CLIMATE 2 |
| 2015 | 3,095,690                            | 17,579,525  | 17,541,791  | 6,314,952                                  | 6,298,500  | 203                          | 202       | 32.11                       | 32.11     |
| 2020 | 12,079,552                           | 54,632,308  | 54,197,991  | 12,238,050                                 | 12,065,140 | 393                          | 387       | 32.11                       | 32.11     |
| 2025 | 43,808,831                           | 98,695,064  | 98,100,635  | 5,377,396                                  | 5,307,587  | 173                          | 170       | 32.11                       | 32.11     |
| 2030 | 79,658,232                           | 155,583,732 | 154,842,779 | 9,173,120                                  | 9,109,236  | 295                          | 292       | 32.11                       | 32.11     |

Advanced (more efficient) coal power plants

Table 5.6 presents new investments for advanced (more efficient) coal power plants under the BAU, CLIMATE1 and CLIMATE 2 scenarios. The size and investments made for new additional power plants under CLIMATE 1 and 2 are both lower than under the BAU scenario. The reduced use of coal plants is the result of two factors: reduced demand for power plant capacity due to conservation efforts and the shift of power plants from coal-based to natural gas power plants (i.e, less GHG emitting power plants) and to renewable power plants (geothermal, hydropower, solar, wind, biofuel, biomass wastes). The table also shows that new coal power plants in Indonesia in the near future (until 2030) will consist of two types: advanced coal (efficiency of 38%) and conventional coal (efficiency of 32%). As a result of reduced use of coal power plants and the use of advanced coal power plants GHG emissions from coal power plants for the CLIMATE 1 and CLIMATE 2 scenarios are both lower than that of the BAU.

**Table 5.6 New investments for advanced (more efficient) coal power plants under the BAU, CLIMATE1 and CLIMATE 2 scenarios**

| Year | BAU           |                      |              |             |                            | CLIMATE 1     |                |              |             |                            | CLIMATE 2     |                |          |             |                            |
|------|---------------|----------------------|--------------|-------------|----------------------------|---------------|----------------|--------------|-------------|----------------------------|---------------|----------------|----------|-------------|----------------------------|
|      | Installed Cap | New power plant, M/V |              | Investment  | Emission                   | Installed Cap | New plant, M/V |              | Investment  | Emission                   | Installed Cap | New plant, M/V |          | Investment  | Emission                   |
|      |               | Advanced             | Conventional |             |                            |               | Advanced       | Conventional |             |                            |               | MW             | Advanced |             |                            |
|      | MW            | Eff 38%              | Eff 32%      | million USD | mill.ton CO <sub>2</sub> e | MW            | Eff 38%        | Eff 32%      | million USD | mill.ton CO <sub>2</sub> e | Climate 2     | Eff 38%        | Eff 32%  | million USD | mill.ton CO <sub>2</sub> e |
| 2010 | 19,890        |                      |              |             | 133                        | 19,890        |                |              |             | 133                        | 19,890        |                |          |             | 133                        |
| 2011 | 24,131        | -                    | 4,241        | 5,089       | 161                        | 19,890        | -              | -            |             | 110                        | 19,890        |                |          |             | 98                         |
| 2012 | 28,630        | -                    | 4,499        | 5,399       | 191                        | 19,890        | -              | -            |             | 125                        | 19,890        |                |          |             | 108                        |
| 2013 | 32,249        | -                    | 3,619        | 4,343       | 215                        | 22,359        | -              | 2,469        | 2,962       | 149                        | 19,890        |                |          |             | 117                        |
| 2014 | 35,043        | -                    | 2,794        | 3,353       | 234                        | 23,897        | 1,539          | -            | 3,077       | 155                        | 19,890        |                |          |             | 118                        |
| 2015 | 37,032        | -                    | 1,989        | 2,387       | 247                        | 26,399        | 2,502          | -            | 5,004       | 171                        | 19,890        |                |          |             | 123                        |
| 2016 | 38,896        | -                    | 1,864        | 2,237       | 260                        | 28,329        | 695            | 1,235        | 2,872       | 183                        | 19,949        | 59             | -        | 118         | 129                        |
| 2017 | 40,575        | -                    | 1,679        | 2,015       | 271                        | 31,024        | 269            | 2,425        | 3,449       | 201                        | 21,052        | 1,103          | -        | 2,206       | 136                        |
| 2018 | 42,782        | -                    | 2,207        | 2,648       | 285                        | 33,801        | 278            | 2,499        | 3,554       | 219                        | 23,153        | 2,101          | -        | 4,201       | 150                        |
| 2019 | 48,609        | 4,079                | 1,748        | 10,256      | 319                        | 38,050        | 4,250          | -            | 8,499       | 243                        | 25,648        | 2,495          | -        | 4,990       | 163                        |
| 2020 | 54,835        | 623                  | 5,603        | 7,968       | 360                        | 42,404        | 1,089          | 3,266        | 6,096       | 270                        | 28,363        | 1,494          | 1,222    | 4,454       | 181                        |
| 2021 | 64,063        | 923                  | 8,305        | 11,812      | 421                        | 49,724        | 1,757          | 5,563        | 10,188      | 317                        | 32,683        | 864            | 3,455    | 5,874       | 208                        |
| 2022 | 73,482        | 5,652                | 3,768        | 15,825      | 475                        | 57,070        | 6,612          | 735          | 14,106      | 358                        | 37,473        | 4,311          | 479      | 9,197       | 235                        |
| 2023 | 83,145        | 1,933                | 7,730        | 13,142      | 538                        | 65,110        | 8,040          | -            | 16,080      | 397                        | 42,579        | 5,106          | -        | 10,213      | 260                        |
| 2024 | 92,172        | 1,805                | 7,221        | 12,276      | 596                        | 72,470        | 7,360          | -            | 14,721      | 442                        | 47,345        | 4,766          | -        | 9,532       | 289                        |
| 2025 | 104,418       | 2,449                | 9,797        | 16,654      | 676                        | 82,458        | 7,990          | 1,997        | 18,376      | 503                        | 53,180        | 4,376          | 1,459    | 10,502      | 324                        |
| 2026 | 115,946       | 2,306                | 9,222        | 15,678      | 750                        | 91,842        | 3,754          | 5,631        | 14,265      | 560                        | 59,492        | 2,840          | 3,472    | 9,847       | 363                        |
| 2027 | 129,345       | 2,680                | 10,720       | 18,223      | 837                        | 102,487       | 5,323          | 5,323        | 17,032      | 625                        | 66,350        | 3,429          | 3,429    | 10,973      | 405                        |
| 2028 | 142,912       | 2,713                | 10,854       | 18,451      | 925                        | 113,712       | 5,612          | 5,612        | 17,959      | 694                        | 72,731        | 3,191          | 3,191    | 10,210      | 428                        |
| 2029 | 159,556       | 3,329                | 13,315       | 22,636      | 1,032                      | 127,034       | 6,661          | 6,661        | 21,315      | 775                        | 80,553        | 3,911          | 3,911    | 12,515      | 476                        |
| 2030 | 178,147       | 3,718                | 14,872       | 25,283      | 1,153                      | 142,260       | 7,613          | 7,613        | 24,361      | 868                        | 88,627        | 4,037          | 4,037    | 12,918      | 510                        |

### Less GHG-emitting power plants

Table 5.7 presents the installed capacity, additional investment and GHG emissions of natural gas power plants from 2010-2030. The installed capacity of natural gas power plants in the CLIMATE 1 scenario is less than that in the BAU scenario. This occurs because the demand of electricity in the CLIMATE 1 scenario is lower than that of the BAU and there is not much effort to shift from coal to natural gas plants. However, as a result of greater efforts to use less GHG emitting plants, the installed capacity of natural gas plants under the CLIMATE 2 scenario is much greater than that of the BAU, especially after 2020. Due to the higher installed capacity of natural gas plants, the required investment and the GHG emissions in the CLIMATE 2 scenario are higher than that of the BAU. Although the GHG emissions of gas power plants are higher in CLIMATE 2, the overall emissions from the power sector in CLIMATE 2 are lower than that of the BAU because the increase in emissions from natural gas plants is compensated by the decrease in emissions from coal power plants.

**Table 5.7 The installed capacity, additional investment and GHG emissions of natural gas power plants from 2010-2030**

| YEAR | BAU               |                                    |                                       | CLIMATE 1         |                                    |  | CLIMATE 2         |                                    |  |
|------|-------------------|------------------------------------|---------------------------------------|-------------------|------------------------------------|--|-------------------|------------------------------------|--|
|      | Installed Cap, MW | Investment of new plant, Mill. USD | Emissions mill. Ton CO <sub>2</sub> e | Installed Cap, MW | Investment of new plant, Mill. USD | Emissions, mill. Ton CO <sub>2</sub> e | Installed Cap, MW | Investment of new plant, Mill. USD | Emissions, mill. Ton CO <sub>2</sub> e |
| 2010 | 12,875            |                                    | 30                                    | 12,875            | -                                  | 30                                     | 12,875            | -                                  | 30                                     |
| 2011 | 12,988            | 90                                 | 30                                    | 12,875            | -                                  | 28                                     | 13,560            | 548                                | 31                                     |
| 2012 | 14,008            | 816                                | 32                                    | 13,070            | 156                                | 31                                     | 15,353            | 1,434                              | 36                                     |
| 2013 | 14,008            | -                                  | 32                                    | 13,070            | -                                  | 30                                     | 16,993            | 1,312                              | 39                                     |
| 2014 | 14,008            | -                                  | 32                                    | 13,569            | 399                                | 31                                     | 18,534            | 1,233                              | 43                                     |
| 2015 | 15,756            | 1,398                              | 37                                    | 14,474            | 725                                | 34                                     | 19,974            | 1,152                              | 46                                     |
| 2016 | 18,094            | 1,870                              | 42                                    | 15,522            | 838                                | 36                                     | 21,519            | 1,236                              | 50                                     |
| 2017 | 20,183            | 1,671                              | 47                                    | 16,735            | 970                                | 39                                     | 23,518            | 1,599                              | 54                                     |
| 2018 | 21,481            | 1,038                              | 50                                    | 18,668            | 1,546                              | 43                                     | 25,771            | 1,803                              | 60                                     |
| 2019 | 23,544            | 1,650                              | 55                                    | 20,340            | 1,337                              | 47                                     | 28,034            | 1,810                              | 65                                     |
| 2020 | 25,414            | 1,496                              | 59                                    | 21,832            | 1,194                              | 51                                     | 30,458            | 1,940                              | 71                                     |
| 2021 | 26,252            | 670                                | 61                                    | 22,480            | 518                                | 52                                     | 33,657            | 2,559                              | 78                                     |
| 2022 | 27,864            | 1,290                              | 65                                    | 23,805            | 1,060                              | 55                                     | 37,278            | 2,897                              | 86                                     |
| 2023 | 29,313            | 1,159                              | 68                                    | 25,021            | 973                                | 58                                     | 40,979            | 2,961                              | 95                                     |
| 2024 | 31,089            | 1,421                              | 72                                    | 26,484            | 1,171                              | 61                                     | 44,683            | 2,963                              | 104                                    |
| 2025 | 31,604            | 412                                | 73                                    | 26,874            | 312                                | 62                                     | 49,011            | 3,463                              | 114                                    |
| 2026 | 33,539            | 1,548                              | 78                                    | 28,478            | 1,283                              | 66                                     | 53,584            | 3,658                              | 124                                    |
| 2027 | 35,756            | 1,773                              | 83                                    | 30,303            | 1,460                              | 70                                     | 58,711            | 4,102                              | 136                                    |
| 2028 | 36,700            | 755                                | 85                                    | 30,982            | 543                                | 72                                     | 63,680            | 3,975                              | 148                                    |
| 2029 | 37,644            | 755                                | 87                                    | 31,690            | 567                                | 73                                     | 69,801            | 4,896                              | 162                                    |
| 2030 | 37,644            | -                                  | 87                                    | 31,690            | -                                  | 73                                     | 76,020            | 4,976                              | 176                                    |

Renewable energy (e.g., geothermal, hydro, biomass-based PP, MSW-based PP, etc.)

Table 5.8 presents the installed capacity and the required investment for renewable power plants BAU, CLIMATE 1 and CLIMATE 2 scenarios. The installed capacity of renewable power plants under the CLIMATE 1 and 2 scenarios is higher than that under the BAU scenario. Conventional renewables such as hydropower and geothermal power plants under the two CLIMATE scenarios are not much different from the BAU due to geographical limitation of resource availability with respect to demand center. For example, although the water resources

available for power is relatively high (up to 75 GW), many of the resource locations (i.e., 25 GW in Papua) do not match with the location of demand center (i.e., Jawa, Sumatera). Similar situations also occur with regard to geothermal resources (26 GW). The significant difference between the two CLIMATE scenarios and BAU is that other renewables such as solar power, wind, biomass based power plants are utilized. However, these plants are still very small compared to the total required power plant capacity and therefore their contribution towards reducing GHG emissions is not significant. Further deployment of these types of power plants are limited because such technologies are much more expensive than conventional power plants. Further development and utilization of wind power is limited because wind resources in Indonesia are not sufficient (i.e., low speed winds).

**Table 5.8 The installed capacity and the required investment for renewable power plants under the BAU, CLIMATE 1 and CLIMATE 2 scenarios**

| Year | BAU               |            |             |                       |            |             | CLIMATE 1         |            |             |                       |            |             | CLIMATE 2         |            |             |                       |            |             |
|------|-------------------|------------|-------------|-----------------------|------------|-------------|-------------------|------------|-------------|-----------------------|------------|-------------|-------------------|------------|-------------|-----------------------|------------|-------------|
|      | Installed Cap, MW |            |             | Investment, Mill. USD |            |             | Installed Cap, MW |            |             | Investment, Mill. USD |            |             | Installed Cap, MW |            |             | Investment, Mill. USD |            |             |
|      | Hydro             | Geothermal | Other renew | Hydro                 | Geothermal | Other renew | Hydro             | Geothermal | Other renew | Hydro                 | Geothermal | Other renew | Hydro             | Geothermal | Other renew | Hydro                 | Geothermal | Other renew |
| 2010 | 3,745             | 1,093      | 145         |                       |            |             | 3,745             | 1,093      | 145         |                       |            |             | 3,745             | 1,093      | 145         |                       |            |             |
| 2011 | 3,983             | 1,370      | 154         | 476                   | 831        | 24          | 3,745             | 1,093      | 145         |                       |            |             | 4,332             | 1,695      | 360         | 1,173                 | 1,806      | 428         |
| 2012 | 4,243             | 2,114      | 169         | 520                   | 2,232      | 30          | 4,368             | 2,333      | 456         | 1,247                 | 3,719      | 623         | 4,877             | 2,460      | 458         | 1,091                 | 2,294      | 218         |
| 2013 | 4,270             | 3,009      | 175         | 54                    | 2,685      | 13          | 4,434             | 3,127      | 620         | 132                   | 2,384      | 302         | 5,229             | 3,221      | 633         | 703                   | 2,284      | 337         |
| 2014 | 5,307             | 3,819      | 176         | 2,074                 | 2,430      | 7           | 5,367             | 4,000      | 769         | 1,865                 | 2,617      | 249         | 6,077             | 4,051      | 794         | 1,696                 | 2,490      | 279         |
| 2015 | 5,382             | 3,875      | 185         | 150                   | 168        | 49          | 5,702             | 4,359      | 919         | 671                   | 1,077      | 290         | 6,513             | 4,776      | 1,036       | 873                   | 2,176      | 394         |
| 2016 | 6,167             | 4,087      | 194         | 1,570                 | 636        | 51          | 6,492             | 4,722      | 1,083       | 1,580                 | 1,088      | 315         | 7,619             | 4,944      | 1,252       | 2,211                 | 502        | 395         |
| 2017 | 7,139             | 4,783      | 199         | 1,944                 | 2,088      | 39          | 7,338             | 5,278      | 1,227       | 1,691                 | 1,669      | 254         | 8,661             | 5,753      | 1,392       | 2,084                 | 2,428      | 320         |
| 2018 | 8,067             | 5,650      | 217         | 1,856                 | 2,601      | 83          | 8,132             | 5,940      | 1,431       | 1,589                 | 1,986      | 417         | 9,303             | 6,271      | 1,604       | 1,283                 | 1,553      | 455         |
| 2019 | 8,146             | 5,998      | 284         | 159                   | 1,044      | 164         | 8,198             | 6,496      | 1,586       | 132                   | 1,670      | 293         | 9,543             | 6,785      | 1,768       | 482                   | 1,542      | 408         |
| 2020 | 8,211             | 6,060      | 306         | 128                   | 187        | 83          | 8,313             | 7,053      | 1,769       | 230                   | 1,671      | 327         | 9,750             | 7,333      | 1,927       | 413                   | 1,644      | 344         |
| 2021 | 8,318             | 6,158      | 344         | 216                   | 293        | 121         | 8,511             | 7,354      | 1,928       | 396                   | 903        | 349         | 10,097            | 7,529      | 2,082       | 694                   | 588        | 454         |
| 2022 | 8,335             | 6,192      | 365         | 34                    | 103        | 76          | 8,647             | 7,630      | 2,108       | 272                   | 826        | 364         | 10,247            | 7,807      | 2,288       | 299                   | 835        | 457         |
| 2023 | 8,338             | 6,253      | 381         | 5                     | 184        | 59          | 8,785             | 7,729      | 2,155       | 276                   | 297        | 113         | 10,458            | 8,004      | 2,355       | 423                   | 590        | 206         |
| 2024 | 8,479             | 6,359      | 402         | 282                   | 317        | 97          | 9,029             | 7,705      | 2,319       | 487                   | -          | 355         | 10,650            | 8,219      | 2,553       | 383                   | 646        | 469         |
| 2025 | 8,479             | 6,475      | 422         | 1                     | 348        | 66          | 9,177             | 7,866      | 2,478       | 296                   | 483        | 341         | 10,737            | 8,463      | 2,728       | 174                   | 733        | 461         |
| 2026 | 8,553             | 6,540      | 442         | 147                   | 195        | 97          | 9,255             | 7,974      | 2,549       | 158                   | 325        | 149         | 10,854            | 8,518      | 2,792       | 234                   | 166        | 228         |
| 2027 | 8,618             | 6,601      | 463         | 131                   | 183        | 72          | 9,480             | 8,159      | 2,741       | 448                   | 554        | 395         | 10,934            | 8,687      | 2,996       | 159                   | 505        | 503         |
| 2028 | 8,927             | 6,745      | 495         | 618                   | 431        | 101         | 9,546             | 8,206      | 2,902       | 133                   | 142        | 363         | 11,152            | 8,889      | 3,181       | 437                   | 607        | 527         |
| 2029 | 9,087             | 6,923      | 510         | 319                   | 535        | 51          | 9,835             | 8,469      | 2,926       | 578                   | 789        | 61          | 11,457            | 9,342      | 3,228       | 610                   | 1,358      | 259         |
| 2030 | 9,171             | 7,055      | 534         | 169                   | 395        | 69          | 10,091            | 8,706      | 3,119       | 512                   | 710        | 411         | 11,651            | 9,550      | 3,438       | 388                   | 625        | 620         |

**Increasing efficiency on the energy demand side**

Improvements in energy efficiency will be approximately 17% in 2020, 20% in 2025 and 23.8% in 2030. The efficiency potential in power generation is approximately 18%, whereby the efficiency potential on the demand side (i.e., household, commercial and industry) will reach 11% in 2020, 13% in 2025 and 16% in 2030. The detailed energy efficiency of each sector from 2020-2030 is presented in Table 5.9.

Table 5.9 Energy efficiency improvements by sector, 2020-2030

| Key Sources of Activity  | 2010 - 2025       |            |                              |                          |  |                 |              |                              |                          |                                  |
|--------------------------|-------------------|------------|------------------------------|--------------------------|--|-----------------|--------------|------------------------------|--------------------------|----------------------------------|
|                          | CLIMATE 1         |            |                              |                          |  | CLIMATE 2       |              |                              |                          |                                  |
|                          | Efficiency, BOE % | Efficiency | Investment Cost, Million USD | Cost Saving, Million USD | Reduction Potential, Ton CO <sub>2</sub> e | Efficiency, BOE | % Efficiency | Investment Cost, Million USD | Cost Saving, Million USD | Reduction, Ton CO <sub>2</sub> e |
| Transportation           | 50,946,486        | 1%         | 16                           | 3,645                    | 116  | 84,727,097      | 1%           | 21                           | 6,062                    | 179                              |
| Industry                 | 1,546,393,912     | 17%        | 115                          | 110,644                  | 22   | 1,923,871,786   | 22%          | 150                          | 137,653                  | 23                               |
| Households (w/o biomass) | 100,916,073       | 6%         | 10                           | 267                      | 14   | 100,916,073     | 6%           | 10                           | 7,221                    | 26                               |
| Commercial               | 86,618,531        | 11%        | 23                           | 230                      | 25   | 106,718,845     | 15%          | 30                           | 7,636                    | 49                               |
| ACM                      | -                 |            |                              |                          |  | -               |              |                              |                          |                                  |
| Total                    | 1,784,875,002     | 10%        |                              | 114,787                  | 177  | 2,216,233,802   |              |                              |                          | 277                              |

## 5.2 Industry Sector

### 5.2.1 Policies Framework

The major GHG emissions generated from industrial processes are CO<sub>2</sub>, methane (CH<sub>4</sub>), nitrous oxide (N<sub>2</sub>O) and perfluorocarbons (PFCs) in the form of CF<sub>4</sub> and C<sub>2</sub>F<sub>6</sub>; CO<sub>2</sub> emissions account for the vast majority, representing 93% of the total GHG emissions from industrial processes. However, current Indonesian regulations and policies do not directly control CO<sub>2</sub> and other GHG emissions from industrial processes.

In terms of legal aspect, these gases are still allowable under Indonesian regulations. Refer to the Decree of the State Minister of Environment/Kep. MENLH No. 129 of 2003 concerning standard quality of emissions for industrial processes; the requirement for stationary emissions sources from the industrial processes are opacity, hydrogen sulfide (H<sub>2</sub>S) and other hydrocarbon gases, which does not cover the CO<sub>2</sub> or other GHG emissions. There is no specific threshold value for these types of emissions from the industrial sector.

### 5.2.2 Emissions Projections and Mitigation Measures

The total number of medium- and large-scale industries of varying types in Indonesia is more than 20,000. However, not all of the GHG emissions from those industries are covered in the SNC. The SNC only includes GHG emissions specifically associated with industrial processes and excludes emissions from energy consumption in industrial activities (which are already covered in the energy sector). The types of industrial processes covered in the SNC are industries that have the potential to generate GHGs, including mineral production processes (e.g., cement, lime, carbonate products), chemical production processes (e.g., ammonia, nitric acid, caprolactam),



and metal production processes (e.g., iron/steel production and aluminum smelter), as covered in the 2006 IPCC Guidelines.

The methodology of GHG emissions projections for industrial processes is principally similar to other sectors, as emissions are based on projected activity data. The projections used simple econometric extrapolation techniques taking into consideration the historical trends. Activity data projections are based on projection models, which use GDP growth of the industrial sector as the growth driver. Although the average national GDP growth from 2000-2005 was only 4.7% per year, the GDP growth from the industrial sector was still high at 9.3% per annum. Considering the development that has occurred in the past five years, future GDP from the industrial sector is projected to grow at a slower rate of 6.4% per year. Future GHG emissions generated from the industrial sector are estimated to increase, consistent with the projected GDP growth of the industrial sector at 6.4% per year.

The industrial sector will reduce the rate of emissions from the BAU scenario by increasing the efficiency in production processes, introduction of new technologies, or by changing the raw materials (i.e., using waste as an alternative material in the cement industry). The emissions reductions can also be achieved through the clean development mechanism (CDM) scheme and private sector participation. Public funding is still needed for the dissemination of programs related to CO<sub>2</sub> emissions reductions in industrial processes.

In the analysis of future emissions from the industrial sector, two scenarios have been established by assuming that emissions reductions are carried out by the private sector through the CDM scheme and supported by government dissemination programs and other international funds/grants. Table 5.10 presents the projected emissions generated by the industrial sector from 2010-2025 under BAU and the two alternative scenarios.

**Table 5.10 Projected emissions from the industrial sector under BAU and mitigation scenarios (Ggram CO<sub>2</sub>e)**

| Scenario  | 2010   | 2015   | 2020   | 2025   |
|---|--------|--------|--------|--------|
| BAU   | 52,850 | 57,296 | 62,117 | 67,343 |
| Private scenario through CDM  | 49,438 | 52,422 | 56,754 | 61,493 |
| Reduction Potential   | 3,412  | 4,875  | 5,362  | 5,850  |
| % Reduction   | 6.5%   | 8.5%   | 8.6%   | 8.7%   |
| Private scenario through CDM with support from the government through dissemination program and other international funds | 46,152 | 49,136 | 53,469 | 58,207 |
| Reduction Potential   | 6,698  | 8,160  | 8,648  | 9,135  |
| % Reduction   | 12.7%  | 14.2%  | 13.9%  | 13.6%  |

The potential for GHG emissions reductions are estimated to occur through the following processes (i) process improvement; (ii) operation system improvement; (iii) technology change; (iv) raw material substitution; and (v) dissemination/promotion program.

## 5.3 Forestry Sector

### 5.3.1 Policy Framework

Mitigation of climate change is one of the long-term national forestry plans for the period from 2010-2029 (MoF, 2008). Five key policies that support mitigation of climate change in the forestry sector have been issued, as stated in Forestry Strategic Plan for 2005-2009 through Ministerial Decree No. P.04/Menhut-II/2005 dated on 14 February 2005. These include the following:

1. **Combating illegal logging and its associated trade**, through (i) securing forest areas and (ii) controlling forest product administration;
2. **Revitalization of the forestry sector, particularly the forestry industries**, through (i) natural production forest management under non-concessionaire license; (ii) development of plantation forest; (iii) management (utilization) of primary production forest; and (iv) restructuring of the primary forestry industry;
3. **Conservation and rehabilitation of forest resources**, through (i) plantation seeds development, (ii) management of watershed areas; (iii) forest and land rehabilitation; (iv) self-management of forest and land rehabilitation; (v) development of national park management; (vi) management of nature preservation/sanctuary reserve/hunting park areas; (vii) controlling forest fires; (viii) management of biological diversity; (ix) management of protected forest; and (x) utilization of wildlife and plant products and environmental services;
4. **Empowering the economy of the community within and surrounding the forest area**, through (i) development of community forest and community plantation forest; (ii) development of non timber forest product (NTFP) utilization; (iii) development of buffer zone area surrounding conservation areas; and (iv) development of social forestry; and
5. **Stabilization of forest area for promoting and strengthening sustainable forest management**, through (i) inventory and mapping of forest resources; (ii) development of Information Assessment System on Forestry Development; (iii) gazettement of forest area, (iv) preparation and evaluation of the utilization and conversion of forest area; and (v) development of Forest Management Units (FMUs).

In 2009, Ministry of Forestry issued eight policy strategies for forestry sector through Ministerial Decree No.70/Menhut-II/2009 dated 7 December 2009, as the basic for forestry activities in 2009-2014. Mitigation and adaptation of climate change for forestry sector are clearly stated as one of the strategies that include:

1. Area establishment,
2. Forest rehabilitation and improvement the carrying capacity of watershed area,
3. Forest protection and forest fire control

4. Conservation of biodiversity
5. Revitalization of forest utilization and forestry industries,
6. Empowerment of forest community and forest industries,
7. Mitigation and adaptation of climate change for forestry sector,
8. Strengthening the forestry institutions.

Consistent with the above programs, targets for the long-term national forestry strategic plans include: (1) Confirmation of 60 million hectares of forest area by 2014; (2) Rehabilitation and establishment of new plantations with the following targets for 2012: industrial forest plantations (7.2 million ha), community plantation forest (5.4 million ha), community forest (4 million ha) and social forestry (2 million ha); (3) Forest protection and natural resources conservation (120.3 million ha); (4) Development of economic and social functions; (5) Establishment of institutions; and (6) Research and development.

The legal and policy framework already exists through which Indonesia could undergo a significant reduction in future carbon emissions. Key pieces of legislation include:

- Government Regulation (PP) No. 35/1995 on reforestation fund;
- Government Regulation (PP) No. 34/2002 on land use, forest management planning, forest and forest land use;
- Government Regulation (PP) No. 45/2004 on forest protection;
- Government Regulation (PP) No. 6/2006 on forest management and utilization;
- Government Regulation (PP) No. 6/2007 on forest planning management and forest utilization;
- Government Regulation (PP) No. 10/2010 on procedure for exchanging forest utilization and forest function
- Government Regulation (PP) No. 11/2010 on rules on the optimizing the use of abandoned land
- Presidential Instruction (Inpres) No. 4/2005 on combating illegal logging;
- Presidential Instruction (Inpres) No. 2/2007 on rehabilitation of the ex-Mega Rice Project in Central Kalimantan;
- Minister of Forestry Regulation (Permenhut) No. 14 /2004 on Afforestation and Reforestation (AR) CDM
- Minister of Forestry Regulation (Permenhut) No. 1/2004 on social forestry.
- Minister of Forestry Regulation (Permenhut) No 68/Menhut-II/2008 on the implementation of REDD demonstration activities.
- Minister of Forestry Regulation (Permenhut) No. 36/Menhut-II/2009 on procedure for getting permit for implementing carbon sequestration, carbon conservation on production forest and protection forest.
- Minister of Forestry Regulation (Permenhut) No 30/Menhut-II/2009 on procedure for implementing REDD activities
- Economic and Finance Ministerial Decree (Kepmen Ekuin) No. 14/2001 on Integrated Water Resources;

- Forest Ministerial Decree (Kepmenhut) No. 159/2004 on restoration of degraded ecosystem in productive forest areas;

### 5.3.2 Emissions Projections and Mitigation Measures

The forestry sector developed emissions and carbon sequestration projections based on historical data. Under the BAU scenario, the future deforestation rate is assumed to be constant at a rate of 1.1 million ha per year (see Figure 5.9). Figure 5.9 shows that the rate of deforestation from 2000-2006 has decreased compared to the previous period. In the analysis, this trend was not taken into account because of the potentially increasing threat of the conversion of forest area to non-forest area due to the formation of new districts in the future. Similarly, the future capacity to establish forest depends on rehabilitation programs (reforestation and afforestation), community forestry, community forest plantation, etc. and are assumed to be the same as historical rate. From 2000 and 2006 satellite data, it was suggested that the rate of forest establishment between 1996 and 2006 was about 198 thousand ha per year (MoF, 2009).

Using the above assumption, under the BAU scenario, the emissions from biomass removal due to deforestation was projected to be constant at a rate of about 0.898 Gt CO<sub>2</sub> per year. The rate of sequestration is assumed to increase from 0.505 Gt CO<sub>2</sub> per year in 2005 to 0.753 Gt CO<sub>2</sub> in 2020. Sequestration of carbon occurs as a result of regeneration of secondary forests, land rehabilitation (i.e., afforestation and reforestation), and regrowth of woody vegetation (i.e., perennial crops and shrubs). Based on various studies, the mean annual growth rate of secondary forest is assumed to be approximately 5.32 tCO<sub>2</sub>/ha, forest plantation is 36.7 t CO<sub>2</sub>/ha, and other perennial crops/shrubs is 13.5 tCO<sub>2</sub>/ha.

Under the BAU scenario, this sector will continue as net emitter. The Government of Indonesia, as part of its non-binding emissions reduction target, will increase the rate of planting from 198 thousand hectare per year to 500 thousand ha per year, which is referred to as mitigation scenario "1." In its Strategic Plan, the forestry sector set up a progressive target with a rate of planting of between 1.6 and 2.2 Mha per year (Table 5.11), which is called as Mitigation Scenario 2. With this rate of planting this sector will become net sinker after 2015. After 2015, the rate of carbon sequestration is projected to be greater than the rate of emissions (Figure 5.9).

Table 5.11 Rate of planting in the Strategic Plan of the Ministry of Forestry (MoF, 2009)

| Year           | Forest Plantation | Community forest | Forest restoration        | Watershed rehabilitation  | Partnership forest | Total |
|----------------|-------------------|------------------|---------------------------|---------------------------|--------------------|-------|
| 2010           | 450               | 500              | 300                       | 300                       | 50                 | 1600  |
| 2011           | 550               | 500              | 350                       | 300                       | 50                 | 1750  |
| 2012           | 500               | 500              | 450                       | 300                       | 50                 | 1800  |
| 2013           | 600               | 500              | 650                       | 350                       | 50                 | 2150  |
| 2014           | 550               | 500              | 750                       | 350                       | 50                 | 2200  |
| 2015           | 450               | 500              | 300                       | 300                       | 50                 | 1600  |
| 2016           | 550               | 500              | 350                       | 300                       | 50                 | 1750  |
| 2017           | 500               | 500              | 450                       | 300                       | 50                 | 1800  |
| 2018           | 600               | 500              | 650                       | 350                       | 50                 | 2150  |
| 2019           | 550               | 500              | 750                       | 350                       | 50                 | 2200  |
| 2020           | 500               | 500              | 750                       | 350                       | 50                 | 2150  |
| Total          | 5800              | 5500             | 5750                      | 3550                      | 550                | 21150 |
| Source of Fund | Private           | Government       | Private and International | Private and International | CSR and Community  |       |




Figure 5.9 Rate of carbon sequestration and emissions under the BAU scenario and the two mitigation scenarios (Ministry of Forestry)

The progressive target in the Strategic Plan will be hard to meet without international support (see Table 5.12). Government of Indonesia has identified potential funding sources that can be used to achieve such targets, namely the REDD Fund, private investment, grants through bilateral and multilateral channels such as Overseas Development Assistance (ODA), Adaptation

Fund under the Kyoto Protocol, Global Forest Fund (GFF), and Debt for Nature Swaps (DNS). Indonesia applied for debt reduction under the Tropical Forest Conservation Act of the USA and used the fund for forest conservation activities.

In addition, the previously (albeit briefly) mentioned UN Collaborative Programme on Reducing Emissions from Deforestation and Forest Degradation (REDD) also has high potential for providing additional support to Indonesia and several countries have made already made commitments [to support Indonesia]. Both REDD and the Forest Carbon Partnership Facility (FCPF) have made commitments to support Indonesia in developing an effective and accountable framework for REDD, together with Australia, the United Kingdom, Germany, Japan and Korea. Table 5.12 presents some of the international support for Indonesia related to implementing forestry mitigation projects.

**Table 5. 12 Program support from international agencies for forestry mitigation projects in Indonesia**

|                  |  |
|------------------|--|
| <b>Title</b>     | <b>Cooperation to Support Forest Governance and Multi-stakeholder Forestry Programme</b>   |
| <b>Duration</b>  | 2007-2010  |
| <b>Objective</b> | <ol style="list-style-type: none"> <li>1. Support governance reforms to reduce and eventually eliminate illegal logging and its associated timber trade, with a particular focus on support for negotiation and implementation of the EU-Gol FLEGT VPA and other international arrangements;</li> <li>2. Through a multi-stakeholder approach, help build the capacity of central and local government and civil society, support partnerships between government and civil society, promote policy analysis and development and support poverty reduction through more equitable and sustainable management of natural resources, with a particular focus on the rights and opportunities through community forestry for disadvantaged and women's groups;</li> <li>3. Explore the opportunities for governance reforms that are necessary for REDD.</li> </ol> |
| <b>Title</b>     | <b>Financial Cooperation for Climate Change and REDD Issue;<br/>Project title:<br/>Forestry-Climate Change Project in Central and East Kalimantan (FCCP)</b>   |
| <b>Duration</b>  | 7 years, started 2009  |
| <b>Objective</b> | To support Indonesia with the reduction of GHG emissions from deforestation and forest degradation. Specific objectives: <ol style="list-style-type: none"> <li>1. To support policy priorities of the Ministry of Forestry in REDD; and</li> <li>2. To support the implementation of mechanisms related to avoiding deforestation by development of pilot projects in Indonesia with the involvement of local communities in sustainable forest management.</li> </ol>  |
| <b>Title</b>     | <b>Technical Cooperation (TC)<br/>Supporting implementation of the Ministry of Forestry's Strategic Plan</b>   |
| <b>Duration</b>  | 3 years, started mid-2008  |
| <b>Objective</b> | To implement the Strategic Plan which is integrated and synergized with other sector planning – specifically,, provincial and districts programs   |

|                  |   |
|------------------|---|
| <b>Title</b>     | <b>Technical Cooperation Forestry Program in Implementing The Heart of Borneo (HoB) Initiative (Malaysia, Indonesia and Brunei Darussalam)</b>  |
| <b>Duration</b>  | 3 years, started mid-2008   |
| <b>Objective</b> | <ol style="list-style-type: none"> <li>1. To establish trilateral, national and local (states, provincial and district) institutional arrangement to support the implementation of the HoB Initiative;</li> <li>2. To develop mechanisms, including inter-alia action plans at all levels, on the implementation of HoB programs; and</li> <li>3. To strengthen the capacity of stakeholders related to the implementation of HoB programs.</li> </ol>  |
| <b>Title</b>     | <b>Kalimantan Forests and Climate Partnership</b>   |
| <b>Duration</b>  | 2007-2012   |
| <b>Objective</b> | The goal of the program is to support the GOI's efforts to reduce GHG emissions associated with deforestation in Indonesia, through actions to reduce rates of deforestation, support reforestation and promote sustainable forest management and deliver improvements to rural livelihoods and environmental benefits.   |
| <b>Title</b>     | <b>The United Nations Collaborative Programme on Reducing Emissions from Deforestation and Forest Degradation in Developing Countries (UN-REDD Programme)</b>   |
| <b>Duration</b>  | 2009-2011   |
| <b>Objective</b> | <p>A Food and Agriculture Organization (FAO), UN Development Programme (UNDP) and UN Environment Programme (UNEP) collaboration, UN-REDD in Indonesia aims to:</p> <ol style="list-style-type: none"> <li>1. Strengthen multi-stakeholder participation and consensus at the national level;</li> <li>2. Successfully demonstrate the establishment of the Reference Emission Level (REL), Monitoring, Reporting and Verification (MRV) and fair payment systems based on the national REDD architecture; and</li> <li>3. Build capacity to implement REDD at decentralized levels through demonstration activities and other support.</li> </ol> |
| <b>Title</b>     | <b>Korea-Indonesia Joint Program on Adaptation and Mitigation of Climate Change in Forestry through A/R CDM and other Related Mechanisms</b>  |
| <b>Duration</b>  | 2008-2012   |
| <b>Objective</b> | <ol style="list-style-type: none"> <li>1. Analyze REDD applications and acquire a framework of carbon credits by preventing forest conversion as a post-2012 preventive measure;</li> <li>2. Implement capacity building programs, including expert exchange and training courses; and</li> <li>3. Acquire cost-effective potential A/R CDM sites and establish a foundation for carbon credits in preparation of post-2012 emissions commitments.</li> </ol>   |

Source: Ministry of Forestry, Republic of Indonesia

### 5.3.3 Mitigation Costs

Studies on forestry mitigation in Indonesia have been conducted since early 1990s (e.g., Directorate for Nature and Management [DNM] Norway and Ministry of State of Environment [MSE] Indonesia, 1993; Adi *et al.*, 1999; Boer *et al.*, 1999; Boer, 2000; Fuad, 2000; Boer, 2001; Boer *et al.*, 2001). In terms of cost for forestry mitigation, the results of the studies varied considerably (Table 5.13). A recent study conducted by DNPI (2009) suggested that the average cost for reducing emissions in the forestry sector was approximately 9 USD/tCO<sub>2</sub> or about 2.5 USD/tC, while the MoF (2007) estimate was between 10 and 23 USD/tC.

Table 5.13 Mitigation potential and cost for forestry sector

| Mitigation                                | Mitigation potential (tC/ha) | Life cycle cost (USD/tC) |
|---|------------------------------|--------------------------|
| <b>Conservation and forest management</b> |                              |                          |
| Forest Protection                         | 55-220                       | 1.18                     |
| Reduced Impact Logging                    | 49                           | 0.07                     |
| Enrichment planting                       | 70                           | 0.25                     |
| <b>Sink enhancement</b>                   |                              |                          |
| Reforestation without rotation/harvesting |                              |                          |
| - Fast growing species                    | 40-101                       | 0.85-22.5                |
| - Slow growing species                    | 94-336                       | 0.48-2.34                |
| Reforestation with rotation/harvesting    |                              |                          |
| - Short rotation                          | 56-122                       | 3.87-33.20               |
| - Long rotation                           | 134-334                      | 1.04-5.70                |
| Agroforestry                              | 94                           | 4.44                     |
| <b>Fossil fuel substitution</b>           |                              |                          |
| Bioelectricity                            | 50-185                       | 20.81                    |

Source: Based on various sources, including Adi et al., 1999; Boer et al., 1999; Fuad, 2000; Boer, 2001; and MoF, 2007.

## 5.4 Peatland

### 5.4.1 Policy Framework

To support the implementation of programs related to reduce emissions from land and forest fires, forest conversion and peat land from agriculture plantation, the Government of Indonesia has issued a number legal instruments. Relevant legal instruments designed to control fire include the following:

- Ministry of Agriculture Regulation No. 14/Permentan/PL.110/2/2009 on the Guidance for the Utilization of Peat Land for Palm Oil;
- PP No. 4/2001, forbids all forest and land fires;
- Ministry of Forestry Decree Kepmen No. 260/Kep-II/1995 on Guidelines for Prevention and Control of Forest Fire, supplemented with the implementation guidelines;
- Director General of Forest Protection and Nature Conservation (PHPA) Decree No. 243/Kpts/DJ.VI/1995 on Technical Guidelines for Forest Fire Prevention and Control in concession areas and other land use;
- Director General of Estate Crops Decree No. 38/KB.110/DJ.BUN/05.95 on Technical Guidelines for Land Clearance without Burning to Develop Plantations;



- While strict legal penalties for persons causing fire are stipulated in UU No. 41/1999, article 78 clauses 3, 4 and 11, are as follows :
  - o Intentionally setting fire to forest: Prison sentence maximum of 15 years and a maximum fine of 5 billion Indonesian Rupee (IDR)
  - o Negligence leading to forest fire: Prison sentence maximum of 5 years and a maximum fine of 1.5 billion IDR
  - o Dumping of materials which can cause forest fire: Prison sentence maximum of 3 years and a maximum fine of 1 billion IDR
- Government Regulation No. 28/1985 on Forest Protection:
  - o Article 10, point 1: burning the forest on purpose will be fined with a maximum of 10 years in prison or 100 million IDR
  - o Article 18, point 3b: burning of forest as a result of carelessness will be fined a maximum of 1 year in prison or 10 million IDR

### 5.4.2 Emissions Projections

Under the BAU scenario, it was assumed that all peat land under convertible production forest (HPK) irrespective of depth will be released and used for non-forestry activities (APL). Peat land in HPK will be released for APL within the next 15 years and used for agriculture activities (BAPPENAS, 2010). The emissions from peat fire are assumed to be the same as the average rate of emissions from 2000-2006 estimated by van der Werf *et al.* (2008). Based on this assumption, the rate of emissions from peat land in 2020 might reach 1.4 GtCO<sub>2</sub>, about approximately four times the 2000 peat emissions.

## 5.5 Agriculture Sector

### 5.5.1 Policy Framework

There is no specific policy in the agriculture sector for climate change mitigation. However, a number of mitigation programs are already in place. The key mitigation programs include (MoA, 2007; Las *et al.*, 2008; Bappenas, 2010):

1. Implementation of no-burning technology for land clearing and land preparation, in food crop, horticulture, and plantation sub-sectors;
2. Development of a Fire Early Warning System to reduce the risk of fire during extreme drought years, particularly in peat areas;
3. Introduction of low methane emitting rice varieties. Currently, there are four varieties used that have low methane emissions – Ciherang, Cisantana, Tukad Belian and Way Apo Buru.
4. Introduction of low methane emitting technologies, such as the use of agriculture waste for bio-energy and composting;
5. Development and application of organic fertilizer and bio-pesticides

6. Introduction of biogas technology and improved feed to reduce methane emissions from the livestock. A number of related activities that have been implemented are the 'Batamas' program, mixed-farming and mini animal feed industries;
7. Optimization of existing agricultural lands by increasing productivity and cropping index through technology innovation support as new improved varieties
8. Expansion of agriculture areas in unproductive land such as grassland and abandoned land particularly to mineral soils either in upland or wet lands
9. Rehabilitation, reclamation and revitalization of abandoned peat lands, and management of peat land for sustainable agriculture,
10. Research and development on low emission technology and MRV in agriculture sector

In the framework of green economy development program, Indonesia is developing carbon efficient farming (ICEF) by organizing farmers in villages or sub-districts. ICEF is an integrated of crop-livestock with low emission, energy efficient and high economic added values. ICEF optimally utilizes all the carbon from crop production and waste or by products from both the crop and livestock.

### 5.5.2 Emissions Projections and Mitigation Measures

The main sources of CH<sub>4</sub> emissions from the agriculture sector are rice fields and livestock. Projections of emissions under these two sub-sectors were developed using a number of assumptions to model the impacts of different policy options. The assumptions were developed by considering historical conditions and a trend of population growth. For rice, two baseline scenarios were developed:

1. **BAU1.** This baseline adopted the assumption that all irrigated rice fields are continuously flooded and applied inorganic fertilizers to Cisadane rice variety. The area of the irrigated rice in Java will not change due to the adoption of regulations related to sustainable agriculture land; outside Java, the land area devoted to irrigated rice will increase at a rate of 50 thousand ha per year. The cropping index is assumed to be 2.4 if the source of irrigation came from a reservoir and 1.4 if source of irrigation is not from reservoir.
2. **BAU2:** Similar to BAU1, but the rice paddy area in Java is assumed to be converted to non-rice paddy area at a rate of about 50 thousand ha per year; outside Java, rice paddy area will increase at a rate of about 150 thousand ha per year.


To reduce emissions from the rate under BAU, six mitigation scenarios were proposed, as shown in Table 5.14 (S1-S6). In the mitigation scenarios, the rice variety used was changed from Cisadane to IR64. By 2030, the total rice paddy area that applied mitigation technologies is less than 30% - specifically, 21% for S1, 5% for S2, 15% for S3, 14% for S4, 6% for S5 and 1.6% for S6. Using these assumptions, it was found that total methane emissions in 2030 under BAU1 and BAU2 would be about 38,804 and 42,434 Gg CO<sub>2</sub>e, respectively. The most effective mitigation scenario would be the adoption of less methane-emitting varieties (Figure 5.10). Adoption of

new less methane emitting varieties would reduce the rate of methane emissions from BAU by approximately 20%.

**Table 5.14 Mitigation scenarios for reducing methane emissions from paddy rice**

| Scenario | Mitigation Technology  | Potential applicability (%) | Rate of the technology adoption (Percent of potential applicability area) |      |      |      |      |      |
|----------|--|-----------------------------|---|------|------|------|------|------|
|          |  |                             | 2005*   | 2010 | 2015 | 2020 | 2025 | 2030 |
| S0       | Flooded with inorganic fertilizer (according to Kepmentan)             | 100                         | 100   | 100  | 100  | 100  | 100  | 100  |
| S1       | Intermittent irrigation (including SRI, PTT)                           | 70                          | 3   | 5    | 10   | 20   | 25   | 30   |
| S2       | Fertilizer supplement (ZA and urea brisket)                            | 100                         | 0   | 0    | 1    | 2    | 3    | 5    |
| S3       | Adoption of less methane emissions-varieties*                          | 30                          | 20  | 25   | 30   | 40   | 45   | 50   |
| S4       | Intermittent + fertilizer supplement (Combine S1 and S2)               | 70                          | 3   | 5    | 8    | 10   | 15   | 20   |
| S5       | Intermittent + fertilizer supplement + varieties (Combine S1, S2 & S3) | 30                          | 3   | 5    | 8    | 10   | 15   | 20   |
| S6       | S5 + iron material/silica  | 30                          | 0   | 0    | 1    | 2    | 4    | 6    |

Note: Total paddy rice area was assumed to be about 7.8 million ha. Potential applicability area refers to the total area applicable for the mitigation technologies. The remaining areas that do not use the mitigation technologies still use the baseline technologies. Less methane emitting varieties include Ciherang, Cisantana, Tukad Belian and Way Apo Buru (Setyanto *et al.*, 2009)



**Figure 5.10 The projection of methane emissions from rice paddies under the baseline and mitigation scenarios (Setyanto *et al.*, 2009)**

The projection of emissions for livestock was consistent with the population growth projection. The annual population growth rate of the livestock was assumed to follow the historical population growth rate, whereas beef cattle and dairy cattle increase at an annual rate of 5%, broiler and layer 3%, buffalo, sheep, goat, pig and local chicken 2%, and horse and duck 1%. With these growth rates, the domestic demand for these animals will not be fully met, implying that Indonesia may still need to import meat and milk.

Mitigation technologies that have been identified for this sub-sector are the following:


1. **Feeding Quality Improvement.** Currently, beef and dairy cattle feeding practices are less efficient (increasing losses of feed energy in the form of methane and reduced amounts of energy stored in animal product). Utilization of low digestible feedstuff, a higher proportion of forage and low quality pasture (especially in eastern parts of Indonesia) are the main problems. Increasing conversions of feed energy to animal products will not only decrease methane emissions, but also benefit the farmer. Feed quality improvement programs of beef and dairy cattle will reduce up to 20% of methane emissions. Some possible activities for reducing methane emissions through improved feed quality are:
  - a. Introduction of high quality local feed sources such as legume and high quality grass, as well as utilization of agriculture byproducts;
  - b. Improvement of low quality byproducts using physical, chemical or biological pre-treatment;
  - c. Increasing concentrate to forage ratio; and
  - d. Increasing animal productivity by intensity of production (feeding as required). The equal amount of animal product will be produced by less animal number.
2. **Supplementation.** Supplementation of minerals in the form of feed supplement blocks (FBS) has been tested on a limited scale by smallholder farmers. One study reported that mineral supplements increased animal productivity, reducing methane formation by up to 15% (Suryahadi *et al.*, 2007). Mineral supplementation programs have a lot of advantages due to their simplicity of application and high success rate (less risk of failure).
3. **Long-term Breeding Program.** Improvement of animal breeding could mitigate up to 10% of methane emissions from livestock. This improvement could be achieved through appropriate long-term breeding programs. The animals will be bred for adaptation to climate, higher productivity and less methanogenic bacteria/methane production. A potential breeding program would need high investment and would yield long-term return rates, thus likely requiring government support.
4. **Bio-energy.** Utilization of animal sewage as bio-energy is a clear option because of its relevance to national government programs. Currently, the Indonesian government is trying to incentivize the deployment of renewable and environmentally-friendly energy resources. Bio-energy from animal sewage could reduce methane emissions up to 80%

compared to untreated sewage. Furthermore, a co-product of the digester used is an organic fertilizer, which could be used on crops and plants, would have high economic value. This program has disadvantages, however, because of the significant investment needs related to digester technology deployment.

With the adoption of these options (Table 5.15) the rate of methane emissions from this sub-sector would reduce slightly (Figure 5.11).

**Table 5.15 Projected mitigation technology options adopted for livestock**

| No. | Mitigation Options            | Animal population mitigation applied to*                     |
|-----|-------------------------------|--|
| 1.  | Supplementation               | 1% of beef cattle and 3% of dairy cattle population per year |
| 2.  | Improvement feeding practices | 2% of beef cattle and 5% of dairy cattle population per year |
| 3.  | Manure management/biogas      | 1% of beef cattle and 1% of dairy cattle population per year |
| 4.  | Long-term breeding program    | 1% of dairy cattle population per year                       |
| 5.  | Tree legume introduction      | 1% of dairy cattle of population per year                    |



**Figure 5.11 Projected methane emissions from livestock (Suryahadi *et al.*, 2009)**

## 5.6 Waste

Municipal solid waste (MSW) landfills are the most significant of GHG emission sources within this sector. Other major sources are industrial solid waste, domestic liquid waste and industrial liquid waste. The national generation of MSW was approximately 47.8 million tons in 2000 and increased to 48.7 million tons in 2005 (estimated from population data and waste generation factors given by Profil Nasional Program Bangun Praja from Ministry of Environment, 2004]. In urban areas of big cities, almost 60% of waste is brought to a solid waste disposal site (SWDS), while in small cities and rural areas, only 30% of waste is brought to a SWDS (Indonesian Statistical Data on Environment, BPS, 2000-2007). The major components of the solid waste brought to SWDS are organic compounds because other types of waste (i.e., plastics, metal, etc.) are generally recycled for reutilization. The SWDSs in most big cities of Indonesia are considered to be unmanaged SWDS (an open dumping system). Currently, incinerators for MSW are generally not used in Indonesia. Although several statistical data indicate that incineration is already used for eliminating MSW, the so-called 'incinerator' is actually an 'open burning' system.

The main source of industrial solid waste is from agro industries, particularly empty fruit bunches (EFBs) generated from palm oil mills. The total amount of waste in 2000 was 6.3 million tons and increased to 12.5 million tons in 2005. The amount of this waste was estimated from waste generation factor (CDM Project Surveys by EcoSecurities, 2007) and palm oil production rates (Central Bureau of Statistic Indonesia, 2007). The current practice for handling EFBs is open dumping (unmanaged) within the vicinity of the palm oil mills.

Domestic wastewater, represented by biological oxygen demand (BOD), was equal to 3 million tons BOD in 2000 and it increased to 3.2 million tons BOD in 2005. Industrial wastewater produced, expressed in terms of TOW was equal to about 23.6 million tons COD in 2000 and 23.7 million tons COD in 2005. The major contributor to industrial wastewater COD is pulp and paper industries. Production of this waste will increase in the future.

To reduce emissions from solid waste, the Government of Indonesia enacted Municipal Solid Management Law No. 18/2008. According to the Law, open dumping practices will be prohibited in the year 2013. It is expected that the Law will encourage the development of a more managed waste handling system (i.e., a sanitary landfill equipped with gas flaring or utilization systems). The GOI expects that after 2020, around 80% of the domestic liquid waste will be handled by sewerage systems.


To estimate future emissions, three scenarios have been established: (a) the BAU scenario assumes that emissions growth will follow historical trends, without any efforts or programs related to climate change mitigation; (b) emissions reductions are to be achieved through implementation of Municipal Solid Management Law No. 18/2008; and (c) further emissions reductions from the waste sector are to be achieved through the implementation of Municipal Solid Management Law No. 18/2008 combined with international funding (CDM scheme or

other schemes). The projection of the BAU and the other two alternative scenarios is presented in Table 5.16.

**Table 5.16 Projection of GHG emissions from waste sector under BAU and alternative scenarios (Dewi *et al.*, 2009)**

| Scenario Mitigation          | 2010           |                |                | 2020           |                |                | 2025           |                |                |
|------------------------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|
|                              | BAU            | S1             | S2             | BAU            | S1             | S2             | BAU            | S1             | S2             |
| Total MSW                    | 19,691         | 19,199         | 18,706         | 22,198         | 19,423         | 17,204         | 23,562         | 20,028         | 17,672         |
| Unmanaged Dumpsite (CPO)     | 11,289         | 10,725         | 10,443         | 13,269         | 10,615         | 9,288          | 14,385         | 10,069         | 8,631          |
| Domestic WWT and Discharge   | 13,568         | 12,890         | 12,551         | 15,287         | 12,230         | 10,701         | 16,227         | 11,359         | 9,736          |
| Industrial WWT and Discharge | 149,818        | 142,327        | 138,582        | 199,477        | 159,581        | 134,647        | 224,411        | 157,088        | 123,426        |
| <b>Total Emission</b>        | <b>194,367</b> | <b>185,141</b> | <b>180,282</b> | <b>250,231</b> | <b>201,849</b> | <b>171,839</b> | <b>278,585</b> | <b>198,544</b> | <b>159,465</b> |

Based on the emissions projection analysis, it was found that by 2020, the rate of net emissions under mitigation scenario 1 reached 31% of the BAU emissions and that under mitigation scenario 2, it could reduce up to 48% (Figure 5.12). Net emissions under BAU in 2020 are expected to be approximately 2.95 Gt CO<sub>2</sub> – more than double the emissions in year 2000.



**Figure 5.12 Projection of emissions of all sectors under BAU and mitigation scenarios (MoE, 2009). Note: Contribution of transportation sector to the total energy emission is approximately 20% (based on emissions data from 2000-2006).**






## CHAPTER VI

# OTHER INFORMATION

Other information considered relevant to the achievement of the objectives of the Convention includes (i) technology transfer, (ii) Research, (iii) systematic observations, (iv) information on education, training, public awareness and capacity-building and (v) efforts to promote information sharing.

### 6.1 Transfer of Technology

Transfer of technologies related to climate change occurs through the implementation of CDM projects. At present, Designated National Authorities for CDM, KOMNAS MPB, has approved about 104 CDM projects. Of this, about 24 projects have been registered at the Executive Board ([www.dna-cdm.menlh.go.id/id/database](http://www.dna-cdm.menlh.go.id/id/database)). Most of the proponents of the CDM projects are private companies. However, NGOs play important roles in the implementation of adaptation and mitigation activities with local communities. Based on data from the SGP-GEF UNDP program, there are about 18 NGOs working on climate change mitigation projects. This NGO work is predominantly centered in Java (Figure 6.1) and the focus of their activities is mostly on the energy and waste sectors.



**Figure 6.1** Number of NGOs working in climate change mitigation activities on energy and waste sectors (Boer *et al.*, 2008)

Several NGOs, especially those who have technical competencies, act also as CDM or voluntary carbon project developers. NGOs or some international NGOs who have technical expertise on digester construction, in collaboration Agency for Rural Technology Development (LPTP, Lembaga Pengembangan Teknologi Pedesaan), have provided technical assistance for designing the necessary infrastructure for community. NGOs receive funds as grants from a number of development agencies. Often these funds are distributed through development aid programs and INGOs (International NGOs).

## 6.2 Research

Research activities related to climate are implemented by many research agencies and universities. Some international organizations based in Indonesia recently have also focused their research activities on climate change. Based on a survey conducted in 2007 of nine organizations working on climate related activities (Boer and Kartikasari, 2007), it was found that research activities conducted by the agencies was quite variable. Research activities on climate applications were mainly conducted by the Research Agency on Agro-climatology and Hydrology (BALITKILIMAT) and Bogor Agriculture University (IPB), climate early warning systems by LAPAN (National Agency for Aviation and Space) and BMKG (National Agency for Meteorology, Climatology and Geophysics), and weather modification by the Agency for Technology Assessment and Application (BPPT). Basic research on weather and climatology modeling were conducted by LAPAN, Bandung Institute of Technology (ITB), BMKG and IPB. Most NGOs have no research activities. However at present, many NGOs also conduct climate research activities and mostly the research were conducted in collaboration with universities and research agencies. At present, research activities on climate change for agriculture sector is coordinated by Indonesian Agency for Agriculture Research and Development through Indonesia Centre for Land Resources Research and Development that have four research agencies namely Agro-climate and Hydrology Research Institute (BALITKILIMAT), Agriculture Environment Research Institute (BALINGTAN), Soil Research Institute (BALITTANAH), and Swampy Land Research Institute (BALITRA).

The number of research scientists in the surveyed research agencies and the universities that hold advanced degree is about 90 people (33 PhD and 57 Master). About 29% have expertise in the area of applied meteorology or applied climatology, 23% meteorology, 16% climatology or climate modeling, 10% atmospheric chemistry of air pollution, 10% hydrology or hydrometeorology, the remaining are atmospheric physics/radars atmosphere, GIS/Remote sensing, microclimatology, geography, astronomy and meteorology instrumentation. The scientists with expertise in applied climatology concentrates in IPB and BALITKILIMAT, while those with expertise in climatology/meteorology concentrate in ITB, BPPT, LAPAN and BMKG (Boer and Kartikasari, 2007). However, recently some of the agencies have expanded their research activities to climate application. For example, BMKG has established a new agency called the Research Centre on Climate Change and Air Quality. Some universities have also established new climate research centers.

To accelerate the development of mitigation and adaptation technologies, in 2008 the Government of Indonesia under the coordination of the Ministry of Agriculture has established the Research Consortium on Climate Variability and Climate Change (especially for Agriculture Sector). This consortium plays role in (i) coordinating each sector's adaptation research resources to more effectively support climate change decision-making, including the brokering of research partnerships and providing a vehicle to commission new integrated research; (ii) Synergizing research programs and activities on climate change and establishing a road map in mitigating and adapting to climate change, (iii) building the capacity of the Indonesia research community to generate information relevant to decision-makers; (iv) establishing an interface between researchers and decision-makers; (v) promoting coordinated programmes of work on impacts and adaptation across Indonesia, working in collaboration with stakeholders and other researchers in national, regional and sectoral contexts, (vi) synthesizing policy and developing sectoral roadmap for addressing climate change for 2010-2050, (vii) delivering the information to support climate change adaptation decision-making at the national, regional and local levels through coordination, integration, synthesis and communication of research.

The consortium conducts research and development activities on inventories, mitigation and adaptation, which include the following: (i) development of GHG emission and removal factors and methods for improving activity data, (ii) identification and development of mitigation technologies potential for carbon trading as well as the development of baseline and monitoring methodologies for the technologies as required by the CDM Executive Board, and (iii) conducting policy research on maintaining food crop self sufficiency under changing demography, socio-economic, land use and climate (e.g. mapping vulnerable areas to climate change, strategies for coping with the climate changes etc). The Ministry of Agriculture for the fiscal year of 2008/2009 and 2009/2010 have allocated research funds from the National Budget for the consortium by about 0.4 Million USD. The Ministry of Research and Technology through BPPT for the fiscal year of 2008/09 has also allocated about 1.5 Million USD to implement research activities on climate change. There are many climate change research activities implemented by other research agencies using national budget. However the data and information on this is not well documented.

### 6.3 Systematic Observations

Systematic observation of the climate in Indonesia is coordinated by the National Agency for Meteorology, Climatology and Geophysics (BMKG). Rainfall Observation Networks exist since Dutch Colonial Era. After independence, more rainfall stations were installed, however, the efforts are still far from enough to have a representative number of stations across the country. The total number of rainfall observation network (rain gauges) is about 5,000 gauges. Most of the gauges are located in Java and then followed in Sumatra, Sulawesi, Maluku, Kalimantan and Papua (Table 6.1). The ideal number of rain gauges to meet the WMO standard for Indonesia is about 20,000 gauges (Sribimawati, 1999). In addition to the rain gauges stations, BMKG also installed a number of automatic weather stations (AWS) and Radar in a number of places


particularly in areas prone to weather/climate hazards. In collaboration with WMO, BMKG has also established one Global Atmospheric Watch (GAW) in Bukittinggi, West Sumatra. With the presence of the GAW, Indonesia becomes part of the global atmospheric observation under World Weather Watch.

**Table 6.1 Rain gauges density in Indonesia**

| No. | Island       | Rain gauge density per 100 km <sup>2</sup> |
|-----|--------------|--|
| 1   | Java         | 11.6                                       |
| 2   | Sumatra      | 0.87                                       |
| 3   | Sulawesi     | 0.67                                       |
| 4   | Maluku       | 0.46                                       |
| 5   | Kalimantan   | 0.18                                       |
| 6   | Papua        | 0.05                                       |
| 7   | Other island | 0.67-1.14                                  |


Source: Sribimawati et al. (1999)

Indonesia has installed a number of instruments to monitor sea level (Figure 6.2). The existing Indonesia Sea Level Monitoring Network consists of 65 operational stations. More stations will be installed through an ongoing program called the establishment of Indonesia Tsunami Early Warning System (IndTEWS). The network will consist of 120 stations of which 80 stations will use real time data transmission and at least two quality sea level recordings. Solar cell power supplies for each station have been installed to ensure an availability of backup power for continuous measurement.



**Figure 6.2 Existing operational Sea Level Monitoring Stations in Indonesia (MoE, 2008)**

The National Network of Agroclimatic Stations has also been installed in a number of agriculture research stations. The network is maintained by the National Agency for Agriculture Research and Development. There are about 59 agroclimatic stations but stations in good condition are around half (Surmaini *et al.*, 2010). The distribution of the agroclimatic stations is presented in Figure 6.3.



**Figure 6.3** Number of agroclimatic stations by province (Surmaini *et al.*, 2010)

In forestry sector, development of systematic forest and land use monitoring systems is also crucial to support the implementation of programs for reducing emission from deforestation and forest degradation. The Government of Indonesia is in the process of establishing Indonesia's National Carbon Accounting System (INCAS), based on Australia's system but tailored to Indonesia's (INCAS) unique circumstances. The INCAS will provide a comprehensive and credible account of Indonesia's land based emissions profile and sinks capacity. It will support Indonesia's reporting requirements under the United Nations Framework Convention on Climate Change (UNFCCC) and the Reduced Emissions from Deforestation and Forest Degradation post-2012 global climate protection regime. The INCAS will contribute significantly to Indonesia's carbon accounting, resulting in positive implications both domestically and internationally. It will allow Indonesia to develop a robust modeling and projections capacity for land based carbon accounting, and therefore robust emissions and removal estimates.

The INCAS will pull together information on deforestation, land use change and land use from Indonesia's forest lands and other lands (primarily agricultural lands) to:

- support Indonesia's position in the international development of policy and guidelines on sinks activity and greenhouse gas emissions and their mitigation from land based systems

- Reduce the scientific uncertainties (particularly about peat) that surround estimates of emissions and removals of both CO<sub>2</sub> and non-CO<sub>2</sub> greenhouse gases from land use change.
- Provide monitoring capabilities for existing emissions and sinks, and scenario development and modeling capabilities that support greenhouse gas mitigation and the sinks development agenda through to 2010 and beyond.
- Provide the scientific and technical basis to international negotiations and promote Indonesia's national interests in international fora.
- Develop a comprehensive GIS that includes digital map-based information such as soil maps, remotely sensed images covering the whole of Indonesia and climate and vegetation data.
- Support Indonesia's negotiations on REDD and provide the necessary inputs required for establishing a credible Reference Emission Level.

In relation to forestry sector, Indonesia is developing Indonesia's Forest Resource Information System (FRIS). FRIS will be a comprehensive and transparent information management system to support effective planning and forest management decision making for forest lands in Indonesia. The FRIS will allow the Ministry of Forestry to monitor forest productivity, yield and growth, harvesting rates, age class, species and forest area among other things. It will also compile critical information on deforestation, land use and land use change within Indonesia's forest lands to support a post-Kyoto climate protection regime that seeks to reduce emissions from deforestation and forest degradation (REDD). Combining this data within a single system will provide a good basis for planning and decisions on sustainable forest management.

A lack of reliable information on forest resources has been identified as a major impediment to sustainable forest management in Indonesia. Such information gaps are hampering efforts to monitor deforestation and forest degradation, sustainably manage forest resources, combat illegal logging, protect biodiversity and reduce carbon emissions from land use change. Improving forest information is crucial in the context of the emerging need for the Government of Indonesia to document carbon stocks in forests and greenhouse gas emissions related to land use change and to participate in carbon markets for avoided deforestation and degradation.


## 6.4 Information on Education, Training, Public Awareness and Capacity Building

Between 2000 and 2008, there were about 91 training, public awareness and capacity building activities related to climate change conducted in Indonesia. This record may be much lower than the actual activities since many of climate change capacity building activities have not been recorded by related agencies. There is no system in place at present to record and monitor the outcome of the activities. However, this data may reflect the level of Annex 1 participation in supporting capacity building programs in Indonesia. The data suggests that most capacity building activities were supported by Japan, Germany, ADB and the World Bank (Figure 6.4). Again, the number of activities does not necessarily reflect the amount of funding being

provided. A number of government agencies, local NGOs and universities also conducted such activities with the national budget. Furthermore, types of capacity building activities conducted before 2006 were dominated by workshops and after 2006 most of activities were in the form of climate change seminars. The number of climate change capacity building activities tended to increase after 2000 (Figure 6.5).

### 6.5 Efforts to Promote Information Sharing

BMKG is the main agency responsible for producing climate information used in various ways for sharing information. Climate forecasts information is disseminated through media and printed materials. These printed materials are disseminated to all provinces and sectors. In 2008, BMKG started to develop a Web-based climate information system. The system was initially developed in four regional offices (Palembang, Semarang, Makasar and Banjarbaru; Figure 6.6). This effort is continuing to be expanded to 34 Meteorological Stations all over Indonesia.



**Figure 6.4 Donor countries/agencies who contribute to the implementation of capacity building activities on climate change (Based on data collected from the Ministry of Environment)**




Figure 6.5 Types of capacity building activities on climate change by year (Based on data collected from the Ministry of Environment)




Figure 6.6 Development of web-based climate information system at four regional offices (Provided by the BMKG)



## CHAPTER VII

# OBSTACLES, SHORTAGES AND NEEDS RELATED TO FUNDING, TECHNOLOGY AND CAPACITY

As a country with high vulnerability to the impacts of climate change, Indonesia must have a response to address climate change both in mitigation and adaptation activities. However, in its position as developing country, Indonesia has several limitations in funding in the frame of climate change anticipation as well as developing and disseminating environmentally friendly technology. Therefore, the problems of technology transfer and funding are two issues that should receive support from international community. Additional important activities, which will ensure success of mitigation and adaptation program, are awareness, education, empowerment and capacity building.

Different from climate change mitigation, climate change adaptation has received limited attention from key-stakeholders in Indonesia and donors. Consultations reveal that various activities are ongoing, but that there is an urgent need to start systematically addressing climate change adaptation in a coordinated manner. Coordination within and between the governments and the donor communities is challenging. Activities related to climate change adaptation are still limited but likely to increase rapidly. Both government and development partners have expressed the need to ensure that addressing climate change adaptation will be effectively coordinated. This is of great importance considering the expected scale of required interventions. It has become clear that the national capacity needs to be substantially strengthened, especially in terms of policy advice, technical assistance, and capacity development.

From national consultation, it was revealed that strengthening national capacity to adapt to increasing future climate will need (UNDP CO, Jakarta, 2007):

1. Strong political consensus and multi-sectoral policy coordination among key-stakeholders to address climate change adaptation in the framework of national development planning needs. Climate change, a cross-sectoral issue, will require long term commitment. Key-ministries, institutions and development partners should have a common platform on the necessity to address climate change adaptation. Full support for the process of developing a national strategy and a longer-term programme is essential. Related to the future implementation of the strategy and programme, donor coordination and dissemination of information should be strengthened.
2. A broadly supported National Strategy for Climate Change Adaptation developed in a transparent and consultative manner.

3. Availability of a longer-term climate change adaptation programme. Considering the complexities, scale and cross-sectoral nature of climate change impacts, the programme should provide an umbrella framework to systematically address climate change adaptation at the national, regional, and local levels. Expertise and experience from all these levels and abroad will be brought in the preparation to ensure a comprehensive and high quality programme. Active involvement of the key-stakeholders, particularly of the relevant ministries, institutions and development partners is needed to ensure the programme is fully supported in its development and implementation.

## 7.1 Climate Change Funding

Due to limited funding capacity through the national budget, the GOI Indonesia will try to create various funding schemes, from domestic sources as well as from bilateral and multilateral sources to support the national planning effort on climate change mitigation and adaptation. The GOI recognizes several funding mechanisms as important vehicles for climate change related activities (see Figure 7.1), as follows:




Figure 7.1 Climate Change Financing Scheme (BAPPENAS, 2009)

1. United Nation Convention for Climate Change (UNFCCC) mechanism: Global Environmental Fund (GEF), Special Climate Change Fund (SCCF), Adaptation fund (Protocol)
2. Official Development Assistance (ODA) mechanism:
  - a. Current scheme:
    - Bilateral grant and multilateral institution (ODA and ODA+)
    - Loan (ODA dan ODA+):
    - Sectoral/project loan – funds activities in a specific sector or project
    - Program loan – support policy reform on climate change and fiscal deficit
  - b. Alternative scheme:
    - Trust Fund International (Climate Investment Fund/CIF - ODA)
    - Indonesian Climate Change Trust Fund Indonesia (Managed by GOI and Donors) – (ODA)
3. Carbon investment: Bilateral, Multilateral, and Unilateral

The GOI will apply the rule and procedures under UNFCCC and ODA financing mechanism for climate change financing schemes. The GOI prioritizes grant utilization to finance climate change sectoral priorities implementation. Loan resources can only be utilized when grant funding is insufficient. However, utilization of loans should be the last alternative for climate change financing. Climate change grant financing must comply with the following guidance:

1. Grant financing focus on sector priorities as mentioned in National Development Planning Response to Climate Change called the Yellow Book
2. Grant financing that originates from UNFCCC, ODA, or other financing mechanism must comply with GOI laws and regulation
3. Grant financing must be recorded into the state budget and must follow the Decree of Director General on State Treasury No. PER-67/PB/2006 on Procedures on Registration and Legalization of Foreign Grant

To efficiently and effectively utilize the foreign financing that has been allocated to address climate change related problem, then strengthening the project planning system, improving the monitoring and evaluation system, and synchronizing the financial and procurement systems between government and donors should continuously be made. Taking these steps will ensure that foreign grants and loans match national development priorities and leads to an enhanced level of accountability.

## 7.2 Technology

Research and technology are needed to adapt and mitigate climate change. Low carbon emission and energy efficient technology is needed immediately to reduce GHG emission in the sectors that consume energy such as power plants, industry, transportation, as well as household and commercial sectors. Real time monitoring and the use of sophisticated fire fighting technology and peat management is required to control emission from forest/peat fire

and peat decomposition. Peat lands store large amounts of carbon stocks that, if released, can greatly contribute to global climate change. Although there are many uncertainties regarding Indonesia's GHG emissions, it is generally agreed that peatland degradation is the main source, together with deforestation. Proper management of peatlands is therefore critical to Indonesia's efforts in combating climate change.

Furthermore, for adaptation need in agriculture sector, technology to develop plant varieties that are resilient to climate change impact is required. Agricultural technology that saves water also should be implemented to anticipate water scarcity during longer dry seasons. To increase the preparedness of various parties in addressing the conditions of extreme climate, Indonesia needs weather prediction technology that can produce more accurate results. With this technology, fisherman and farmers can plan more proactively for their activities, thus reducing the losses that they may suffer because of the extreme climate and weather events (drought, storm etc.). In addition, early warning technology should also be implemented to reduce the general public's exposure to the negative impacts related with extreme climate. International cooperation and transfer of technology should be strengthened to share the benefit of technological knowledge worldwide. Indonesia needs to prioritize the development of low-cost technologies with substantial local content.

Indonesia assigns priority to climate change adaptation programs in the four following areas: (i) coastal areas, (ii) food insecure areas in eastern Indonesia, (iii) small islands, and (iv) Java's northern coast, the nation's key rice production areas (BAPPENAS, 2009). Coastal areas are among the areas most affected by sea-level rise and land subsidence-induced environmental degradation. Both have exposed coastal areas to regular flooding and have significant economic and social impacts. As these communities are at the forefront of adapting to climate change, focused regional approach is warranted. Investing in coastal areas to adapt to sea-level rise is a key to meet the MDGs.

In food insecure areas in eastern Indonesia such as Sulawesi, NTT and NTB, the ENSO signal is very strong (Gianinie *et al.*, 2007; Moron *et al.*, 2009). Strong El-Niño events normally cause the problem of false rains, in which isolated rainfall events around the expected onset date do not signal the sustained onset of the monsoon thus shortening the length of rainy season. This condition leads to farmers experiencing repeated crop failures and rendering it possible only to plant crops once in that year. Crop failure due to drought during the 2006/07 El-Niño for example, increased the reliance of NTT farmers to government aid. In normal years, contribution of government aid to farmers' total income was only 5% but in 2006 El-Niño year, it increased to 60% (Kieft and Soekarjo, 2007). This, combined with already low general human development, has triggered a decline in child health nutrition, a key MDG indicator. Current development approaches employed have proven ineffective, thus calling for efforts to enhance local capacity to adapt to the changing climate.

Java's northern coast is also strongly affected by ENSO. Local action will place an emphasis on the provinces of West and East Java. Increased pressure on these lands due to continued

urban growth and relative limited expansion of agricultural lands outside Java has deepened the strategic importance of the Java “rice basket” to ensure national food security. Increased frequency of ENSO events is leading to increased crop losses in these areas. An integrated approach that is smallholder-driven and that includes technology development and spatial planning is a key.

Sea level rise and changing rainfall patterns in combination with extreme wave and high tide are threatening small islands. These undermine the local economy and ecology and may also lead to the loss of the small islands. Programs in these areas should focus on enhancing general human development while strengthening ecological sustainability. More effort will be put on developing adaptation and mitigation measures for small islands.

To increase the resilience of the sectors and community to current and future climate risk, the development and implementation of a comprehensive communications strategy to increase the capacity in using climate information is needed as well as institutionalizing the use of climate information. The development and promotion of tools for adaptation planning tailored to user’s requirements that include (i) decision-support tools such as methods for assessing the costs and benefits of adaptation strategies, and guides for risk management, (ii) methods for understanding social impacts, and (iii) a national ‘one stop shop’ website where decision-makers and their advisers can access information about climate projections, likely climate change impacts, tools, guides and approaches to adaptation planning will be required.

### **7.3 Awareness, Education, Empowerment and Capacity Building**

Raising the awareness of every citizen on the importance of climate change and integrating climate change issues into national education is essential to ensuring public buy-in and the success of mitigation and adaptation programs. Changing human behaviour in order to respect natural environment, and thereby minimizing human impacts on the environment, must occur at all level of society. Climate change is an effective entry point to raise public awareness about the need to develop in a more responsible and suitable manner.

Human resources will need to be strengthened to ensure the efficient and effective implementation of climate change initiatives. Capacity building and empowerment of all related stakeholders will take place, with emphasis on local government and civil society as they will be the key implementers of change at the regional and local level. Capacity building will include immediate steps as well as longer-term efforts and it will consider the complexity of the many sectors involved in climate change.

Our knowledge on the uncertain future climate and its impact is still limited. In spite of this uncertainty, the ability to generate access and interpret information on climate change and its likely impacts is crucial. Further, it is equally important to develop suitable methods for identifying and assessing potential adaptation strategies. Knowledge and methods will need

to span across disciplines, including climate science, biophysical sciences, engineering, social sciences, economics, and planning. Implementation of pilot projects on addressing climate change is very important to generate best practices on how we can improve our resilience to future climate risk.

---

## REFERENCES

- Achanta, A.N., and R. Kanetkar. 1996. *Impact of Climate Change on Forest Productivity: A Case Study of Kerala, India*. Paper presented at the Asia and Pacific Workshop on Climate Change Vulnerability and Adaptation Assessment, January 15-19, Manila, the Philippines.
- ADB and BAPPENAS. 1999. Planning for the Fire Prevention and Drought Management. Final Report. Asian Development Bank and National Planning Agency, Jakarta.
- Adi, A.C., Boer, R., Gunardi, Boedoyo, S., and Bey. A.: 1999. 'Integrated Analysis of GHG Emissions and Sequestration'. Paper presented at International Symposium on Land Use and Forest Management to Mitigate Climate Change.
- ADPC, 2000. ENSO impact and potential forecast applications in Indonesia. Extreme Climate Events Program. Asian Disaster Preparedness Center, Bangkok, Thailand. 90p.
- ADRC. 2005. Total Disaster Risk Management: Good Practices. Asian Disaster preparedness Centre. Kobe, Japan.
- Agus, F., I. Irawan, H. Suganda, W. Wahyunto, A. Setyanto, and M. Kundarto. 2006. Environmental multifunctionality of Indonesian agriculture. *Journal Paddy Water Environment* 4:181-188.
- Aldrian, E and Djamil, S.D. 2006. Long term rainfall trend of the brantas catchment area, East Java. *Indonesian Journal of Geography* 38:26-40.
- Aldrian, E. 2008. PEAT Carbon, fire and climate Interactions. *Jurnal Hidrosfir Indonesia* 3: 23-32.
- Ardiansyah, M. and R. Boer. 2009. Forestry Sector. In MoE. Technical Report on Vulnerability and Adaptation Assessment to Climate Change for Indonesia's Second National Communication. Ministry of Environment and United Nations Development Programme, Jakarta.
- Balitbangtan. 2007. Prospek dan Arah Pengembangan Komoditas Pertanian: Tinjauan Aspek Sumber daya Lahan. Badan Litbang Pertanian. Jakarta.
- Bappenas and Bakornas PB. 2006. National action plan for disaster reduction 2006-2009. Office of The State Minister for National Development Planning/ National Development Planning Agency and National Coordinating Agency for Disaster Management. Jakarta.

- Bappenas, 2008. National Development Planning: Indonesia Response to Climate Change. National Agency for Planning and Development (Bappenas), Republic of Indonesia
- Bappenas, 2009. National Development Plan: Indonesian Response to climate change (*Yellow Book*). National Agency for Planning and Development, Jakarta.
- Bappenas, 2009a. Reducing carbon emissions from Indonesia's peat land. Interim Report of a Multi-Disciplinary Study. National Planning and Development Agency, Jakarta
- Bappenas, 2010. Sectoral Roadmap
- Bappenas. 2009b. Blueprint for Indonesia Climate Change Trust Fund (ICCTF). National Agency for Planning and Development (Bappenas), Republic of Indonesia.
- BAPPENAS/ADB. 1999. *Causes, Extent, Impact and Costs of 1997/98 Fires and Drought Final Report, Annex 1 and 2, Planning for Fire Prevention and Drought Management Project* (Asian Development Bank TA 2999-INO Fortech, Pusat Pengembangan Agribisnis, Margueles Pöyry, Jakarta, Indonesia.
- Boer, R. 2001. Economic Assessment of Mitigation Options for Enhancing and Maintaining Carbon Sink Capacity in Indonesia. *Mitigation and Adaptation Strategies for Global Change* 6:257-290
- Boer, R. A. Buono., A. Rakhman, and A. Turyanti. 2009a. Historical and Future Change of Indonesian Climate. In MoE. Technical Report on Vulnerability and Adaptation Assessment to Climate Change for Indonesia's Second National Communication. Ministry of Environment and United Nations Development Programme, Jakarta.
- Boer, R. and K. Kartikasari, 2007. Assessment of Capacity and Needs to Address Vulnerabilities, Adaptations and Resilience to Climate Risks in Indonesia. Paper submitted to USER - Unit for Social and Environmental Research Faculty of Social Sciences at Chiang Mai University, Bangkok.
- Boer, R. and Perdinan. 2008. Adaptation to climate variability and climate change: its socio-economic aspect. Paper presented in the *the EEPSEA Conference On Climate Change: Impacts, Adaptation, And Policy In South East Asia With A Focus On Economics, Socio-Economics And Institutional Aspects, 13-15 February 2008, Bali* (<http://www.eepsea.cc-sea.org>)
- Boer, R. and Subbiah, A.R. 2005. Agriculture drought in Indonesia. In V.K. Boken, A.P. Cracknell and R.L. Heathcote (eds). *Monitoring and predicting agriculture drought: A global study*, pp: 330-344. Oxford University Press, New York.



- Boer, R., A. Buono, Sumaryanto, E. Surmaini, A. Rakhman, W. Estiningtyas, K. Kartikasari, and Fitriyani. 2009b. Agriculture Sector. Technical Report on Vulnerability and Adaptation Assessment to Climate Change for Indonesia's Second National Communication. Ministry of Environment and United Nations Development Programme, Jakarta.
- Boer, R., A.R. Subbiah, K. Tamkani, H. Hardjanto., and S. Alimoeso. 2006. Institutionalizing Climate Information Applications: Indonesian Case. In Edited by R. Motha, M.V.K. Sivakumar, and M. Bernardi. Strengthening Operational Agrometeorological Services at the National Level. Proceedings of a Inter-Regional Workshop held 22-26 March 2004 in Manila, Philippines. AGM-9, WMO/TD-No. 1277. Available on line at <http://www.wmo.ch/web/wcp/agm/Publications/html/AGM9.htm>
- Boer, R., and K. Kartikasari. 2007. Assessment of Capacity and Needs to Address Vulnerabilities, Adaptations and Resilience to Climate Risks in Indonesia. Technical Report submitted to USER - Unit for Social and Environmental Research Faculty of Social Sciences at Chiang Mai University. [http://www.sea-user.org/download\\_pubdoc.php?doc=3421](http://www.sea-user.org/download_pubdoc.php?doc=3421)
- Boer, R., *et al.*, 2007. Deteksi perubahan iklim dan dampak sosio-ekonominya. Laporan Proyek Kerjasama BMG dan IPB, Bogor.
- Boer, R., Gintings, A.N. and Bey, A. 1999. Greenhouse Gases Inventory and abatement strategy for forestry and land use change sector. *Journal of Agrometeorology* 13:26-26
- Boer, R., R.G. Dewi, Y. Ismawati, S.D. Anggraini., Y. Ardiansyah, E. Admawaty, Irmaida, and A. Suharto. 2008. Study on local actions in Asia contributing to climate change mitigation and alternative financial mechanism. Project Report Submitted to Institute for Global Environmental Studies. Carbon Environmental Research, Bogor
- Boerema, J. 1938. Rainfall Types in Nederlands Indie. *Verhandelingen* No. 18.
- Boland, G.J. M.S. Melzer, A. Hopkin, V. Higgins, and A. Nassuth. 2004. Climate change and plant diseases in Ontario. *Can. J. Plant Pathol.* 26: 335–350
- Bonaro, O., A Lurette,, C Vidal, J Fargues. 2007. Modelling temperature-dependent bionomics of *Bemisia tabaci* (Q-biotype) *Physiological Entomology*,32: 50-55
- BPS, 2005, "Indonesian Input – Output Data", Jakarta, Indonesia
- BPS, 2007a, "Statistic of Indonesia", Jakarta, Indonesia
- BPS, 2007b, "Stastistik Industri Besar dan Sedang Indonesia (Statistics of Indonesian Large and Medium Industries)", Jakarta, Indonesia

- BPS, 2009, "Proyeksi Kependudukan Indonesia (The Projection of Indonesian Population) 2000 - 2025", Jakarta, Indonesia
- Bryant, D., L. Burke, J. McManus and M. Spalding. 1998. Reefs at Risk: A Map-Based Indicator of Threats to the World's Coral Reefs. Washington, D.C.: World Resources Institute.
- Chen X, Kimoto M, Takahashi M (2005) Changes in ENSO in response to greenhouse warming as simulated by the CCSR/NIES/FRCGC coupled GCM. SOLA 1:149–152
- Collins M (2000a) Understanding uncertainties in the response of ENSO to greenhouse warming. Geophysical Research Letter 27:3509–3513
- Collins M (2000b) The El Niño Southern Oscillation in the second Hadley centre coupled model and its response to greenhouse warming. Journal of Climate 13:1299–1312
- Cubasch U, Meehl GA, Boer GJ, Stouffer RJ, Dix M, Noda A, Senior CA, Raper S, Yap KS (2001) Projections of future climate change. In: Houghton JT., Ding Y, Griggs DJ, Noguer M, van der Linden P, Dai X, Maskell K, Johnson CI (eds) Climate change 2001: the scientific basis. Contribution of working group I to the 3rd assessment report of the intergovernmental panel on climate change, Cambridge University Press, Cambridge, pp 525–582
- Depsos, 2009. Tahun 2009 angka pengangguran membengkak. Departement Sosial Republik Indonesia. <http://www.depsos.go.id>
- Dewi, R.G., M.H. Nasution, S.D. Anggraeni, Y. Ismawati, U. Solihin, A. Syafe'i, and U.S.A Kamil. 2009. GHG mitigation analysis for waste sector. In MoE. Projection of GHG Emission and Mitigation Analysis for Indonesia's Second National Communication. Ministry of Environment and United Nations Development Programme, Jakarta.
- Dewi, R.G., U. Siagian, M. H. Nasution, P. Lestari, Farida Z., M. Saleh, A. Wahyudi and Prasetyo. 2009. GHG mitigation analysis for energy, industry and transportation sector. In MoE. Projection of GHG Emission and Mitigation Analysis for Indonesia's Second National Communication. Ministry of Environment and United Nations Development Programme, Jakarta.
- Ditjenbun (Direktorat Jenderal Perkebunan; Directorate General of Plantation). 2007. Road Map Kelapa Sawit. Ditjenbun, Jakarta, Indonesia.
- DNM Norway and MSE Indonesia. 1993. Climate change and forestry Indonesia: Eco-strategies for terrestrial CO<sub>2</sub>-fixation. Directorate for Nature and Management-Norway and Ministry of State for Environment-Indonesia. 117p.

- DNPI, 2009. *Indonesian GHG Abatement Cost Curve* with McKinsey. National Council for Climate Change, Jakarta.
- Duncan, B.N., I. Bey, M. Chin, L. J. Mickley, T. D. Fairlie, R. V. Martin, and H. Matsueda. 2003. Indonesian wildfires of 1997: Impact on tropospheric chemistry. *Journal of Geophysical Research* 108(D15):4458
- Emanuel, K. 2005. Increasing destructiveness of tropical cyclones over the past 30 years. *Nature* 36: 686-688
- Fuad, E.D: 2000. 'Analisis potensi dan efektivitas biaya opsi mitigasi gas rumah kaca pada sektor kehutanan Indonesia dengan menggunakan model COMAP', Program Pasca sarjana, Institut Pertanian Bogor (Unpublished).
- Giannini, A., A.W. Robertson and J.H. Qian, (2007) A role for tropical tropospheric temperature adjustment to ENSO in the seasonality of monsoonal Indonesia precipitation predictability, *J. Geophys. Res. (Atmosphere)*, 112,
- Guilyardi E (2005) El Niño-mean state-seasonal cycle interactions in a multi-model ensemble. *Clim Dyn* DOI 10.1007/s00382-005- 0084-6
- Hansen, J., Sato, M., Ruedy, R., Lo, K., Lea, D.W., and Medina-Elizade, M. 2006. Global temperature change. *Proceeding of National Academy of Science* 103: 14288-14293.
- Hariati, F. B.D. Dasanto, R. Boer, A. Herienschah, Fitriyani and D. Octavariani. Marine and Coastal. In MoE. Technical Report on Vulnerability and Adaptation Assessment to Climate Change for Indonesia's Second National Communication. Ministry of Environment and United Nations Development Programme, Jakarta.
- Hasan, H., Pawitan, H., Boer, B dan Yahya, S. 1998. Model simulasi produksi kelapa sawit berdasarkan karakteristik kekeringan (*A simulation model of oil palm production based on drought characteristics*). *Jurnal Agromet* 13:41-54.
- Heil, A. B. Langmann and E. Aldrian. 2006. Indonesian peat and vegetation fire emissions: Study on factors influencing large-scale smoke haze pollution using a regional atmospheric chemistry model. *Mitig Adapt Strat Glob Change* 12:113-133
- Herienschah, A. R. Boer, Yanuar, and B.D. Dasanto. 2009. Water Resources. In MoE. Technical Report on Vulnerability and Adaptation Assessment to Climate Change for Indonesia's Second National Communication. Ministry of Environment and United Nations Development Programme, Jakarta.

- Hidayati, R., A. Buono, A. Rakhman, Fitriyani, R. Boer. 2009. Health Sector. Technical Report on Vulnerability and Adaptation Assessment to Climate Change for Indonesia's Second National Communication. Ministry of Environment and United Nations Development Programme, Jakarta.
- Hooijer, A., Silvius, M., Wösten, H., Page, S (2006): PEAT-CO<sub>2</sub>, Assessment of CO<sub>2</sub> emissions from drained peatlands in SE Asia. Delft Hydraulics report Q3943.
- IPCC. 2006. IPCC Guidelines for National Greenhouse Gas Inventories. Prepared by The National Greenhouse Gas Inventories Programme, Eggleston H.S., Buendia, L., Miwa, K., Ngara, T. and Tanabe, K. (eds.). Published by IGES Japan.
- IPCC. 2007. Climate Change 2007: The Physical Science Basis. Summary for Policy makers. Intergovernmental Panel on Climate Change, Geneva.
- IPCC. 1997. Revised 1996 IPCC Guidelines for National Greenhouse Gas Inventories – Workbook (Volume 2). <http://www.ipcc.ch>
- IPCC. 2000. Emission scenarios. A Special Report of Working Group III of the IPCC. Intergovernmental Panel on Climate Change. Cambridge University Press.
- Kalshoven, LGE. 1981. Pests of Crops in Indonesia. PT Ichtar Baru-van Hoeve. Jakarta.
- Kieft, J. & Soekarjo, D. 2007. Food and nutritional security assessment March 2007: Initial impact analysis of the 2006/2007 crop season in comparison to 1997/1998 and 2002/2003 El Nino events for the Eastern NTT region. Jakarta, CARE International Indonesia
- Kirono, D. and I. J. Partridge. 2002. The climate and the SOI. p. 17-24. In I.J. Partridge and M. Ma'shum (ed) Will It Rain?: The effect of the Southern Oscillation and El Niño in Indonesia. Queensland Government, Department of Primary Industry, Australia.
- Las, I and Team. 2008. Matrix Program and Strategy Roadmap for addressing climate change in agriculture sector. Agriculture Research and Development Agency Ministry of Agriculture. Report submitted to National Planning Agency (BAPPENAS).
- Las, I., Boer, R., Syahbudin, H. *et al.* 1999. Analisis peluang penyimpangan iklim dan ketersediaan air pada wilayah pengembangan IP Padi 300 (*Probability analysis of climate extreme in major rice growing areas*). Pusat Penelitian Tanah dan Agroklimat. Laporan Proyek ARMP-II, Badan Penelitian dan Pengembangan Pertanian. Bogor, 155p
- Levine JS. 1999. The 1997 fires in Kalimantan and Sumatra, Indonesia: gaseous and particulate emissions. *Geophysical Research Letters* 26:815–818.

- Masripatin, N., Ng. Ginting, A. Wibowo, and A. Pribadi. 2009 b. Forestry Sector. In Indonesia's Technology Needs Assessment on Climate Change Mitigation. Republic of Indonesia.
- Masripatin, N., Ng. Gintings and I.W.S. Dharmawan. 2009b. Penanganan perubahan iklim sektor kehutanan. Dalam Bappenas. Roadmap Pengarusutamaan Isu Perubahan Iklim dalam Perencanaan Pembangunan Nasional. Badan Perencanaan Pembangunan Nasional (Draft)
- MEMR. 2008, RUKN *Rencana Umum Ketenagalistrikan Nasional* (General National Electricity Plan) 2008 – 2027. Ministry of Energy and Mineral Resources, Jakarta
- MEMR. 2009, "*Blueprint Pengelolaan Energi Nasional (Blueprint of National Energy Management) 2010 - 2025*", Ministry of Energy and Mineral Resources, Jakarta.
- MoA. 2007. Strategy and technology innovation to cope with global climate change. Ministry of Agriculture Republic of Indonesia, Bogor.
- MoE, 2009. Technical Report: National Greenhouse Gas Inventory for the Second National Communication. Ministry of Environment and United Nation Development Program, Jakarta.
- MoE. 2003. Environmental status report. The State Ministry of Environment
- MoE. 2007. State of environment report in Indonesia 2006. The State Ministry of Environment
- MoE. 2007a. National Action Plan on Climate Change. Ministry of Environment, Republic of Indonesia
- MoE. 2007b. Climate variability and climate changes, and their implication. Ministry of Environment, Republic of Indonesia
- MoF, 2009. Rencana Strategis (Strategic Plan) 2010 – 2014. Ministry of Forestry, Jakarta
- MoF. 2006. Forestry Long Term Development Plan Year 2006-2025. Ministry of Forestry. Jakarta.
- MoF. 2007. Forestry Statistics of Indonesia. Ministry of Forestry. Jakarta
- MoF. 2008. National Forestry Plan (RKTN) Year 2010-2029. Ministry of Forestry. Jakarta.
- Moji K., Tsuyuoka, R., Makunike, B. and Tanaka, H. 2002. Meteorological factors influencing the intensity of malaria outbreak in Zimbabwe Japanese Journal of Tropical Medicine and Hygiene 30:283-287
- Wiyono, S. 2007. Climate change and pests and diseases explosion. Paper presented in One day Seminar on Biodiversity in the middle of global warming. KEHATI Foundation, Jakarta 28 June 2007 (in Indonesian)

- Moron, V., A.W. Robertson and R. Boer. 2009. Spatial coherence and seasonal predictability of monsoon onset over Indonesia. *Journal of Climate*.
- Nastari Bogor dan Klinik Tanaman IPB. 2007. Laporan Safari Gotong Royong *Sambung Keperluan untuk Petani Indonesia* di 24 Kabupaten-Kota di Pulau Jawa 4 April-2 Mei 2007. Yayasan Nastari Bogor- Klinik Tanaman IPB.
- Naylor, R.L., Battisti, D.S. Vimont, D.J., Falcon, W.P. and Burke, M.B. 2007. Assessing risks of climate variability and climate change for Indonesian rice agriculture. *Proceeding of the National Academic of Science* 114:7752-7757
- Page, S.E., F. Siegert, J. O. Rieley, Hans-Dieter V. Boehm<sup>3</sup>, A. Jaya & S. Limin. 2002. The amount of carbon released from peat and forest fires in Indonesia during 1997. *Nature* 420: 61-65
- Parry, M. L., A. R. Magalhaes, and N. H. Nih. 1992. *The potential socio-economic effects of climate change: A summary of three regional assessments*. Nairobi, Kenya: United Nations Environment Programme (UNEP).
- PEACE, World Bank and DFID, 2007. *Indonesia and Climate Change: Current Status and Policies*. The World Bank, Jakarta
- PT PLN, 2009. RUPTL *Rencana Usaha Penyediaan Tenaga Listrik* (Electricity Service Supply Plan) 2009 – 2018, Jakarta, Indonesia
- Pusdatin – MEMR (2006). *Handbook of Energy and Economic Statistics of Indonesia, 2006*. Jakarta, Indonesia
- Pusdatin – MEMR (2009a). *Handbook of Energy and Economic Statistics of Indonesia, 2009*. Jakarta, Indonesia
- Pusdatin – MEMR (2009b). *Indonesian Energy Outlook 2009*. Jakarta, Indonesia
- Quinn, W.H., Zopf, D.D., Sort, K.S., and Kuo Yang, T.R.W. 1978. Historical trends and statistics of the southern oscillation-El-Niño and Indonesian drought. *Fishery Bulletin* 76, p.3.
- Rodo, X. Pascual, M., Fuchs.G., Faruque, A.S.G. 2002. ENSO and cholera: A nonstationary link related to climate change?. *Proceeding of National Academy of Science* 99: 12901–12906
- Rohman, L. and R. Boer, 2001. Penggunaan indeks kenyamanan untuk mengevaluasi kesesuaian wilayah untuk proses reproduksi ternak domba (The use of comfort index to evaluate the suitability of region for reproduction process of sheep. *Jurnal Agromet*.

- Saddler, GS. 2000. IMI Description of Fungi and Bacteria No. 146 sheet 1457.
- Salinger, M.J. 2005. Increasing climate variability and change: reducing the vulnerability. *Climate Change* 70:1-3.
- Schimel, D., and Baker, D. 2002. The wildfire factor. *Nature* 420:29-30
- Setyanto, P., E. Surmaini and R. Boer. 2009. Mitigation of Methane Emission from Rice Field. In MoE. Projection of GHG Emission and Mitigation Analysis for Indonesia's Second National Communication. Ministry of Environment and United Nation Development Program, Jakarta.
- Siregar, H. 2008. Problems facing the Indonesian economy and current policy responses for agriculture. In Proceeding on the 33rd Annual Conference of The Federation of ASEAN's Economic Associations: "ASEAN's Co-operation and Agricultural and Rural Development in the Globalization Era. Hanoi, Vietnam.
- Sivakumar, M.V.K. 2005. Impacts of natural disasters in Agriculture, rangeland and forestry: An overview. In M.V.K. Sivakumar, R.P. Motha and H.P. Das (eds). *Natural disasters and extreme events in Agriculture*, page:1-22. Springer, Berlin.
- Sofian, I. 2010. Scientific Basis: Analysis and Projection of Sea Level Rise and Extreme Weather Event. Indonesia Climate Change Sectoral Roadmap – ICCSR. National Planning and Development Agency (Bappenas), Jakarta.
- Surmaini, E., I. Amien, E. Susanti, M.W. Trinugroho, Y. Sarvina. 2010. Pengembangan dan pemanfaatan sistem informasi prediksi curah hujan mk 2010 dan mh 2010/2011 berbasis web. Balai Penelitian Agroklimat Dan Hidrologi, Balai Besar Litbang Sumberdaya Lahan Pertanian, Badan Penelitian dan Pengembangan Pertanian, Departemen Pertanian.
- Suryahadi, and I. Permana. 2009. Mitigation of Methane Emission from Livestock. In MoE. Projection of GHG Emission and Mitigation Analysis for Indonesia's Second National Communication. Ministry of Environment and United Nation Development Program, Jakarta.
- Suryahadi, S., Wulandari, S., and Hidayat, R. (2001). Identification of less GHG emission technology on agriculture energy sector (livestock sub-sector). In MoE, *The identification of less GHG emission technologies*". State Ministry for Environment of Indonesia, Jakarta.
- Timmermann A, Oberhuber J, Bacher A, Esch M, Latif M, Roeckner E (1999) Increased El Niño frequency in a climate model forced by future greenhouse warming. *Nature* 398:694–696

- Tjasyono, B. 1997. Mekanisme fisis para, selama, dan pasca El Niño. Paper disajikan pada Workshop Kelompok Peneliti Dinamika Atmosfer, 13-14 Maret 1997.
- Tondok, E. 2001. The Causal Agent of Twisting Disease of Shallot. Master Thesis. University of Goettingen, Germany
- Trenberth, KE and Houghton, JT, 1996. *The 1990-1995 El Nino-Southern Oscillation Event: Longest on Record*. Geographical Research Letter 23: 57-60.
- UNDP CO. 2007. Strengthening national capacity to address climate change adaptation. United Nations Development Programme Country Office, Jakarta.
- USDA. 1984. World indices of agriculture and food production, 1974-1983. Statistical Bulletin 710. Economic Research Service, U.S. Department of Agriculture, Washington DC.
- Van der Werf, G. R., Randerson, J. T., Giglio, L., Gobron, N., Dolman, A. J. (2008). Climate controls on the variability of fires in the tropics and subtropics. *Global Biogeochemical Cycles*, 22, GB3028, 1-13
- Wibowo, A. and R. Boer. 2009. Mitigation Analysis for Forestry Sector. In MoE. Projection of GHG Emission and Mitigation Analysis for Indonesia's Second National Communication. Ministry of Environment and United Nations Development Programme, Jakarta.
- Wiyono, S. 2007. Climate change and pests and diseases explosion. Paper presented in One day Seminar on Biodiversity in the middle of global warming. KEHATI Foundation, Jakarta 28 June 2007 (in Indonesian)
- World Bank, 2008. Climate Change in Indonesia: Low Carbon Development Options Study. Phase 1 Status Report. Paper presented at the National Consultation on 'A Regional Review of the Economic of Climate Change in South East Asia (PRECCS). Jakarta 23-24 May 2008.
- Yoshino, M., K. Urushibara-Yoshino, and W. Suratman. 2000. Agriculture production and climate change in Indonesia. *Global Environ. Res.* 3:187-197.